

Alabama, 2013

Primary and secondary syphilis

- CASES: Alabama reported 183 cases in 2013.
- RATES: Alabama ranked 23 among 50 states reporting at least one case of P&S syphilis with 3.8 cases per 100,000 population compared to the U.S. rate of 5.5.

Rates of primary and secondary syphilis, by sex

- In Alabama, the rate among males was 6.9 cases per 100,000 population compared to the U.S. male rate of 10.3.
- The rate among females was 0.9 compared to the U.S. female rate of 0.9.


Rates of primary and secondary syphilis, by race/ethnicity

- In Alabama, the race/ethnicity adjusted rates per 100,000 population were 1.3 among whites, 9.1 among blacks, 1.0 among Hispanics, 0.0 among Asians, 0.0 among Native Hawaiian/Other Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 6.8 times that of whites.


Congenital syphilis

- CASES: Alabama reported 2 cases in 2013.
- RATES: Alabama ranked 20 among 25 states reporting at least one case of congenital syphilis with a rate of 3.3 cases per 100,000 live births compared to the U.S. rate of 8.7.

Syphilis: P&S Rates and Congenital Cases, Alabama, 2004–2013



Total P&S Syphilis Rates by County, Alabama, 2013



Rate (per 100,000 population)

≤0.4

0.41 - 2.2

>2.2

Alaska, 2013

Primary and secondary syphilis

- CASES: Alaska reported 23 cases in 2013.
- RATES: Alaska ranked 32 among 50 states reporting at least one case of P&S syphilis with 3.1 cases per 100,000 population compared to the U.S. rate of 5.5.

Rates of primary and secondary syphilis, by sex

- In Alaska, the rate among males was 5.5 per 100,000 population compared to the U.S. male rate of 10.3 cases per 100,000.
- The rate among females was 0.6 compared to the U.S. female rate of 0.9.


Rates of primary and secondary syphilis, by race/ethnicity*

- In Alaska, the race/ethnicity adjusted rates per 100,000 population were 1.7 among whites, 9.4 among blacks, 2.2 among Hispanics, 3.8 among Asian/Pacific Islanders, and 7.7 among American Indians/Alaska Natives.
- The rate among blacks was 5.5 times that of whites.
- The rate among Hispanics was 1.3 times that of whites.
- The rate among Asian/Pacific Islanders was 2.2 times that of whites.
- The rate among American Indians/Alaska Natives was 4.5 times that of whites.


Congenital syphilis

- CASES: Alaska reported 1 case in 2013.
- RATES: Alaska ranked 12 among 25 states reporting at least one case of congenital syphilis with a rate of 8.7 cases per 100,000 live births compared to the U.S. rate of 8.7.

Syphilis: P&S Rates and Congenital Cases, Alaska, 2004–2013



Total P&S Syphilis Rates by County, Alaska, 2013



Rate (per 100,000 population)

≤ 0.4
 0.41 - 2.2
 > 2.2

*OMB-compliant race values not available. NCHS bridged population used to calculate rates.

Arizona, 2013

Primary and secondary syphilis

- CASES: Arizona reported 287 cases in 2013.
- RATES: Arizona ranked 17 among 50 states reporting at least one case of P&S syphilis with 4.4 cases per 100,000 population compared to the U.S. rate of 5.5.

Rates of primary and secondary syphilis, by sex

- In Arizona, the rate among males was 8.0 cases per 100,000 population compared to the U.S. male rate of 10.3.
- The rate among females was 0.8 compared to the U.S. female rate of 0.9.


Rates of primary and secondary syphilis, by race/ethnicity

- In Arizona, the race/ethnicity adjusted rates per 100,000 population were 3.4 among whites, 13.7 among blacks, 5.1 among Hispanics, 2.1 among Asians, 0.0 among Native Hawaiian/Other Pacific Islanders, and 6.0 among American Indians/Alaska Natives.
- The rate among blacks was 4 times that of whites.
- The rate among Hispanics was 1.5 times that of whites.
- The rate among American Indians/Alaska Natives was 1.8 times that of whites.


Congenital syphilis

- CASES: Arizona reported 13 cases in 2013.
- RATES: Arizona ranked 7 among 25 states reporting at least one case of congenital syphilis with a rate of 14.9 cases per 100,000 live births compared to the U.S. rate of 8.7.

Syphilis: P&S Rates and Congenital Cases, Arizona, 2004–2013



Total P&S Syphilis Rates by County, Arizona, 2013



Rate (per 100,000 population)

≤0.4

0.41 - 2.2

>2.2

Arkansas, 2013

Primary and secondary syphilis

- CASES: Arkansas reported 177 cases in 2013.
- RATES: Arkansas ranked 10 among 50 states reporting at least one case of P&S syphilis with 6.0 cases per 100,000 population compared to the U.S. rate of 5.5.

Rates of primary and secondary syphilis, by sex

- In Arkansas, the rate among males was 9.2 cases per 100,000 population compared to the U.S. male rate of 10.3.
- The rate among females was 2.9 compared to the U.S. female rate of 0.9.


Rates of primary and secondary syphilis, by race/ethnicity

- In Arkansas, the race/ethnicity adjusted rates per 100,000 population were 2.8 among whites, 22.1 among blacks, 2.5 among Hispanics, 5.0 among Asians, 62.3 among Native Hawaiian/Other Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 7.9 times that of whites.
- The rate among Asians was 1.8 times that of whites.
- The rate among Native Hawaiian/Other Pacific Islanders was 22.3 times that of whites.


Congenital syphilis

- CASES: Arkansas reported 12 cases in 2013.
- RATES: Arkansas ranked 2 among 25 states reporting at least one case of congenital syphilis with a rate of 31.1 cases per 100,000 live births compared to the U.S. rate of 8.7.

Syphilis: P&S Rates and Congenital Cases, Arkansas, 2004–2013



Total P&S Syphilis Rates by County, Arkansas, 2013



Rate (per 100,000 population)

≤0.4

0.41 - 2.2

>2.2

California, 2013

Primary and secondary syphilis

- CASES: California reported 3,532 cases in 2013.
- RATES: California ranked 2 among 50 states reporting at least one case of P&S syphilis with 9.3 cases per 100,000 population compared to the U.S. rate of 5.5.

Rates of primary and secondary syphilis, by sex

- In California, the rate among males was 17.5 cases per 100,000 population compared to the U.S. male rate of 10.3.
- The rate among females was 1.1 compared to the U.S. female rate of 0.9.


Rates of primary and secondary syphilis, by race/ethnicity

- In California, the race/ethnicity adjusted rates per 100,000 population were 8.4 among whites, 22.5 among blacks, 8.7 among Hispanics, 3.9 among Asians, 13.1 among Native Hawaiian/Other Pacific Islanders, and 10.3 among American Indians/Alaska Natives.
- The rate among blacks was 2.7 times that of whites.
- The rate among Native Hawaiian/Other Pacific Islanders was 1.6 times that of whites.
- The rate among American Indians/Alaska Natives was 1.2 times that of whites.


Congenital syphilis

- CASES: California reported 56 cases in 2013.
- RATES: California ranked 10 among 25 states reporting at least one case of congenital syphilis with a rate of 11.0 cases per 100,000 live births compared to the U.S. rate of 8.7.

Syphilis: P&S Rates and Congenital Cases, California, 2004–2013



Total P&S Syphilis Rates by County, California, 2013



Rate (per 100,000 population)

≤0.4

0.41 - 2.2

>2.2

Colorado, 2013

Primary and secondary syphilis

- CASES: Colorado reported 163 cases in 2013.
- RATES: Colorado ranked 33 among 50 states reporting at least one case of P&S syphilis with 3.1 cases per 100,000 population compared to the U.S. rate of 5.5.

Rates of primary and secondary syphilis, by sex

- In Colorado, the rate among males was 6.1 cases per 100,000 population compared to the U.S. male rate of 10.3.
- The rate among females was 0.2 compared to the U.S. female rate of 0.9.


Rates of primary and secondary syphilis, by race/ethnicity

- In Colorado, the race/ethnicity adjusted rates per 100,000 population were 2.6 among whites, 5.0 among blacks, 3.6 among Hispanics, 1.4 among Asians, 15.8 among Native Hawaiian/Other Pacific Islanders, and 3.0 among American Indians/Alaska Natives.
- The rate among blacks was 1.9 times that of whites.
- The rate among Hispanics was 1.4 times that of whites.
- The rate among Native Hawaiian/Other Pacific Islanders was 6.1 times that of whites.
- The rate among American Indians/Alaska Natives was 1.2 times that of whites.


Congenital syphilis

- CASES: Colorado reported 0 cases in 2013.
- RATES: Colorado had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.7.

Syphilis: P&S Rates and Congenital Cases, Colorado, 2004–2013



Total P&S Syphilis Rates by County, Colorado, 2013



Rate (per 100,000 population)



Connecticut, 2013

Primary and secondary syphilis

- CASES: Connecticut reported 56 cases in 2013.
- RATES: Connecticut ranked 44 among 50 states reporting at least one case of P&S syphilis with 1.6 cases per 100,000 population compared to the U.S. rate of 5.5.

Rates of primary and secondary syphilis, by sex

- In Connecticut, the rate among males was 2.7 cases per 100,000 population compared to the U.S. male rate of 10.3.
- The rate among females was 0.4 compared to the U.S. female rate of 0.9.


Rates of primary and secondary syphilis, by race/ethnicity

- In Connecticut, the race/ethnicity adjusted rates per 100,000 population were 0.7 among whites, 5.5 among blacks, 2.3 among Hispanics, 0.0 among Asians, 0.0 among Native Hawaiian/Other Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 8.2 times that of whites.
- The rate among Hispanics was 3.5 times that of whites.


Congenital syphilis

- CASES: Connecticut reported 0 cases in 2013.
- RATES: Connecticut had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.7.

Syphilis: P&S Rates and Congenital Cases, Connecticut, 2004–2013



Total P&S Syphilis Rates by County, Connecticut, 2013



Rate (per 100,000 population)

<=0.4

0.41 - 2.2

>2.2

Delaware, 2013

Primary and secondary syphilis

- CASES: Delaware reported 52 cases in 2013.
- RATES: Delaware ranked 12 among 50 states reporting at least one case of P&S syphilis with 5.7 cases per 100,000 population compared to the U.S. rate of 5.5.

Rates of primary and secondary syphilis, by sex

- In Delaware, the rate among males was 11.0 cases per 100,000 population compared to the U.S. male rate of 10.3.
- The rate among females was 0.6 compared to the U.S. female rate of 0.9.


Rates of primary and secondary syphilis, by race/ethnicity

- In Delaware, the race/ethnicity adjusted rates per 100,000 population were 2.0 among whites, 17.6 among blacks, 7.6 among Hispanics, 0.0 among Asians, 0.0 among Native Hawaiian/Other Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 8.7 times that of whites.
- The rate among Hispanics was 3.8 times that of whites.


Congenital syphilis

- CASES: Delaware reported 1 case in 2013.
- RATES: Delaware ranked 11 among 25 states reporting at least one case of congenital syphilis with a rate of 8.8 cases per 100,000 live births compared to the U.S. rate of 8.7.

Syphilis: P&S Rates and Congenital Cases, Delaware, 2004–2013



Total P&S Syphilis Rates by County, Delaware, 2013



Rate (per 100,000 population)

<=0.4

0.41 - 2.2

>2.2

District of Columbia, 2013

Primary and secondary syphilis

- CASES: District of Columbia reported 168 cases in 2013.
- RATES: District of Columbia reported 26.6 cases per 100,000 population compared to the U.S. rate of 5.5.

Rates of primary and secondary syphilis, by sex

- In District of Columbia, the rate among males was 49.8 cases per 100,000 population compared to the U.S. male rate of 10.3.
- The rate among females was 5.7 compared to the U.S. female rate of 0.9.


Rates of primary and secondary syphilis, by race/ethnicity

- In District of Columbia, the race/ethnicity adjusted rates per 100,000 population were 18.7 among whites, 33.9 among blacks, 9.6 among Hispanics, 12.9 among Asians, 266 among Native Hawaiian/Other Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 1.8 times that of whites.
- The rate among Native Hawaiian/Other Pacific Islanders was 14.2 times that of whites.


Congenital syphilis

- CASES: District of Columbia reported 2 cases in 2013.
- RATES: District of Columbia had a rate of 21.8 cases per 100,000 live births compared to the U.S. rate of 8.7.

Syphilis: P&S Rates and Congenital Cases, District of Columbia, 2004–2013



Total P&S Syphilis Rate, District of Columbia, 2013



Rate (per 100,000 population)

<=0.4

0.41 - 2.2

>2.2

Florida, 2013

Primary and secondary syphilis

- CASES: Florida reported 1,513 cases in 2013.
- RATES: Florida ranked 4 among 50 states reporting at least one case of P&S syphilis with 7.8 cases per 100,000 population compared to the U.S. rate of 5.5.

Rates of primary and secondary syphilis, by sex

- In Florida, the rate among males was 14.6 cases per 100,000 population compared to the U.S. male rate of 10.3.
- The rate among females was 1.4 compared to the U.S. female rate of 0.9.


Rates of primary and secondary syphilis, by race/ethnicity

- In Florida, the race/ethnicity adjusted rates per 100,000 population were 4.3 among whites, 18.5 among blacks, 8.4 among Hispanics, 2.2 among Asians, 8.3 among Native Hawaiian/Other Pacific Islanders, and 4.0 among American Indians/Alaska Natives.
- The rate among blacks was 4.4 times that of whites.
- The rate among Hispanics was 2 times that of whites.
- The rate among Native Hawaiian/Other Pacific Islanders was 2 times that of whites.


Congenital syphilis

- CASES: Florida reported 37 cases in 2013.
- RATES: Florida ranked 5 among 25 states reporting at least one case of congenital syphilis with a rate of 17.2 cases per 100,000 live births compared to the U.S. rate of 8.7.

Syphilis: P&S Rates and Congenital Cases, Florida, 2004–2013



Total P&S Syphilis Rates by County, Florida, 2013



Rate (per 100,000 population)

≤0.4

0.41 - 2.2

>2.2

Georgia, 2013

Primary and secondary syphilis

- CASES: Georgia reported 1,017 cases in 2013.
- RATES: Georgia ranked 1 among 50 states reporting at least one case of P&S syphilis with 10.3 cases per 100,000 population compared to the U.S. rate of 5.5.

Rates of primary and secondary syphilis, by sex

- In Georgia, the rate among males was 19.2 cases per 100,000 population compared to the U.S. male rate of 10.3.
- The rate among females was 1.7 compared to the U.S. female rate of 0.9.


Rates of primary and secondary syphilis, by race/ethnicity

- In Georgia, the race/ethnicity adjusted rates per 100,000 population were 3.3 among whites, 25.5 among blacks, 5.1 among Hispanics, 0.9 among Asians, 34.1 among Native Hawaiian/Other Pacific Islanders, and 4.4 among American Indians/Alaska Natives.
- The rate among blacks was 7.8 times that of whites.
- The rate among Hispanics was 1.5 times that of whites.
- The rate among Native Hawaiian/Other Pacific Islanders was 10.4 times that of whites.
- The rate among American Indians/Alaska Natives was 1.4 times that of whites.


Congenital syphilis

- CASES: Georgia reported 20 cases in 2013.
- RATES: Georgia ranked 6 among 25 states reporting at least one case of congenital syphilis with a rate of 14.9 cases per 100,000 live births compared to the U.S. rate of 8.7.

Syphilis: P&S Rates and Congenital Cases, Georgia, 2004–2013



Total P&S Syphilis Rates by County, Georgia, 2013



Rate (per 100,000 population)

≤0.4

0.41 - 2.2

>2.2

Hawaii, 2013

Primary and secondary syphilis

- CASES: Hawaii reported 46 cases in 2013.
- RATES: Hawaii ranked 30 among 50 states reporting at least one case of P&S syphilis with 3.3 cases per 100,000 population compared to the U.S. rate of 5.5.

Rates of primary and secondary syphilis, by sex

- In Hawaii, the rate among males was 6.6 cases per 100,000 population compared to the U.S. male rate of 10.3.
- The rate among females was 0.0 compared to the U.S. female rate of 0.9.


Rates of primary and secondary syphilis, by race/ethnicity

- In Hawaii, the race/ethnicity adjusted rates per 100,000 population were 3.8 among whites, 7.5 among blacks, 3.0 among Hispanics, 2.5 among Asians, 3.8 among Native Hawaiian/Other Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 2 times that of whites.


Congenital syphilis

- CASES: Hawaii reported 0 cases in 2013.
- RATES: Hawaii had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.7.

Syphilis: P&S Rates and Congenital Cases, Hawaii, 2004–2013



Total P&S Syphilis Rates by County, Hawaii, 2013



Rate (per 100,000 population)

≤0.4

0.41 - 2.2

>2.2

Idaho, 2013

Primary and secondary syphilis

- CASES: Idaho reported 15 cases in 2013.
- RATES: Idaho ranked 45 among 50 states reporting at least one case of P&S syphilis with 0.9 cases per 100,000 population compared to the U.S. rate of 5.5.

Rates of primary and secondary syphilis, by sex

- In Idaho, the rate among males was 1.9 cases per 100,000 population compared to the U.S. male rate of 10.3.
- The rate among females was 0.0 compared to the U.S. female rate of 0.9.


Rates of primary and secondary syphilis, by race/ethnicity

- In Idaho, the race/ethnicity adjusted rates per 100,000 population were 0.8 among whites, 0.0 among blacks, 2.2 among Hispanics, 0.0 among Asians, 0.0 among Native Hawaiian/Other Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among Hispanics was 2.6 times that of whites.


Congenital syphilis

- CASES: Idaho reported 0 cases in 2013.
- RATES: Idaho had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.7.

Syphilis: P&S Rates and Congenital Cases, Idaho, 2004–2013



Total P&S Syphilis Rates by County, Idaho, 2013



Rate (per 100,000 population)

≤0.4
 0.41 - 2.2
 >2.2

Illinois, 2013

Primary and secondary syphilis

- CASES: Illinois reported 798 cases in 2013.
- RATES: Illinois ranked 9 among 50 states reporting at least one case of P&S syphilis with 6.2 cases per 100,000 population compared to the U.S. rate of 5.5.

Rates of primary and secondary syphilis, by sex

- In Illinois, the rate among males was 11.6 cases per 100,000 population compared to the U.S. male rate of 10.3.
- The rate among females was 1.0 compared to the U.S. female rate of 0.9.


Rates of primary and secondary syphilis, by race/ethnicity

- In Illinois, the race/ethnicity adjusted rates per 100,000 population were 3.0 among whites, 21.3 among blacks, 6.2 among Hispanics, 3.4 among Asians, 29.5 among Native Hawaiian/Other Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 7.2 times that of whites.
- The rate among Hispanics was 2.1 times that of whites.
- The rate among Asians was 1.1 times that of whites.
- The rate among Native Hawaiian/Other Pacific Islanders was 10 times that of whites.


Congenital syphilis

- CASES: Illinois reported 23 cases in 2013.
- RATES: Illinois ranked 8 among 25 states reporting at least one case of congenital syphilis with a rate of 13.9 cases per 100,000 live births compared to the U.S. rate of 8.7.

Syphilis: P&S Rates and Congenital Cases, Illinois, 2004–2013



Total P&S Syphilis Rates by County, Illinois, 2013



Rate (per 100,000 population)

≤0.4

0.41 - 2.2

>2.2

Indiana, 2013

Primary and secondary syphilis

- CASES: Indiana reported 215 cases in 2013.
- RATES: Indiana ranked 31 among 50 states reporting at least one case of P&S syphilis with 3.3 cases per 100,000 population compared to the U.S. rate of 5.5.

Rates of primary and secondary syphilis, by sex

- In Indiana, the rate among males was 6.1 cases per 100,000 population compared to the U.S. male rate of 10.3.
- The rate among females was 0.5 compared to the U.S. female rate of 0.9.


Rates of primary and secondary syphilis, by race/ethnicity

- In Indiana, the race/ethnicity adjusted rates per 100,000 population were 1.8 among whites, 14.9 among blacks, 5.6 among Hispanics, 1.8 among Asians, 0.0 among Native Hawaiian/Other Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 8.5 times that of whites.
- The rate among Hispanics was 3.2 times that of whites.


Congenital syphilis

- CASES: Indiana reported 0 cases in 2013.
- RATES: Indiana had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.7.

Syphilis: P&S Rates and Congenital Cases, Indiana, 2004–2013



Total P&S Syphilis Rates by County, Indiana, 2013



Rate (per 100,000 population)



Iowa, 2013

Primary and secondary syphilis

- CASES: Iowa reported 106 cases in 2013.
- RATES: Iowa ranked 28 among 50 states reporting at least one case of P&S syphilis with 3.4 cases per 100,000 population compared to the U.S. rate of 5.5.

Rates of primary and secondary syphilis, by sex

- In Iowa, the rate among males was 6.3 cases per 100,000 population compared to the U.S. male rate of 10.3.
- The rate among females was 0.6 compared to the U.S. female rate of 0.9.


Rates of primary and secondary syphilis, by race/ethnicity

- In Iowa, the race/ethnicity adjusted rates per 100,000 population were 2.7 among whites, 19.4 among blacks, 2.5 among Hispanics, 5.1 among Asians, 0.0 among Native Hawaiian/Other Pacific Islanders, and 11.4 among American Indians/Alaska Natives.
- The rate among blacks was 7.1 times that of whites.
- The rate among Asians was 1.9 times that of whites.
- The rate among American Indians/Alaska Natives was 4.2 times that of whites.


Congenital syphilis

- CASES: Iowa reported 0 cases in 2013.
- RATES: Iowa had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.7.

Syphilis: P&S Rates and Congenital Cases, Iowa, 2004–2013



Total P&S Syphilis Rates by County, Iowa, 2013



Rate (per 100,000 population)



Kansas, 2013

Primary and secondary syphilis

- CASES: Kansas reported 51 cases in 2013.
- RATES: Kansas ranked 41 among 50 states reporting at least one case of P&S syphilis with 1.8 cases per 100,000 population compared to the U.S. rate of 5.5.

Rates of primary and secondary syphilis, by sex

- In Kansas, the rate among males was 3.3 cases per 100,000 population compared to the U.S. male rate of 10.3.
- The rate among females was 0.3 compared to the U.S. female rate of 0.9.


Rates of primary and secondary syphilis, by race/ethnicity

- In Kansas, the race/ethnicity adjusted rates per 100,000 population were 1.4 among whites, 8.9 among blacks, 1.3 among Hispanics, 0.0 among Asians, 0.0 among Native Hawaiian/Other Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 6.2 times that of whites.


Congenital syphilis

- CASES: Kansas reported 0 cases in 2013.
- RATES: Kansas had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.7.

Syphilis: P&S Rates and Congenital Cases, Kansas, 2004–2013



Total P&S Syphilis Rates by County, Kansas, 2013



Rate (per 100,000 population)

≤0.4
 0.41 - 2.2
 >2.2

Kentucky, 2013

Primary and secondary syphilis

- CASES: Kentucky reported 122 cases in 2013.
- RATES: Kentucky ranked 35 among 50 states reporting at least one case of P&S syphilis with 2.8 cases per 100,000 population compared to the U.S. rate of 5.5.

Rates of primary and secondary syphilis, by sex

- In Kentucky, the rate among males was 4.9 cases per 100,000 population compared to the U.S. male rate of 10.3.
- The rate among females was 0.8 compared to the U.S. female rate of 0.9.


Rates of primary and secondary syphilis, by race/ethnicity

- In Kentucky, the race/ethnicity adjusted rates per 100,000 population were 1.6 among whites, 14.8 among blacks, 4.9 among Hispanics, 1.8 among Asians, 0.0 among Native Hawaiian/Other Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 9 times that of whites.
- The rate among Hispanics was 3 times that of whites.
- The rate among Asians was 1.1 times that of whites.


Congenital syphilis

- CASES: Kentucky reported 4 cases in 2013.
- RATES: Kentucky ranked 15 among 25 states reporting at least one case of congenital syphilis with a rate of 7.2 cases per 100,000 live births compared to the U.S. rate of 8.7.

Syphilis: P&S Rates and Congenital Cases, Kentucky, 2004–2013



Total P&S Syphilis Rates by County, Kentucky, 2013



Rate (per 100,000 population)

≤0.4

0.41 - 2.2

>2.2

Louisiana, 2013

Primary and secondary syphilis

- CASES: Louisiana reported 423 cases in 2013.
- RATES: Louisiana ranked 3 among 50 states reporting at least one case of P&S syphilis with 9.2 cases per 100,000 population compared to the U.S. rate of 5.5.

Rates of primary and secondary syphilis, by sex

- In Louisiana, the rate among males was 13.7 cases per 100,000 population compared to the U.S. male rate of 10.3.
- The rate among females was 4.9 compared to the U.S. female rate of 0.9.


Rates of primary and secondary syphilis, by race/ethnicity

- In Louisiana, the race/ethnicity adjusted rates per 100,000 population were 3.1 among whites, 22.3 among blacks, 3.4 among Hispanics, 1.3 among Asians, 0.0 among Native Hawaiian/Other Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 7.2 times that of whites.
- The rate among Hispanics was 1.1 times that of whites.


Congenital syphilis

- CASES: Louisiana reported 32 cases in 2013.
- RATES: Louisiana ranked 1 among 25 states reporting at least one case of congenital syphilis with a rate of 51.3 cases per 100,000 live births compared to the U.S. rate of 8.7.

Syphilis: P&S Rates and Congenital Cases, Louisiana, 2004–2013



Total P&S Syphilis Rates by County, Louisiana, 2013



Rate (per 100,000 population)

≤0.4

0.41 - 2.2

>2.2

Maine, 2013

Primary and secondary syphilis

- CASES: Maine reported 10 cases in 2013.
- RATES: Maine ranked 47 among 50 states reporting at least one case of P&S syphilis with 0.8 cases per 100,000 population compared to the U.S. rate of 5.5.

Rates of primary and secondary syphilis, by sex

- In Maine, the rate among males was 1.4 cases per 100,000 population compared to the U.S. male rate of 10.3.
- The rate among females was 0.1 compared to the U.S. female rate of 0.9.


Rates of primary and secondary syphilis, by race/ethnicity

- In Maine, the race/ethnicity adjusted rates per 100,000 population were 0.6 among whites, 6.0 among blacks, 0.0 among Hispanics, 0.0 among Asians, 0.0 among Native Hawaiian/Other Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 10.8 times that of whites.


Congenital syphilis

- CASES: Maine reported 0 cases in 2013.
- RATES: Maine had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.7.

Syphilis: P&S Rates and Congenital Cases, Maine, 2004–2013



Total P&S Syphilis Rates by County, Maine, 2013



Rate (per 100,000 population)

≤0.4

0.41 - 2.2

>2.2

Maryland, 2013

Primary and secondary syphilis

- CASES: Maryland reported 456 cases in 2013.
- RATES: Maryland ranked 5 among 50 states reporting at least one case of P&S syphilis with 7.7 cases per 100,000 population compared to the U.S. rate of 5.5.

Rates of primary and secondary syphilis, by sex

- In Maryland, the rate among males was 13.9 per 100,000 population compared to the U.S. male rate of 10.3 cases per 100,000.
- The rate among females was 2.0 compared to the U.S. female rate of 0.9.


Rates of primary and secondary syphilis, by race/ethnicity*

- In Maryland, the race/ethnicity adjusted rates per 100,000 population were 2.1 among whites, 19.6 among blacks, 5.1 among Hispanics, 2.4 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 9.3 times that of whites.
- The rate among Hispanics was 2.4 times that of whites.
- The rate among Asian/Pacific Islanders was 1.1 times that of whites.


Congenital syphilis

- CASES: Maryland reported 14 cases in 2013.
- RATES: Maryland ranked 4 among 25 states reporting at least one case of congenital syphilis with a rate of 19.0 cases per 100,000 live births compared to the U.S. rate of 8.7.

Syphilis: P&S Rates and Congenital Cases, Maryland, 2004–2013



Total P&S Syphilis Rates by County, Maryland, 2013



Rate (per 100,000 population)

≤0.4

0.41 - 2.2

>2.2

*OMB-compliant race values not available. NCHS bridged population used to calculate rates.

Massachusetts, 2013

Primary and secondary syphilis

- CASES: Massachusetts reported 360 cases in 2013.
- RATES: Massachusetts ranked 14 among 50 states reporting at least one case of P&S syphilis with 5.4 cases per 100,000 population compared to the U.S. rate of 5.5.

Rates of primary and secondary syphilis, by sex

- In Massachusetts, the rate among males was 10.7 cases per 100,000 population compared to the U.S. male rate of 10.3.
- The rate among females was 0.5 compared to the U.S. female rate of 0.9.


Rates of primary and secondary syphilis, by race/ethnicity

- In Massachusetts, the race/ethnicity adjusted rates per 100,000 population were 4.0 among whites, 11.6 among blacks, 10.1 among Hispanics, 2.4 among Asians, 0.0 among Native Hawaiian/Other Pacific Islanders, and 33.6 among American Indians/Alaska Natives.
- The rate among blacks was 2.9 times that of whites.
- The rate among Hispanics was 2.5 times that of whites.
- The rate among American Indians/Alaska Natives was 8.4 times that of whites.


Congenital syphilis

- CASES: Massachusetts reported 4 cases in 2013.
- RATES: Massachusetts ranked 17 among 25 states reporting at least one case of congenital syphilis with a rate of 5.5 cases per 100,000 live births compared to the U.S. rate of 8.7.

Syphilis: P&S Rates and Congenital Cases, Massachusetts, 2004–2013



Total P&S Syphilis Rates by County, Massachusetts, 2013



Rate (per 100,000 population)

≤0.4

0.41 - 2.2

>2.2

Michigan, 2013

Primary and secondary syphilis

- CASES: Michigan reported 487 cases in 2013.
- RATES: Michigan ranked 16 among 50 states reporting at least one case of P&S syphilis with 4.9 cases per 100,000 population compared to the U.S. rate of 5.5.

Rates of primary and secondary syphilis, by sex

- In Michigan, the rate among males was 9.4 cases per 100,000 population compared to the U.S. male rate of 10.3.
- The rate among females was 0.6 compared to the U.S. female rate of 0.9.


Rates of primary and secondary syphilis, by race/ethnicity

- In Michigan, the race/ethnicity adjusted rates per 100,000 population were 1.7 among whites, 24.0 among blacks, 3.5 among Hispanics, 0.8 among Asians, 0.0 among Native Hawaiian/Other Pacific Islanders, and 1.8 among American Indians/Alaska Natives.
- The rate among blacks was 13.9 times that of whites.
- The rate among Hispanics was 2 times that of whites.


Congenital syphilis

- CASES: Michigan reported 9 cases in 2013.
- RATES: Michigan ranked 13 among 25 states reporting at least one case of congenital syphilis with a rate of 7.9 cases per 100,000 live births compared to the U.S. rate of 8.7.

Syphilis: P&S Rates and Congenital Cases, Michigan, 2004–2013



Total P&S Syphilis Rates by County, Michigan, 2013



Rate (per 100,000 population)

≤0.4

0.41 - 2.2

>2.2

Minnesota, 2013

Primary and secondary syphilis

- CASES: Minnesota reported 193 cases in 2013.
- RATES: Minnesota ranked 27 among 50 states reporting at least one case of P&S syphilis with 3.6 cases per 100,000 population compared to the U.S. rate of 5.5.

Rates of primary and secondary syphilis, by sex

- In Minnesota, the rate among males was 6.7 cases per 100,000 population compared to the U.S. male rate of 10.3.
- The rate among females was 0.4 compared to the U.S. female rate of 0.9.


Rates of primary and secondary syphilis, by race/ethnicity

- In Minnesota, the race/ethnicity adjusted rates per 100,000 population were 2.5 among whites, 16.4 among blacks, 3.4 among Hispanics, 3.0 among Asians, 46.7 among Native Hawaiian/Other Pacific Islanders, and 3.5 among American Indians/Alaska Natives.
- The rate among blacks was 6.6 times that of whites.
- The rate among Hispanics was 1.4 times that of whites.
- The rate among Asians was 1.2 times that of whites.
- The rate among Native Hawaiian/Other Pacific Islanders was 19 times that of whites.
- The rate among American Indians/Alaska Natives was 1.4 times that of whites.


Congenital syphilis

- CASES: Minnesota reported 0 cases in 2013.
- RATES: Minnesota had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.7.

Syphilis: P&S Rates and Congenital Cases, Minnesota, 2004–2013



Total P&S Syphilis Rates by County, Minnesota, 2013



Rate (per 100,000 population)

≤0.4

0.41 - 2.2

>2.2

Mississippi, 2013

Primary and secondary syphilis

- CASES: Mississippi reported 78 cases in 2013.
- RATES: Mississippi ranked 37 among 50 states reporting at least one case of P&S syphilis with 2.6 cases per 100,000 population compared to the U.S. rate of 5.5.

Rates of primary and secondary syphilis, by sex

- In Mississippi, the rate among males was 4.1 cases per 100,000 population compared to the U.S. male rate of 10.3.
- The rate among females was 1.2 compared to the U.S. female rate of 0.9.


Rates of primary and secondary syphilis, by race/ethnicity

- In Mississippi, the race/ethnicity adjusted rates per 100,000 population were 0.1 among whites, 6.5 among blacks, 1.2 among Hispanics, 0.0 among Asians, 0.0 among Native Hawaiian/Other Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 55.9 times that of whites.
- The rate among Hispanics was 10.1 times that of whites.


Congenital syphilis

- CASES: Mississippi reported 0 cases in 2013.
- RATES: Mississippi had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.7.

Syphilis: P&S Rates and Congenital Cases, Mississippi, 2004–2013



Total P&S Syphilis Rates by County, Mississippi, 2013



Rate (per 100,000 population)

≤0.4

0.41 - 2.2

>2.2

Missouri, 2013

Primary and secondary syphilis

- CASES: Missouri reported 251 cases in 2013.
- RATES: Missouri ranked 19 among 50 states reporting at least one case of P&S syphilis with 4.2 cases per 100,000 population compared to the U.S. rate of 5.5.

Rates of primary and secondary syphilis, by sex

- In Missouri, the rate among males was 7.9 cases per 100,000 population compared to the U.S. male rate of 10.3.
- The rate among females was 0.6 compared to the U.S. female rate of 0.9.


Rates of primary and secondary syphilis, by race/ethnicity

- In Missouri, the race/ethnicity adjusted rates per 100,000 population were 2.0 among whites, 18.4 among blacks, 6.7 among Hispanics, 0.0 among Asians, 31.2 among Native Hawaiian/Other Pacific Islanders, and 8.1 among American Indians/Alaska Natives.
- The rate among blacks was 9.2 times that of whites.
- The rate among Hispanics was 3.3 times that of whites.
- The rate among Native Hawaiian/Other Pacific Islanders was 15.6 times that of whites.
- The rate among American Indians/Alaska Natives was 4 times that of whites.


Congenital syphilis

- CASES: Missouri reported 3 cases in 2013.
- RATES: Missouri ranked 19 among 25 states reporting at least one case of congenital syphilis with a rate of 3.9 cases per 100,000 live births compared to the U.S. rate of 8.7.

Syphilis: P&S Rates and Congenital Cases, Missouri, 2004–2013



Total P&S Syphilis Rates by County, Missouri, 2013



Rate (per 100,000 population)

≤0.4

0.41 - 2.2

>2.2

Montana, 2013

Primary and secondary syphilis

- CASES: Montana reported 5 cases in 2013.
- RATES: Montana ranked 48 among 50 states reporting at least one case of P&S syphilis with 0.5 cases per 100,000 population compared to the U.S. rate of 5.5.

Rates of primary and secondary syphilis, by sex

- In Montana, the rate among males was 0.8 cases per 100,000 population compared to the U.S. male rate of 10.3.
- The rate among females was 0.2 compared to the U.S. female rate of 0.9.


Rates of primary and secondary syphilis, by race/ethnicity

- In Montana, the race/ethnicity adjusted rates per 100,000 population were 0.3 among whites, 0.0 among blacks, 0.0 among Hispanics, 0.0 among Asians, 0.0 among Native Hawaiian/Other Pacific Islanders, and 1.6 among American Indians/Alaska Natives.
- The rate among American Indians/Alaska Natives was 4.8 times that of whites.


Congenital syphilis

- CASES: Montana reported 0 cases in 2013.
- RATES: Montana had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.7.

Syphilis: P&S Rates and Congenital Cases, Montana, 2004–2013



Total P&S Syphilis Rates by County, Montana, 2013



Rate (per 100,000 population)



Nebraska, 2013

Primary and secondary syphilis

- CASES: Nebraska reported 41 cases in 2013.
- RATES: Nebraska ranked 39 among 50 states reporting at least one case of P&S syphilis with 2.2 cases per 100,000 population compared to the U.S. rate of 5.5.

Rates of primary and secondary syphilis, by sex

- In Nebraska, the rate among males was 4.0 cases per 100,000 population compared to the U.S. male rate of 10.3.
- The rate among females was 0.4 compared to the U.S. female rate of 0.9.


Rates of primary and secondary syphilis, by race/ethnicity

- In Nebraska, the race/ethnicity adjusted rates per 100,000 population were 1.5 among whites, 13.1 among blacks, 0.6 among Hispanics, 2.8 among Asians, 0.0 among Native Hawaiian/Other Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 9 times that of whites.
- The rate among Asians was 1.9 times that of whites.


Congenital syphilis

- CASES: Nebraska reported 0 cases in 2013.
- RATES: Nebraska had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.7.

Syphilis: P&S Rates and Congenital Cases, Nebraska, 2004–2013



Total P&S Syphilis Rates by County, Nebraska, 2013



Rate (per 100,000 population)



Nevada, 2013

Primary and secondary syphilis

- CASES: Nevada reported 205 cases in 2013.
- RATES: Nevada ranked 7 among 50 states reporting at least one case of P&S syphilis with 7.4 cases per 100,000 population compared to the U.S. rate of 5.5.

Rates of primary and secondary syphilis, by sex

- In Nevada, the rate among males was 13.7 cases per 100,000 population compared to the U.S. male rate of 10.3.
- The rate among females was 1.0 compared to the U.S. female rate of 0.9.


Rates of primary and secondary syphilis, by race/ethnicity

- In Nevada, the race/ethnicity adjusted rates per 100,000 population were 5.3 among whites, 16.7 among blacks, 8.4 among Hispanics, 4.9 among Asians, 12.3 among Native Hawaiian/Other Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 3.2 times that of whites.
- The rate among Hispanics was 1.6 times that of whites.
- The rate among Native Hawaiian/Other Pacific Islanders was 2.3 times that of whites.


Congenital syphilis

- CASES: Nevada reported 2 cases in 2013.
- RATES: Nevada ranked 16 among 25 states reporting at least one case of congenital syphilis with a rate of 5.6 cases per 100,000 live births compared to the U.S. rate of 8.7.

Syphilis: P&S Rates and Congenital Cases, Nevada, 2004–2013



Total P&S Syphilis Rates by County, Nevada, 2013



Rate (per 100,000 population)

≤0.4

0.41 - 2.2

>2.2

New Hampshire, 2013

Primary and secondary syphilis

- CASES: New Hampshire reported 28 cases in 2013.
- RATES: New Hampshire ranked 40 among 50 states reporting at least one case of P&S syphilis with 2.1 cases per 100,000 population compared to the U.S. rate of 5.5.

Rates of primary and secondary syphilis, by sex

- In New Hampshire, the rate among males was 4.1 cases per 100,000 population compared to the U.S. male rate of 10.3.
- The rate among females was 0.1 compared to the U.S. female rate of 0.9.


Rates of primary and secondary syphilis, by race/ethnicity

- In New Hampshire, the race/ethnicity adjusted rates per 100,000 population were 1.9 among whites, 12.8 among blacks, 5.0 among Hispanics, 0.0 among Asians, 0.0 among Native Hawaiian/Other Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 6.7 times that of whites.
- The rate among Hispanics was 2.6 times that of whites.


Congenital syphilis

- CASES: New Hampshire reported 0 cases in 2013.
- RATES: New Hampshire had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.7.

Syphilis: P&S Rates and Congenital Cases, New Hampshire, 2004–2013



Total P&S Syphilis Rates by County, New Hampshire, 2013



Rate (per 100,000 population)

<=0.4

0.41 - 2.2

>2.2

New Jersey, 2013

Primary and secondary syphilis

- CASES: New Jersey reported 233 cases in 2013.
- RATES: New Jersey ranked 36 among 50 states reporting at least one case of P&S syphilis with 2.6 cases per 100,000 population compared to the U.S. rate of 5.5.

Rates of primary and secondary syphilis, by sex

- In New Jersey, the rate among males was 5.1 cases per 100,000 population compared to the U.S. male rate of 10.3.
- The rate among females was 0.3 compared to the U.S. female rate of 0.9.


Rates of primary and secondary syphilis, by race/ethnicity

- In New Jersey, the race/ethnicity adjusted rates per 100,000 population were 1.4 among whites, 7.8 among blacks, 4.0 among Hispanics, 0.4 among Asians, 0.0 among Native Hawaiian/Other Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 5.5 times that of whites.
- The rate among Hispanics was 2.8 times that of whites.


Congenital syphilis

- CASES: New Jersey reported 0 cases in 2013.
- RATES: New Jersey had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.7.

Syphilis: P&S Rates and Congenital Cases, New Jersey, 2004–2013



Total P&S Syphilis Rates by County, New Jersey, 2013



Rate (per 100,000 population)

≤0.4

0.41 - 2.2

>2.2

New Mexico, 2013

Primary and secondary syphilis

- CASES: New Mexico reported 78 cases in 2013.
- RATES: New Mexico ranked 25 among 50 states reporting at least one case of P&S syphilis with 3.7 cases per 100,000 population compared to the U.S. rate of 5.5.

Rates of primary and secondary syphilis, by sex

- In New Mexico, the rate among males was 5.6 cases per 100,000 population compared to the U.S. male rate of 10.3.
- The rate among females was 1.9 compared to the U.S. female rate of 0.9.


Rates of primary and secondary syphilis, by race/ethnicity

- In New Mexico, the race/ethnicity adjusted rates per 100,000 population were 2.0 among whites, 16.1 among blacks, 3.8 among Hispanics, 0.0 among Asians, 0.0 among Native Hawaiian/Other Pacific Islanders, and 3.4 among American Indians/Alaska Natives.
- The rate among blacks was 7.9 times that of whites.
- The rate among Hispanics was 1.8 times that of whites.
- The rate among American Indians/Alaska Natives was 1.6 times that of whites.


Congenital syphilis

- CASES: New Mexico reported 2 cases in 2013.
- RATES: New Mexico ranked 14 among 25 states reporting at least one case of congenital syphilis with a rate of 7.2 cases per 100,000 live births compared to the U.S. rate of 8.7.

Syphilis: P&S Rates and Congenital Cases, New Mexico, 2004–2013



Total P&S Syphilis Rates by County, New Mexico, 2013



Rate (per 100,000 population)



New York, 2013

Primary and secondary syphilis

- CASES: New York reported 1,459 cases in 2013.
- RATES: New York ranked 6 among 50 states reporting at least one case of P&S syphilis with 7.5 cases per 100,000 population compared to the U.S. rate of 5.5.

Rates of primary and secondary syphilis, by sex

- In New York, the rate among males was 14.8 per 100,000 population compared to the U.S. male rate of 10.3 cases per 100,000.
- The rate among females was 0.4 compared to the U.S. female rate of 0.9.


Rates of primary and secondary syphilis, by race/ethnicity*

- In New York, the race/ethnicity adjusted rates per 100,000 population were 3.7 among whites, 15.8 among blacks, 10.0 among Hispanics, 2.9 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 4.3 times that of whites.
- The rate among Hispanics was 2.7 times that of whites.


Congenital syphilis

- CASES: New York reported 11 cases in 2013.
- RATES: New York ranked 18 among 25 states reporting at least one case of congenital syphilis with a rate of 4.5 cases per 100,000 live births compared to the U.S. rate of 8.7.

Syphilis: P&S Rates and Congenital Cases, New York, 2004–2013



Total P&S Syphilis Rates by County, New York, 2013



Rate (per 100,000 population)

≤0.4

0.41 - 2.2

>2.2

*OMB-compliant race values not available. NCHS bridged population used to calculate rates.

North Carolina, 2013

Primary and secondary syphilis

- CASES: North Carolina reported 404 cases in 2013.
- RATES: North Carolina ranked 20 among 50 states reporting at least one case of P&S syphilis with 4.1 cases per 100,000 population compared to the U.S. rate of 5.5.

Rates of primary and secondary syphilis, by sex

- In North Carolina, the rate among males was 7.7 cases per 100,000 population compared to the U.S. male rate of 10.3.
- The rate among females was 0.7 compared to the U.S. female rate of 0.9.


Rates of primary and secondary syphilis, by race/ethnicity

- In North Carolina, the race/ethnicity adjusted rates per 100,000 population were 1.6 among whites, 13.5 among blacks, 1.3 among Hispanics, 0.0 among Asians, 0.0 among Native Hawaiian/Other Pacific Islanders, and 3.6 among American Indians/Alaska Natives.
- The rate among blacks was 8.5 times that of whites.
- The rate among American Indians/Alaska Natives was 2.3 times that of whites.


Congenital syphilis

- CASES: North Carolina reported 1 case in 2013.
- RATES: North Carolina ranked 25 among 25 states reporting at least one case of congenital syphilis with a rate of 0.8 cases per 100,000 live births compared to the U.S. rate of 8.7.

Syphilis: P&S Rates and Congenital Cases, North Carolina, 2004–2013



Total P&S Syphilis Rates by County, North Carolina, 2013



North Dakota, 2013

Primary and secondary syphilis

- CASES: North Dakota reported 12 cases in 2013.
- RATES: North Dakota ranked 42 among 50 states reporting at least one case of P&S syphilis with 1.7 cases per 100,000 population compared to the U.S. rate of 5.5.

Rates of primary and secondary syphilis, by sex

- In North Dakota, the rate among males was 3.1 cases per 100,000 population compared to the U.S. male rate of 10.3.
- The rate among females was 0.3 compared to the U.S. female rate of 0.9.


Rates of primary and secondary syphilis, by race/ethnicity

- In North Dakota, the race/ethnicity adjusted rates per 100,000 population were 0.3 among whites, 10.0 among blacks, 5.8 among Hispanics, 13.1 among Asians, 0.0 among Native Hawaiian/Other Pacific Islanders, and 16.4 among American Indians/Alaska Natives.
- The rate among blacks was 30.7 times that of whites.
- The rate among Hispanics was 17.9 times that of whites.
- The rate among Asians was 40.3 times that of whites.
- The rate among American Indians/Alaska Natives was 50.5 times that of whites.


Congenital syphilis

- CASES: North Dakota reported 0 cases in 2013.
- RATES: North Dakota had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.7.

Syphilis: P&S Rates and Congenital Cases, North Dakota, 2004–2013



Total P&S Syphilis Rates by County, North Dakota, 2013



Rate (per 100,000 population)

<=0.4

0.41 - 2.2

>2.2

Ohio, 2013

Primary and secondary syphilis

- CASES: Ohio reported 436 cases in 2013.
- RATES: Ohio ranked 24 among 50 states reporting at least one case of P&S syphilis with 3.8 cases per 100,000 population compared to the U.S. rate of 5.5.

Rates of primary and secondary syphilis, by sex

- In Ohio, the rate among males was 6.6 cases per 100,000 population compared to the U.S. male rate of 10.3.
- The rate among females was 1.1 compared to the U.S. female rate of 0.9.


Rates of primary and secondary syphilis, by race/ethnicity

- In Ohio, the race/ethnicity adjusted rates per 100,000 population were 2.1 among whites, 15.1 among blacks, 4.2 among Hispanics, 1.0 among Asians, 0.0 among Native Hawaiian/Other Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 7.3 times that of whites.
- The rate among Hispanics was 2 times that of whites.


Congenital syphilis

- CASES: Ohio reported 17 cases in 2013.
- RATES: Ohio ranked 9 among 25 states reporting at least one case of congenital syphilis with a rate of 12.2 cases per 100,000 live births compared to the U.S. rate of 8.7.

Syphilis: P&S Rates and Congenital Cases, Ohio, 2004–2013



Total P&S Syphilis Rates by County, Ohio, 2013



Oklahoma, 2013

Primary and secondary syphilis

- CASES: Oklahoma reported 118 cases in 2013.
- RATES: Oklahoma ranked 34 among 50 states reporting at least one case of P&S syphilis with 3.1 cases per 100,000 population compared to the U.S. rate of 5.5.

Rates of primary and secondary syphilis, by sex

- In Oklahoma, the rate among males was 5.6 cases per 100,000 population compared to the U.S. male rate of 10.3.
- The rate among females was 0.7 compared to the U.S. female rate of 0.9.


Rates of primary and secondary syphilis, by race/ethnicity

- In Oklahoma, the race/ethnicity adjusted rates per 100,000 population were 2.8 among whites, 9.7 among blacks, 2.5 among Hispanics, 2.9 among Asians, 0.0 among Native Hawaiian/Other Pacific Islanders, and 1.0 among American Indians/Alaska Natives.
- The rate among blacks was 3.4 times that of whites.


Congenital syphilis

- CASES: Oklahoma reported 0 cases in 2013.
- RATES: Oklahoma had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.7.

Syphilis: P&S Rates and Congenital Cases, Oklahoma, 2004–2013



Total P&S Syphilis Rates by County, Oklahoma, 2013



Oregon, 2013

Primary and secondary syphilis

- CASES: Oregon reported 267 cases in 2013.
- RATES: Oregon ranked 8 among 50 states reporting at least one case of P&S syphilis with 6.8 cases per 100,000 population compared to the U.S. rate of 5.5.

Rates of primary and secondary syphilis, by sex

- In Oregon, the rate among males was 13.2 cases per 100,000 population compared to the U.S. male rate of 10.3.
- The rate among females was 0.6 compared to the U.S. female rate of 0.9.


Rates of primary and secondary syphilis, by race/ethnicity

- In Oregon, the race/ethnicity adjusted rates per 100,000 population were 6.2 among whites, 20.6 among blacks, 10.8 among Hispanics, 2.6 among Asians, 0.0 among Native Hawaiian/Other Pacific Islanders, and 4.6 among American Indians/Alaska Natives.
- The rate among blacks was 3.3 times that of whites.
- The rate among Hispanics was 1.7 times that of whites.


Congenital syphilis

- CASES: Oregon reported 0 cases in 2013.
- RATES: Oregon had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.7.

Syphilis: P&S Rates and Congenital Cases, Oregon, 2004–2013



Total P&S Syphilis Rates by County, Oregon, 2013



Pennsylvania, 2013

Primary and secondary syphilis

- CASES: Pennsylvania reported 471 cases in 2013.
- RATES: Pennsylvania ranked 26 among 50 states reporting at least one case of P&S syphilis with 3.7 cases per 100,000 population compared to the U.S. rate of 5.5.

Rates of primary and secondary syphilis, by sex

- In Pennsylvania, the rate among males was 7.1 cases per 100,000 population compared to the U.S. male rate of 10.3.
- The rate among females was 0.4 compared to the U.S. female rate of 0.9.


Rates of primary and secondary syphilis, by race/ethnicity

- In Pennsylvania, the race/ethnicity adjusted rates per 100,000 population were 1.7 among whites, 16.7 among blacks, 5.0 among Hispanics, 0.8 among Asians, 31.4 among Native Hawaiian/Other Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 10 times that of whites.
- The rate among Hispanics was 3 times that of whites.
- The rate among Native Hawaiian/Other Pacific Islanders was 18.7 times that of whites.


Congenital syphilis

- CASES: Pennsylvania reported 2 cases in 2013.
- RATES: Pennsylvania ranked 24 among 25 states reporting at least one case of congenital syphilis with a rate of 1.4 cases per 100,000 live births compared to the U.S. rate of 8.7.

Syphilis: P&S Rates and Congenital Cases, Pennsylvania, 2004–2013



Total P&S Syphilis Rates by County, Pennsylvania, 2013



Rate (per 100,000 population)

<=0.4

0.41 - 2.2

>2.2

Rhode Island, 2013

Primary and secondary syphilis

- CASES: Rhode Island reported 45 cases in 2013.
- RATES: Rhode Island ranked 18 among 50 states reporting at least one case of P&S syphilis with 4.3 cases per 100,000 population compared to the U.S. rate of 5.5.

Rates of primary and secondary syphilis, by sex

- In Rhode Island, the rate among males was 8.7 cases per 100,000 population compared to the U.S. male rate of 10.3.
- The rate among females was 0.2 compared to the U.S. female rate of 0.9.


Rates of primary and secondary syphilis, by race/ethnicity

- In Rhode Island, the race/ethnicity adjusted rates per 100,000 population were 3.8 among whites, 7.0 among blacks, 4.3 among Hispanics, 9.2 among Asians, 0.0 among Native Hawaiian/Other Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 1.8 times that of whites.
- The rate among Hispanics was 1.2 times that of whites.
- The rate among Asians was 2.4 times that of whites.


Congenital syphilis

- CASES: Rhode Island reported 0 cases in 2013.
- RATES: Rhode Island had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.7.

Syphilis: P&S Rates and Congenital Cases, Rhode Island, 2004–2013



Total P&S Syphilis Rates by County, Rhode Island, 2013



Rate (per 100,000 population)



South Carolina, 2013

Primary and secondary syphilis

- CASES: South Carolina reported 271 cases in 2013.
- RATES: South Carolina ranked 11 among 50 states reporting at least one case of P&S syphilis with 5.7 cases per 100,000 population compared to the U.S. rate of 5.5.

Rates of primary and secondary syphilis, by sex

- In South Carolina, the rate among males was 10.1 cases per 100,000 population compared to the U.S. male rate of 10.3.
- The rate among females was 1.6 compared to the U.S. female rate of 0.9.


Rates of primary and secondary syphilis, by race/ethnicity

- In South Carolina, the race/ethnicity adjusted rates per 100,000 population were 2.0 among whites, 15.8 among blacks, 1.6 among Hispanics, 1.5 among Asians, 0.0 among Native Hawaiian/Other Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 7.8 times that of whites.


Congenital syphilis

- CASES: South Carolina reported 1 case in 2013.
- RATES: South Carolina ranked 23 among 25 states reporting at least one case of congenital syphilis with a rate of 1.7 cases per 100,000 live births compared to the U.S. rate of 8.7.

Syphilis: P&S Rates and Congenital Cases, South Carolina, 2004–2013



Total P&S Syphilis Rates by County, South Carolina, 2013



Rate (per 100,000 population)

≤0.4

0.41 - 2.2

>2.2

South Dakota, 2013

Primary and secondary syphilis

- CASES: South Dakota reported 44 cases in 2013.
- RATES: South Dakota ranked 15 among 50 states reporting at least one case of P&S syphilis with 5.3 cases per 100,000 population compared to the U.S. rate of 5.5.

Rates of primary and secondary syphilis, by sex

- In South Dakota, the rate among males was 6.9 cases per 100,000 population compared to the U.S. male rate of 10.3.
- The rate among females was 3.6 compared to the U.S. female rate of 0.9.


Rates of primary and secondary syphilis, by race/ethnicity

- In South Dakota, the race/ethnicity adjusted rates per 100,000 population were 2.4 among whites, 7.5 among blacks, 11.7 among Hispanics, 0.0 among Asians, 0.0 among Native Hawaiian/Other Pacific Islanders, and 31.1 among American Indians/Alaska Natives.
- The rate among blacks was 3.1 times that of whites.
- The rate among Hispanics was 4.8 times that of whites.
- The rate among American Indians/Alaska Natives was 12.8 times that of whites.


Congenital syphilis

- CASES: South Dakota reported 0 cases in 2013.
- RATES: South Dakota had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.7.

Syphilis: P&S Rates and Congenital Cases, South Dakota, 2004–2013



Total P&S Syphilis Rates by County, South Dakota, 2013



Rate (per 100,000 population)



Tennessee, 2013

Primary and secondary syphilis

- CASES: Tennessee reported 214 cases in 2013.
- RATES: Tennessee ranked 29 among 50 states reporting at least one case of P&S syphilis with 3.3 cases per 100,000 population compared to the U.S. rate of 5.5.

Rates of primary and secondary syphilis, by sex

- In Tennessee, the rate among males was 6.1 cases per 100,000 population compared to the U.S. male rate of 10.3.
- The rate among females was 0.7 compared to the U.S. female rate of 0.9.


Rates of primary and secondary syphilis, by race/ethnicity

- In Tennessee, the race/ethnicity adjusted rates per 100,000 population were 1.5 among whites, 12.3 among blacks, 1.3 among Hispanics, 1.0 among Asians, 0.0 among Native Hawaiian/Other Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 8.1 times that of whites.


Congenital syphilis

- CASES: Tennessee reported 2 cases in 2013.
- RATES: Tennessee ranked 21 among 25 states reporting at least one case of congenital syphilis with a rate of 2.5 cases per 100,000 live births compared to the U.S. rate of 8.7.

Syphilis: P&S Rates and Congenital Cases, Tennessee, 2004–2013



Total P&S Syphilis Rates by County, Tennessee, 2013



Rate (per 100,000 population)

<=0.4

0.41 - 2.2

>2.2

Texas, 2013

Primary and secondary syphilis

- CASES: Texas reported 1,475 cases in 2013.
- RATES: Texas ranked 13 among 50 states reporting at least one case of P&S syphilis with 5.7 cases per 100,000 population compared to the U.S. rate of 5.5.

Rates of primary and secondary syphilis, by sex

- In Texas, the rate among males was 10.0 cases per 100,000 population compared to the U.S. male rate of 10.3.
- The rate among females was 1.4 compared to the U.S. female rate of 0.9.


Rates of primary and secondary syphilis, by race/ethnicity

- In Texas, the race/ethnicity adjusted rates per 100,000 population were 3.4 among whites, 17.1 among blacks, 5.3 among Hispanics, 1.3 among Asians, 14.9 among Native Hawaiian/Other Pacific Islanders, and 1.2 among American Indians/Alaska Natives.
- The rate among blacks was 5 times that of whites.
- The rate among Hispanics was 1.6 times that of whites.
- The rate among Native Hawaiian/Other Pacific Islanders was 4.4 times that of whites.


Congenital syphilis

- CASES: Texas reported 75 cases in 2013.
- RATES: Texas ranked 3 among 25 states reporting at least one case of congenital syphilis with a rate of 19.4 cases per 100,000 live births compared to the U.S. rate of 8.7.

Syphilis: P&S Rates and Congenital Cases, Texas, 2004–2013



Total P&S Syphilis Rates by County, Texas, 2013



Rate (per 100,000 population)

≤0.4

0.41 - 2.2

>2.2

Utah, 2013

Primary and secondary syphilis

- CASES: Utah reported 74 cases in 2013.
- RATES: Utah ranked 38 among 50 states reporting at least one case of P&S syphilis with 2.6 cases per 100,000 population compared to the U.S. rate of 5.5.

Rates of primary and secondary syphilis, by sex

- In Utah, the rate among males was 5.0 cases per 100,000 population compared to the U.S. male rate of 10.3.
- The rate among females was 0.1 compared to the U.S. female rate of 0.9.


Rates of primary and secondary syphilis, by race/ethnicity

- In Utah, the race/ethnicity adjusted rates per 100,000 population were 2.5 among whites, 10.4 among blacks, 2.4 among Hispanics, 5.0 among Asians, 0.0 among Native Hawaiian/Other Pacific Islanders, and 3.6 among American Indians/Alaska Natives.
- The rate among blacks was 4.1 times that of whites.
- The rate among Asians was 1.9 times that of whites.
- The rate among American Indians/Alaska Natives was 1.4 times that of whites.


Congenital syphilis

- CASES: Utah reported 0 cases in 2013.
- RATES: Utah had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.7.

Syphilis: P&S Rates and Congenital Cases, Utah, 2004–2013



Total P&S Syphilis Rates by County, Utah, 2013



Rate (per 100,000 population)

<=0.4

0.41 - 2.2

>2.2

Vermont, 2013

Primary and secondary syphilis

- CASES: Vermont reported 3 cases in 2013.
- RATES: Vermont ranked 49 among 50 states reporting at least one case of P&S syphilis with 0.5 cases per 100,000 population compared to the U.S. rate of 5.5.

Rates of primary and secondary syphilis, by sex

- In Vermont, the rate among males was 1.0 cases per 100,000 population compared to the U.S. male rate of 10.3.
- The rate among females was 0.0 compared to the U.S. female rate of 0.9.


Rates of primary and secondary syphilis, by race/ethnicity

- In Vermont, the race/ethnicity adjusted rates per 100,000 population were 0.5 among whites, 0.0 among blacks, 0.0 among Hispanics, 0.0 among Asians, 0.0 among Native Hawaiian/Other Pacific Islanders, and 0.0 among American Indians/Alaska Natives.


Congenital syphilis

- CASES: Vermont reported 0 cases in 2013.
- RATES: Vermont had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.7.

Syphilis: P&S Rates and Congenital Cases, Vermont, 2004–2013



Total P&S Syphilis Rates by County, Vermont, 2013



Rate (per 100,000 population)

≤0.4

0.41 - 2.2

>2.2

Virginia, 2013

Primary and secondary syphilis

- CASES: Virginia reported 315 cases in 2013.
- RATES: Virginia ranked 22 among 50 states reporting at least one case of P&S syphilis with 3.8 cases per 100,000 population compared to the U.S. rate of 5.5.

Rates of primary and secondary syphilis, by sex

- In Virginia, the rate among males was 7.4 cases per 100,000 population compared to the U.S. male rate of 10.3.
- The rate among females was 0.4 compared to the U.S. female rate of 0.9.


Rates of primary and secondary syphilis, by race/ethnicity

- In Virginia, the race/ethnicity adjusted rates per 100,000 population were 2.3 among whites, 10.7 among blacks, 3.1 among Hispanics, 0.6 among Asians, 0.0 among Native Hawaiian/Other Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 4.6 times that of whites.
- The rate among Hispanics was 1.3 times that of whites.


Congenital syphilis

- CASES: Virginia reported 2 cases in 2013.
- RATES: Virginia ranked 22 among 25 states reporting at least one case of congenital syphilis with a rate of 1.9 cases per 100,000 live births compared to the U.S. rate of 8.7.

Syphilis: P&S Rates and Congenital Cases, Virginia, 2004–2013



Total P&S Syphilis Rates by County, Virginia, 2013



Rate (per 100,000 population)

≤0.4

0.41 - 2.2

>2.2

Washington, 2013

Primary and secondary syphilis

- CASES: Washington reported 284 cases in 2013.
- RATES: Washington ranked 21 among 50 states reporting at least one case of P&S syphilis with 4.1 cases per 100,000 population compared to the U.S. rate of 5.5.

Rates of primary and secondary syphilis, by sex

- In Washington, the rate among males was 7.9 cases per 100,000 population compared to the U.S. male rate of 10.3.
- The rate among females was 0.4 compared to the U.S. female rate of 0.9.


Rates of primary and secondary syphilis, by race/ethnicity

- In Washington, the race/ethnicity adjusted rates per 100,000 population were 3.9 among whites, 12.2 among blacks, 4.8 among Hispanics, 1.2 among Asians, 2.4 among Native Hawaiian/Other Pacific Islanders, and 4.4 among American Indians/Alaska Natives.
- The rate among blacks was 3.1 times that of whites.
- The rate among Hispanics was 1.2 times that of whites.
- The rate among American Indians/Alaska Natives was 1.1 times that of whites.


Congenital syphilis

- CASES: Washington reported 0 cases in 2013.
- RATES: Washington had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.7.

Syphilis: P&S Rates and Congenital Cases, Washington, 2004–2013



Total P&S Syphilis Rates by County, Washington, 2013



Rate (per 100,000 population)



West Virginia, 2013

Primary and secondary syphilis

- CASES: West Virginia reported 15 cases in 2013.
- RATES: West Virginia ranked 46 among 50 states reporting at least one case of P&S syphilis with 0.8 cases per 100,000 population compared to the U.S. rate of 5.5.

Rates of primary and secondary syphilis, by sex

- In West Virginia, the rate among males was 1.2 cases per 100,000 population compared to the U.S. male rate of 10.3.
- The rate among females was 0.4 compared to the U.S. female rate of 0.9.


Rates of primary and secondary syphilis, by race/ethnicity

- In West Virginia, the race/ethnicity adjusted rates per 100,000 population were 0.6 among whites, 6.2 among blacks, 0.0 among Hispanics, 0.0 among Asians, 0.0 among Native Hawaiian/Other Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 9.8 times that of whites.


Congenital syphilis

- CASES: West Virginia reported 0 cases in 2013.
- RATES: West Virginia had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.7.

Syphilis: P&S Rates and Congenital Cases, West Virginia, 2004–2013



Total P&S Syphilis Rates by County, West Virginia, 2013



Wisconsin, 2013

Primary and secondary syphilis

- CASES: Wisconsin reported 95 cases in 2013.
- RATES: Wisconsin ranked 43 among 50 states reporting at least one case of P&S syphilis with 1.7 cases per 100,000 population compared to the U.S. rate of 5.5.

Rates of primary and secondary syphilis, by sex

- In Wisconsin, the rate among males was 3.0 cases per 100,000 population compared to the U.S. male rate of 10.3.
- The rate among females was 0.3 compared to the U.S. female rate of 0.9.


Rates of primary and secondary syphilis, by race/ethnicity

- In Wisconsin, the race/ethnicity adjusted rates per 100,000 population were 1.1 among whites, 6.4 among blacks, 4.8 among Hispanics, 0.0 among Asians, 0.0 among Native Hawaiian/Other Pacific Islanders, and 2.0 among American Indians/Alaska Natives.
- The rate among blacks was 6.1 times that of whites.
- The rate among Hispanics was 4.5 times that of whites.
- The rate among American Indians/Alaska Natives was 1.9 times that of whites.


Congenital syphilis

- CASES: Wisconsin reported 0 cases in 2013.
- RATES: Wisconsin had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.7.

Syphilis: P&S Rates and Congenital Cases, Wisconsin, 2004–2013



Total P&S Syphilis Rates by County, Wisconsin, 2013



Rate (per 100,000 population)

≤0.4

0.41 - 2.2

>2.2

Wyoming, 2013

Primary and secondary syphilis

- CASES: Wyoming reported 1 case in 2013.
- RATES: Wyoming ranked 50 among 50 states reporting at least one case of P&S syphilis with 0.2 cases per 100,000 population compared to the U.S. rate of 5.5.

Rates of primary and secondary syphilis, by sex

- In Wyoming, the rate among males was 0.3 cases per 100,000 population compared to the U.S. male rate of 10.3.
- The rate among females was 0.0 compared to the U.S. female rate of 0.9.


Rates of primary and secondary syphilis, by race/ethnicity

- In Wyoming, the race/ethnicity adjusted rates per 100,000 population were 0.0 among whites, 0.0 among blacks, 1.8 among Hispanics, 0.0 among Asians, 0.0 among Native Hawaiian/Other Pacific Islanders, and 0.0 among American Indians/Alaska Natives.


Congenital syphilis

- CASES: Wyoming reported 0 cases in 2013.
- RATES: Wyoming had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.7.

Syphilis: P&S Rates and Congenital Cases, Wyoming, 2004–2013



Total P&S Syphilis Rates by County, Wyoming, 2013



Guam, 2013

Primary and secondary syphilis

- CASES: Guam reported 6 cases in 2013.
- RATES: Guam reported 3.8 cases per 100,000 population compared to the U.S. rate of 5.5.


Rates of primary and secondary syphilis, by sex

- In Guam, the rate among males was 1.2 cases per 100,000 population compared to the U.S. male rate of 10.3.
- The rate among females was 6.3 compared to the U.S. female rate of 0.9.


Congenital syphilis

- CASES: Guam reported 1 case in 2013.
- RATES: Guam had a rate of 29.3 cases per 100,000 live births compared to the U.S. rate of 8.7.

Syphilis: P&S Rates and Congenital Cases, Guam, 2004–2013



Total P&S Syphilis Rate, Guam, 2013



Puerto Rico, 2013

Primary and secondary syphilis

- CASES: Puerto Rico reported 385 cases in 2013.
- RATES: Puerto Rico reported 10.5 cases per 100,000 population compared to the U.S. rate of 5.5.


Rates of primary and secondary syphilis, by sex

- In Puerto Rico, the rate among males was 19.9 cases per 100,000 population compared to the U.S. male rate of 10.3.
- The rate among females was 1.8 compared to the U.S. female rate of 0.9.


Congenital syphilis

- CASES: Puerto Rico reported 1 case in 2013.
- RATES: Puerto Rico had a rate of 2.4 cases per 100,000 live births compared to the U.S. rate of 8.7.

Syphilis: P&S Rates and Congenital Cases, Puerto Rico, 2004–2013



Total P&S Syphilis Rates by Municipality, Puerto Rico, 2013



Rate (per 100,000 population)

≤0.4

0.41 - 2.2

>2.2

Virgin Islands, 2013

Primary and secondary syphilis

- CASES: Virgin Islands reported 2 cases in 2013.
- RATES: Virgin Islands reported 1.9 cases per 100,000 population compared to the U.S. rate of 5.5.


Rates of primary and secondary syphilis, by sex

- In Virgin Islands, the rate among males was 2.0 cases per 100,000 population compared to the U.S. male rate of 10.3.
- The rate among females was 1.8 compared to the U.S. female rate of 0.9.

Congenital syphilis

- CASES: Virgin Islands reported 0 cases in 2013.
- RATES: Virgin Islands had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.7.

Syphilis: P&S Rates and Congenital Cases, Virgin Islands, 2004–2013



Total P&S Syphilis Rate, Virgin Islands, 2013



Rate (per 100,000 population)

≤0.4

0.41 - 2.2

>2.2