


Self-Study Modules on Tuberculosis



Epidemiology of Tuberculosis

Module 2: Objectives

At completion of this module, learners will be able to:

1. Describe how the number of TB cases reported in the U.S has changed over the last 60 years
2. List 5 factors that contributed to the increase of TB cases between 1985 and 1992
3. List 3 improvements TB programs made with increased funds that have contributed to a decrease in TB cases since 1993
4. List groups of people who are more likely to be exposed to or infected with *M. tuberculosis*
5. List groups of people who are more likely to develop TB disease once infected with *M. tuberculosis*

Module 2 – Epidemiology of Tuberculosis

2

Module 2: Overview

- Introduction to TB Epidemiology
- People at High Risk for TB Infection and TB Disease
- Case Studies

Module 2 – Epidemiology of Tuberculosis

3

Introduction to TB Epidemiology

4

Epidemiology (1)

Epidemiology is the study of the distribution and causes of disease and other health problems in groups of people.

Module 2 – Epidemiology of Tuberculosis

5

Epidemiology (2)

- Epidemiologists:
 - Determine frequency and pattern of health problems in communities
 - Try to figure out why health problems are occurring

Module 2 – Epidemiology of Tuberculosis

6

Module 2 - Epidemiology of Tuberculosis

Global Epidemiology of TB

- TB is one of the leading causes of death due to infectious disease in the world
- Almost 2 billion people are infected with *M. tuberculosis*
- Each year about:
 - 9 million people develop TB disease
 - 1.5 million people die of TB

Module 2 – Epidemiology of Tuberculosis

7

TB Reporting in U.S.

- The Report of Verified Case of Tuberculosis (RVCT) is the national TB surveillance data collection form and is used for reporting all verified TB cases to CDC
 - The 50 states, District of Columbia, New York City, Puerto Rico, and 7 other jurisdictions in the Pacific and Caribbean report TB cases to CDC
- Health care providers are required by law to report TB cases to state or local health departments

Module 2 – Epidemiology of Tuberculosis

8


U.S. Epidemiology of TB 1953 - 1986

- 1953:
 - More than 84,000 cases of TB
- 1953-1984:
 - TB cases declined about 6% each year
- 1985:
 - TB cases reached a low of 22,201
- 1986:
 - Significant increase in TB cases began

Module 2 – Epidemiology of Tuberculosis

9

U.S. TB Resurgence (1) 1986 - 1992



Module 2 – Epidemiology of Tuberculosis

10

U.S. TB Resurgence (2) 1986 - 1992

- Contributing factors:
 - Inadequate funding for TB control and other public health efforts
 - HIV epidemic
 - Increased immigration from countries where TB is common
 - Spread of TB in certain settings (e.g., homeless shelters and correctional facilities)
 - Spread of multidrug-resistant TB (MDR TB)

Module 2 – Epidemiology of Tuberculosis

11


U.S. TB Control and Prevention (1) 1993 - 2014

- 1993-2014:
 - Number of TB cases reported annually in U.S. steadily declined
- Increased federal funds and other resources allowed TB programs to improve control efforts to:
 - Promptly identify persons with TB
 - Start appropriate initial treatment for TB cases
 - Ensure patients complete treatment
 - Conduct contact investigations

Module 2 – Epidemiology of Tuberculosis

12

U.S. TB Control and Prevention (2) 1993 - 2014



Module 2 – Epidemiology of Tuberculosis

13

Continuing Challenges in TB Control

- TB is reported in almost every state and is increasing in some areas
- More than half of all TB cases in the U.S. are among foreign-born persons
- TB affects racial/ethnic minorities disproportionately
- MDR TB and extensively drug-resistant TB (XDR TB) remain serious public health concerns

Module 2 – Epidemiology of Tuberculosis

14

TB Case Rates (1)

- A case rate is the number of TB cases that occur during a certain time period, divided by size of the population at that time
- Often expressed in terms of a population size of 100,000 persons

Module 2 – Epidemiology of Tuberculosis

15

TB Case Rates (2)

Example:

- In the U.S. in 2014, there were 9,421 new TB cases in a population size of 318,857,056


$$\frac{9,421}{318,857,056} \times 100,000 = 2.96$$

- In 2014, the U.S. TB case rate was 2.96 TB cases per 100,000 persons (rounded to 3.0)

Module 2 – Epidemiology of Tuberculosis

16

TB Case Rates by State, 2014



Module 2 – Epidemiology of Tuberculosis

17

TB Case Rates (3)

- Health departments, CDC, and others can compare the occurrence of TB in different places, time periods, and groups of people using case rates
- Comparisons have shown that rates of TB are higher in certain groups than in others

Module 2 – Epidemiology of Tuberculosis

18

Epidemiology of TB Study Question 2.1

What happened to the number of TB cases in the United States between 1953 and 1984?

Module 2 – Epidemiology of Tuberculosis

19

Epidemiology of TB Study Question 2.2

What happened to the number of TB cases in the United States between 1985 and 1992?

Module 2 – Epidemiology of Tuberculosis

20

Epidemiology of TB Study Question 2.3

Name 5 factors that may have contributed to the increase in the number of TB cases between 1985 and 1992.

Module 2 – Epidemiology of Tuberculosis

21

Epidemiology of TB Study Question 2.4

What happened to the number of TB cases in the United States from 1993 to 2014?

Module 2 – Epidemiology of Tuberculosis

22

Epidemiology of TB Study Question 2.5

Name 3 improvements TB programs were able to make with increased federal, state, and other funds that contributed to the decrease in TB cases since 1993.

Module 2 – Epidemiology of Tuberculosis



23

Race and Ethnicity (1)

- **TB affects certain racial and ethnic minorities disproportionately**
- **In 2014, about 85% of TB cases in the U.S. were among racial and ethnic minorities**
- **Percentage of TB cases in racial and ethnic minorities is higher than expected based on percentage of these minorities in the U.S. population**

Module 2 – Epidemiology of Tuberculosis

24



- ### Race and Ethnicity (4)
- Disparities may exist due to racial and ethnic minorities having other risk factors for TB, such as:
 - Birth in a country where TB is common
 - HIV infection
 - Low socioeconomic status
 - Exposure to TB in high-risk settings
- Module 2 – Epidemiology of Tuberculosis 27

Relative Risk for TB (1)

Relative risk is a comparison of case rates between two groups.

Module 2 – Epidemiology of Tuberculosis 28

Relative Risk for TB (2)

Example:

- The case rate for Asians is 17.8 compared to 0.6 for non-Hispanic whites. Therefore, the relative risk for Asians is about 29 times higher than non-Hispanic whites

$$\frac{17.8 \text{ (TB case rate for Asians)}}{0.6 \text{ (TB case rate for non-Hispanic whites)}} = 29.6$$

Module 2 – Epidemiology of Tuberculosis 29

Relative Risk for TB (3) Race and Ethnicity, 2014

Race/Ethnicity	TB Case Rate	Relative Risk
Asians	17.8	29.6
Native Hawaiians or Other Pacific Islanders	16.9	28.1
Blacks or African Americans	5.1	8.5
American Indians or Alaska Natives	5.0	8.3
Hispanics or Latinos	5.0	8.3
Multiple Race	2.8	4.6
Non- Hispanic Whites	0.6	1

Module 2 – Epidemiology of Tuberculosis 30

Race and Ethnicity Study Question 2.6

Which racial and ethnic groups are disproportionately affected by TB?

People at High Risk for TB Infection and TB Disease

High-Risk Groups

- High-risk groups can be divided into two categories:
 - High risk for exposure to or infection with *M. tuberculosis*
 - High risk for developing TB disease after infection with *M. tuberculosis*

People at High Risk for Exposure to or Infection with *M. tuberculosis*

- Contacts
- People who have come to the U.S. within the last 5 years from areas of the world where TB is common
- Persons who visit areas with a high prevalence of TB disease
- People who live or work in high-risk congregate settings
- Health care workers who serve patients at increased risk
- Populations defined locally as having an increased incidence of LTBI or TB disease, possibly medically underserved, low-income populations, or persons who abuse drugs or alcohol
- Infants, children, and adolescents exposed to adults who are at increased risk for LTBI or TB disease

People at High Risk for Developing TB Disease after Infection with *M. tuberculosis* (1)

- People living with HIV
- Children younger than 5 years of age
- People infected with *M. tuberculosis* within past 2 years
- People with a history of untreated or inadequately treated TB disease
- People who are receiving immunosuppressive therapy
- People with silicosis, diabetes mellitus, chronic renal failure, leukemia, or cancer of the head, neck, or lung

People at High Risk for Developing TB Disease after Infection with *M. tuberculosis* (2)

- Persons who have had a gastrectomy or jejunioileal bypass
- Low body weight
- Cigarette smokers and persons who abuse drugs or alcohol
- Populations defined locally as having an increased risk

High-Risk Groups for TB Infection (1) Contacts

- Contacts are persons who have spent time with someone who has infectious TB disease
- May include:
 - Family members
 - Coworkers
 - Friends




Module 2 – Epidemiology of Tuberculosis

37

High-Risk Groups for TB Infection (2) Foreign-Born Persons/Immigrants

In the U.S., LTBI and TB disease often occur among people born in areas of the world where TB is common:


- Asia
- Africa
- Russia
- Eastern Europe
- Latin America



Module 2 – Epidemiology of Tuberculosis

38

High-Risk Groups for TB Infection (3) Foreign-Born Persons/Immigrants



Cases of TB in foreign-born and U.S.-born, 1992 and 2014

Module 2 – Epidemiology of Tuberculosis

39

High-Risk Groups for TB Infection (4) Foreign-Born Persons/Immigrants

- To address high rates of TB in foreign-born persons, CDC and other public health organizations are working to:
 - Improve the overseas and domestic screening process for immigrants and refugees
 - Strengthen the notification system that alerts health departments about the arrival of immigrants and refugees with suspected TB disease
 - Test recent arrivals from countries where TB is common for TB infection and ensure completion of treatment

Module 2 – Epidemiology of Tuberculosis

40

High-Risk Groups for TB Infection (5) Foreign-Born Persons/Immigrants

- Individuals applying for immigration and refugee status from overseas:
 - Must be screened for TB by panel physicians before entering the U.S.
 - Must have completed treatment before entering the U.S. if diagnosed with TB disease

Module 2 – Epidemiology of Tuberculosis

41

High-Risk Groups for TB Infection (6) Foreign-Born Persons/Immigrants

- Immigrants living in the U.S. who apply for permanent residence or citizenship:
 - Must be tested for TB infection and evaluated for TB disease by U.S.- based civil surgeons



Module 2 – Epidemiology of Tuberculosis

42

High-Risk Groups for TB Infection (7) Congregate Settings

- In certain congregate settings, the risk of being exposed to TB is higher than other places. This may include:
 - Correctional facilities
 - Homeless shelters
 - Nursing homes
 - Health care facilities

Module 2 – Epidemiology of Tuberculosis

43

High-Risk Groups for TB Infection (8) Congregate Settings

- Risk of exposure to TB is higher than in other settings
- Risk is higher if facility is crowded

Module 2 – Epidemiology of Tuberculosis

44

High-Risk Groups for TB Infection (9) Correctional Facilities

- Higher risk in correctional facilities may be due to:
 - Incarcerated population includes a high proportion of people at greater risk for TB than overall population (risk factors may include HIV-infection and a history of homelessness or drug use)
 - Physical structure of correctional facilities (e.g., close living quarters, overcrowding, potential for inadequate ventilation)
 - Movement of inmates into and out of facilities can lead to interruption of therapy

Module 2 – Epidemiology of Tuberculosis

45

High-Risk Groups for TB Infection (10) Health Care Workers

- Might be exposed to TB at work
- Risk depends on:
 - Number of persons with TB in facility
 - Job duties
 - Infection control procedures



Module 2 – Epidemiology of Tuberculosis

46

High-Risk Groups for TB Infection (11) Populations Defined Locally

Populations that may have an increased risk include

- Persons experiencing homelessness
- Medically underserved populations
- Low-income groups
- Persons who abuse drugs or alcohol



Module 2 – Epidemiology of Tuberculosis

47

High-Risk Groups for TB Infection (12) Populations Defined Locally

- Low-income is linked to higher risk of TB exposure
- Possible reasons include factors associated with low-income:
 - Inadequate living conditions
 - Crowding
 - Malnutrition
 - Poor access to health care
- TB rates are 10 times higher for people experiencing homelessness

Module 2 – Epidemiology of Tuberculosis

48

High-Risk Groups for TB Infection (13) Children and Adolescents

- High risk if exposed to adults in high-risk groups
- If a child has TB infection or disease, it suggests that:
 - TB was transmitted relatively recently
 - Person who transmitted TB to child may still be infectious
 - Others may have been exposed



Module 2 – Epidemiology of Tuberculosis

49

High-Risk Groups for TB Disease (1) Infants and Children Younger than 5 Years

High risk for rapidly developing TB disease due to underdeveloped immune system



Module 2 – Epidemiology of Tuberculosis

50

High-Risk Groups for TB Disease (2) People Living with HIV

- HIV is the strongest known risk factor for developing TB disease
- TB is the leading cause of death for people with HIV/AIDS
- Risk of developing TB disease is 7% - 10% each year for people who are infected with both TB and HIV (if the HIV is not treated)

Module 2 – Epidemiology of Tuberculosis

51

High-Risk Groups for TB Infection Study Question 2.7

Name 7 groups of people who are more likely to be exposed or infected with *M. tuberculosis*.

Module 2 – Epidemiology of Tuberculosis

52

High-Risk Groups for TB Infection Study Question 2.8

What are public health agencies doing to address the high rate of TB in foreign born persons?

Module 2 – Epidemiology of Tuberculosis

53

High-Risk Groups for TB Infection Study Question 2.9

Why is the risk of being exposed to TB higher in certain settings, such as nursing homes or correctional facilities?

Module 2 – Epidemiology of Tuberculosis

54

**High-Risk Groups for TB Infection
Study Question 2.10**

What are some reasons why rates of TB disease are higher in correctional facilities?

**High-Risk Groups for TB Infection
Study Question 2.11**

When a child has TB infection or disease, what does it tell us about the spread of TB in the child's home or community? Name 3 things.

**High-Risk Groups for TB Disease
Study Question 2.12**

Name 8 groups of people who are more likely to develop TB disease once infected.

**High-Risk Groups for TB Disease
Study Question 2.13**

What is the strongest known risk factor for the development of TB disease?

**High-Risk Groups for TB Disease
Study Question 2.14**

If a person is infected with both *M. tuberculosis* and HIV, what are his or her chances of developing TB disease? How does this compare to the risk for people who are infected only with *M. tuberculosis*?

Case Studies

Module 2: Case Study 2.1

For each of the following people, choose the factor(s) known to increase the risk of being exposed to or infected with TB

Person	Factors
a) Mr. Petrov	<input type="checkbox"/> Works at a nursing home <input type="checkbox"/> Rides the subway every day <input type="checkbox"/> Emigrated from Russia
b) Ms. Montoya	<input type="checkbox"/> Was born in Latin America <input type="checkbox"/> Has a father who had pulmonary TB disease
c) Ms. Parker	<input type="checkbox"/> Volunteers in the emergency room of an inner-city hospital <input type="checkbox"/> Works in a day care center
d) Mr. Dudley	<input type="checkbox"/> Was released from prison last year <input type="checkbox"/> Sleeps in a homeless shelter

Module 2 – Epidemiology of Tuberculosis

61

Module 2: Case Study 2.2

For each of the following people, indicate the factor(s) known to increase the risk of developing TB disease once infected

Person	Factors
a) Mr. Sims	<input type="checkbox"/> Injects heroin <input type="checkbox"/> Has HIV
b) Mr. Allen	<input type="checkbox"/> Has diabetes <input type="checkbox"/> Has high blood pressure
c) Ms. Li	<input type="checkbox"/> Has chest x-ray findings suggestive of previous TB disease <input type="checkbox"/> Has heart problems
d) Mr. Vinson	<input type="checkbox"/> Is overweight <input type="checkbox"/> Became infected with <i>M. tuberculosis</i> 6 months ago

Module 2 – Epidemiology of Tuberculosis

62