Table 39. Extrapulmonary Tuberculosis Cases and Percentages by Site of Disease: Reporting Areas, 2012

Reporting Area	Total Extrapulm. Cases[1]	Total Extrapulm. Sites[2]	Site of Disease: Pleural No.	Pleural (%)	Lymphatic No.	Lymphatic (%)	Bone and/or Joint No.	Bone and/or Joint (%)	Genitourinary No.	Genitourinary (%)	Meningeal No.	Meningeal (%)	Peritoneal No.	Peritoneal (%)	Laryngeal No.	Laryngeal (%)	Other No.	Other (%)
United States	**2,100**	**2,213**	**368**	**(16.6)**	**846**	**(38.2)**	**229**	**(10.3)**	**111**	**(5.0)**	**108**	**(4.9)**	**127**	**(5.7)**	**8**	**(0.4)**	**416**	**(18.8)**
Alabama	21	23	7	(30.4)	5	(21.7)	2	(8.7)	0	(0.0)	1	(4.3)	0	(0.0)	0	(0.0)	8	(34.8)
Alaska	6	6	0	(0.0)	5	(83.3)	0	(0.0)	0	(0.0)	0	(0.0)	0	(0.0)	0	(0.0)	1	(16.7)
Arizona	27	28	5	(17.9)	14	(50.0)	1	(3.6)	1	(3.6)	2	(7.1)	2	(7.1)	0	(0.0)	3	(10.7)
Arkansas	20	20	3	(15.0)	3	(15.0)	2	(10.0)	0	(0.0)	2	(10.0)	0	(0.0)	1	(5.0)	9	(45.0)
California	499	517	70	(13.5)	211	(40.8)	45	(8.7)	24	(4.6)	22	(4.3)	35	(6.8)	2	(0.4)	108	(20.9)
Colorado	19	21	4	(19.0)	6	(28.6)	4	(19.0)	2	(9.5)	2	(9.5)	0	(0.0)	0	(0.0)	3	(14.3)
Connecticut	18	19	6	(31.6)	9	(47.4)	2	(10.5)	1	(5.3)	1	(5.3)	0	(0.0)	0	(0.0)	0	(0.0)
Delaware	11	14	2	(14.3)	0	(0.0)	2	(14.3)	2	(14.3)	3	(21.4)	1	(7.1)	0	(0.0)	4	(28.6)
District of Columbia	14	15	2	(13.3)	4	(26.7)	2	(13.3)	0	(0.0)	3	(20.0)	1	(6.7)	1	(6.7)	2	(13.3)
Florida	105	108	27	(25.0)	39	(36.1)	10	(9.3)	6	(5.6)	5	(4.6)	7	(6.5)	0	(0.0)	14	(13.0)
Georgia	75	75	17	(22.7)	28	(37.3)	5	(6.7)	3	(4.0)	4	(5.3)	1	(1.3)	2	(2.7)	15	(20.0)
Hawaii	23	23	11	(47.8)	6	(26.1)	0	(0.0)	0	(0.0)	2	(8.7)	1	(4.3)	0	(0.0)	3	(13.0)
Idaho	1	1	0	(0.0)	0	(0.0)	0	(0.0)	0	(0.0)	0	(0.0)	0	(0.0)	0	(0.0)	1	(100.0)
Illinois	90	97	10	(10.3)	36	(37.1)	10	(10.3)	3	(3.1)	4	(4.1)	8	(8.2)	0	(0.0)	26	(26.8)
Indiana	17	19	1	(5.3)	4	(21.1)	4	(21.1)	4	(21.1)	2	(10.5)	1	(5.3)	0	(0.0)	3	(15.8)
Iowa	8	8	4	(50.0)	2	(25.0)	2	(25.0)	0	(0.0)	0	(0.0)	0	(0.0)	0	(0.0)	0	(0.0)
Kansas	7	7	1	(14.3)	3	(42.9)	2	(28.6)	0	(0.0)	0	(0.0)	0	(0.0)	0	(0.0)	1	(14.3)
Kentucky	8	9	2	(22.2)	1	(11.1)	0	(0.0)	0	(0.0)	0	(0.0)	1	(11.1)	0	(0.0)	5	(55.6)
Louisiana	31	31	5	(16.1)	13	(41.9)	4	(12.9)	1	(3.2)	1	(3.2)	0	(0.0)	0	(0.0)	7	(22.6)
Maine	4	4	1	(25.0)	1	(25.0)	0	(0.0)	1	(25.0)	0	(0.0)	0	(0.0)	0	(0.0)	1	(25.0)
Maryland	66	71	16	(22.5)	26	(36.6)	3	(4.2)	2	(2.8)	3	(4.2)	1	(1.4)	1	(1.4)	19	(26.8)
Massachusetts	61	64	10	(15.6)	25	(39.1)	6	(9.4)	4	(6.3)	5	(7.8)	4	(6.3)	0	(0.0)	10	(15.6)
Michigan	47	48	5	(10.4)	15	(31.3)	6	(12.5)	1	(2.1)	1	(2.1)	1	(2.1)	0	(0.0)	19	(39.6)
Minnesota	58	63	7	(11.1)	31	(49.2)	6	(9.5)	1	(1.6)	2	(3.2)	5	(7.9)	0	(0.0)	11	(17.5)
Mississippi	10	10	6	(60.0)	1	(10.0)	2	(20.0)	0	(0.0)	0	(0.0)	1	(10.0)	0	(0.0)	0	(0.0)
Missouri	17	18	4	(22.2)	7	(38.9)	1	(5.6)	1	(5.6)	1	(5.6)	1	(5.6)	0	(0.0)	3	(16.7)
Nebraska	10	10	0	(0.0)	5	(50.0)	2	(20.0)	0	(0.0)	0	(0.0)	2	(20.0)	0	(0.0)	1	(10.0)
Nevada	11	12	0	(0.0)	3	(25.0)	1	(8.3)	5	(41.7)	1	(8.3)	1	(8.3)	0	(0.0)	1	(8.3)
New Hampshire	2	2	0	(0.0)	1	(50.0)	0	(0.0)	0	(0.0)	0	(0.0)	0	(0.0)	0	(0.0)	1	(50.0)
New Jersey	69	73	16	(21.9)	30	(41.1)	15	(20.5)	2	(2.7)	2	(2.7)	6	(8.2)	0	(0.0)	2	(2.7)
New Mexico	7	7	1	(14.3)	3	(42.9)	2	(28.6)	0	(0.0)	0	(0.0)	0	(0.0)	0	(0.0)	1	(14.3)

Table 39. (Con't) Extrapulmonary Tuberculosis Cases and Percentages by Site of Disease: Reporting Areas, 2012

Reporting Area	Total Extrapulm. Cases[1]	Total Extrapulm. Sites[2]	Site of Disease: Pleural		Lymphatic		Bone and/or Joint		Genitourinary		Meningeal		Peritoneal		Laryngeal		Other	
			No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)
New York	214	238	33	(13.9)	105	(44.1)	27	(11.3)	11	(4.6)	8	(3.4)	16	(6.7)	1	(0.4)	37	(15.5)
North Carolina	53	56	16	(28.6)	23	(41.1)	3	(5.4)	4	(7.1)	2	(3.6)	3	(5.4)	0	(0.0)	5	(8.9)
North Dakota	5	5	2	(40.0)	1	(20.0)	0	(0.0)	0	(0.0)	0	(0.0)	1	(20.0)	0	(0.0)	1	(20.0)
Ohio	37	38	5	(13.2)	20	(52.6)	5	(13.2)	2	(5.3)	1	(2.6)	0	(0.0)	0	(0.0)	5	(13.2)
Oklahoma	18	18	3	(16.7)	8	(44.4)	3	(16.7)	0	(0.0)	1	(5.6)	0	(0.0)	0	(0.0)	3	(16.7)
Oregon	19	20	3	(15.0)	7	(35.0)	2	(10.0)	2	(10.0)	0	(0.0)	1	(5.0)	0	(0.0)	5	(25.0)
Pennyslvania	69	70	10	(14.3)	25	(35.7)	9	(12.9)	7	(10.0)	4	(5.7)	6	(8.6)	0	(0.0)	9	(12.9)
Rhode Island	7	9	3	(33.3)	2	(22.2)	2	(22.2)	0	(0.0)	0	(0.0)	1	(11.1)	0	(0.0)	1	(11.1)
South Carolina	23	24	8	(33.3)	8	(33.3)	3	(12.5)	1	(4.2)	2	(8.3)	0	(0.0)	0	(0.0)	2	(8.3)
South Dakota	3	3	1	(33.3)	1	(33.3)	0	(0.0)	0	(0.0)	0	(0.0)	0	(0.0)	0	(0.0)	1	(33.3)
Tennessee	34	35	8	(22.9)	17	(48.6)	1	(2.9)	0	(0.0)	4	(11.4)	1	(2.9)	0	(0.0)	4	(11.4)
Texas	119	123	7	(5.7)	42	(34.1)	16	(13.0)	10	(8.1)	12	(9.8)	7	(5.7)	0	(0.0)	29	(23.6)
Utah	12	12	1	(8.3)	2	(16.7)	2	(16.7)	1	(8.3)	0	(0.0)	0	(0.0)	0	(0.0)	6	(50.0)
Vermont	1	1	0	(0.0)	1	(100.0)	0	(0.0)	0	(0.0)	0	(0.0)	0	(0.0)	0	(0.0)	0	(0.0)
Virginia	58	64	10	(15.6)	20	(31.3)	6	(9.4)	4	(6.3)	3	(4.7)	7	(10.9)	0	(0.0)	14	(21.9)
Washington	48	52	9	(17.3)	21	(40.4)	7	(13.5)	4	(7.7)	2	(3.8)	2	(3.8)	0	(0.0)	7	(13.5)
West Virginia	1	1	0	(0.0)	0	(0.0)	0	(0.0)	0	(0.0)	0	(0.0)	0	(0.0)	0	(0.0)	1	(100.0)
Wisconsin	17	21	6	(28.6)	6	(28.6)	2	(9.5)	1	(4.8)	0	(0.0)	2	(9.5)	0	(0.0)	4	(19.0)
Wyoming	0	0	0	(0.0)	0	(0.0)	0	(0.0)	0	(0.0)	0	(0.0)	0	(0.0)	0	(0.0)	0	(0.0)
American Samoa[3]	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...
Fed. States of Micronesia[3]	19	20	11	(55.0)	7	(35.0)	2	(10.0)	0	(0.0)	0	(0.0)	0	(0.0)	0	(0.0)	0	(0.0)
Guam[3]	8	8	3	(37.5)	1	(12.5)	1	(12.5)	0	(0.0)	0	(0.0)	1	(12.5)	0	(0.0)	2	(25.0)
Marshall Islands[3]	25	29	8	(27.6)	13	(44.8)	0	(0.0)	0	(0.0)	0	(0.0)	7	(24.1)	0	(0.0)	1	(3.4)
N. Mariana Islands[3]	2	2	0	(0.0)	0	(0.0)	2	(100.0)	0	(0.0)	0	(0.0)	0	(0.0)	0	(0.0)	0	(0.0)
Puerto Rico[3]	10	10	5	(50.0)	1	(10.0)	0	(0.0)	0	(0.0)	1	(10.0)	2	(20.0)	0	(0.0)	1	(10.0)
Republic of Palau[3]	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...
U.S. Virgin Islands[3]	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...

[1] Excludes cases with pulmonary site of disease.
[2] Patient may have more than one extrapulmonary site of disease.
[3] Not included in U.S. totals.
Note: Ellipses indicate data not available.
See Technical Notes.