

Table 27. Tuberculosis Cases and Percentages by Reason Tuberculosis Therapy Stopped and Type of Move: United States, 2010

Type of Move	Total Cases	Completed Therapy		Adverse Event		Lost		Refused		Died		Other ¹		Unknown	
	No.	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)
Moved in state ²	331	265	(80.1)	1	(0.3)	14	(4.2)	1	(0.3)	17	(5.1)	26	(7.9)	7	(2.1)
Moved out of state ³	305	228	(74.8)	1	(0.3)	23	(7.5)	1	(0.3)	7	(2.3)	35	(11.5)	10	(3.3)
Moved out of country ⁴	436	163	(37.4)	0	(0.0)	47	(10.8)	7	(1.6)	6	(1.4)	199	(45.6)	11	(2.5)
Did not move ⁵	9,692	8,780	(90.6)	27	(0.3)	83	(0.9)	56	(0.6)	609	(6.3)	27	(0.3)	108	(1.1)

¹ Therapy was discontinued for a known reason other than those listed (e.g. patient moved outside the U.S., or patient moved from state A to state B, and though state A notified state B, state B never followed up).

² Includes patients who were alive at diagnosis, started on treatment, and moved in state.

³ Includes patients who were alive at diagnosis, started on treatment, and moved out of state.

⁴ Includes patients who were alive at diagnosis, started on treatment, and moved out of the country; transnational referrals were provided for 262 (51.8%) TB patients who moved out of the country.

⁵ Includes patients who were alive at diagnosis, started on treatment, and did not indicate having moved.

Note: There may be differences in the way jurisdictions determine treatment completion for patients who moved out of the country; some reporting jurisdictions may be classifying all patients who moved out of the country as 'other' for reason therapy stopped.