Table 28. Deaths Prior to Tuberculosis Diagnosis or During Tuberculosis Therapy by Age Group: United States, 2010

	Total			Dead at Diagnosis							Died During Therapy[1]						
	Total Deaths Reported	Deaths Related to TB Disease or Therapy[2]		Total Dead at Diagnosis	TB a Cause of Death		TB Not a Cause of Death		Unknown		Total Died During Therapy	Related to TB[3]		Unrelated to TB		Unknown	
Age Group	No.	No.	(%)	No.	No.	(%)	No.	(%)	No.	(%)	No.	No.	(%)	No.	(%)	No.	(%)
Total	**893**	**320**	**(35.8)**	**252**	**85**	**(33.7)**	**123**	**(48.8)**	**44**	**(17.5)**	**641**	**235**	**(36.7)**	**294**	**(45.9)**	**112**	**(17.5)**
0–4	0	0	...	0	0	...	0	...	0	...	0	0	...	0	...	0	...
5–14	2	1	(50.0)	1	0	(0.0)	1	(100.0)	0	(0.0)	1	1	(100.0)	0	(0.0)	0	(0.0)
15–24	11	5	(45.5)	6	3	(50.0)	3	(50.0)	0	(0.0)	5	2	(40.0)	3	(60.0)	0	(0.0)
25–44	95	41	(43.2)	40	16	(40.0)	16	(40.0)	8	(20.0)	55	25	(45.5)	20	(36.4)	10	(18.2)
45–64	294	114	(38.8)	84	32	(38.1)	37	(44.0)	15	(17.9)	210	82	(39.0)	99	(47.1)	29	(13.8)
≥65	491	159	(32.4)	121	34	(28.1)	66	(54.5)	21	(17.4)	370	125	(33.8)	172	(46.5)	73	(19.7)

[1] Among patients alive at diagnosis. Excludes 16 patients who died during therapy but did not start on therapy or unknown whether or not therapy was started.
[2] Includes patients who were dead at diagnosis or died during therapy, for which TB or TB therapy was indicated as a cause of death.
[3] Eight patient deaths during therapy were related to TB therapy.
Note: Ellipses indicate data not available.