Table 59. Completion of Tuberculosis Therapy (COT) Cases and Percentages[1] by Hispanic Ethnicity and Non-Hispanic Race: Reporting Areas, 2010[2]

Reporting Area	Total Cases[3]	Hispanic[4]		Non-Hispanic													
				American Indian or Alaska Native		Asian		Black		Native Hawaiian or Other Pacific Islander		White		Multiple Race		Unknown or Missing	
		No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)
United States	**9,480**	**2686**	**(87.5)**	**123**	**(90.2)**	**2666**	**(87.7)**	**2312**	**(88.8)**	**84**	**(92.9)**	**1461**	**(88.1)**	**130**	**(92.3)**	**18**	**(94.4)**
Alabama	124	21	(95.2)	1	(100.0)	10	(80.0)	50	(86.0)	0	...	42	(95.2)	0	...	0	...
Alaska	50	0	...	36	(91.7)	7	(100.0)	2	(100.0)	0	...	5	(100.0)	0	...	0	...
Arizona	220	93	(82.8)	21	(95.2)	53	(86.8)	18	(100.0)	0	...	35	(80.0)	0	...	0	...
Arkansas	70	8	(100.0)	0	...	8	(87.5)	22	(77.3)	3	(100.0)	29	(89.7)	0	...	0	...
California	1,967	737	(83.7)	0	...	828	(86.7)	133	(88.7)	12	(83.3)	156	(87.8)	96	(90.6)	5	(80.0)
Colorado	59	19	(89.5)	1	(100.0)	15	(86.7)	15	(100.0)	2	(100.0)	7	(100.0)	0	...	0	...
Connecticut	70	19	(89.5)	0	...	27	(81.5)	10	(90.0)	0	...	12	(83.3)	2	(100.0)	0	...
Delaware	16	2	(100.0)	0	...	6	(83.3)	8	(100.0)	0	...	0	...	0	...	0	...
District of Columbia	34	7	(71.4)	0	...	2	(100.0)	24	(83.3)	0	...	1	(100.0)	0	...	0	...
Florida	718	221	(94.1)	2	(100.0)	78	(93.6)	246	(95.5)	3	(100.0)	167	(91.6)	1	(100.0)	0	...
Georgia	341	53	(86.8)	0	...	71	(93.0)	174	(83.9)	0	...	42	(92.9)	1	(100.0)	0	...
Hawaii	105	1	(100.0)	0	...	80	(95.0)	0	...	18	(94.4)	6	(100.0)	0	...	0	...
Idaho	14	4	(50.0)	0	...	1	(100.0)	4	(100.0)	0	...	4	(100.0)	1	(100.0)	0	...
Illinois	317	98	(87.8)	0	...	86	(83.7)	73	(87.7)	0	...	60	(86.7)	0	...	0	...
Indiana	81	22	(100.0)	0	...	17	(94.1)	16	(93.8)	0	...	26	(84.6)	0	...	0	...
Iowa	41	10	(100.0)	0	...	14	(92.9)	11	(81.8)	0	...	6	(100.0)	0	...	0	...
Kansas	40	11	(100.0)	1	(100.0)	12	(100.0)	12	(100.0)	0	...	4	(100.0)	0	...	0	...
Kentucky	78	10	(90.0)	0	...	11	(72.7)	19	(84.2)	0	...	38	(84.2)	0	...	0	...
Louisiana	179	21	(71.4)	0	...	21	(71.4)	99	(83.8)	0	...	37	(81.1)	1	(100.0)	0	...
Maine	6	1	(100.0)	0	...	0	...	4	(100.0)	0	...	1	(100.0)	0	...	0	...
Maryland	187	37	(81.1)	1	(0.0)	54	(90.7)	81	(96.3)	0	...	14	(85.7)	0	...	0	...
Massachusetts	195	28	(85.7)	0	...	78	(78.2)	51	(88.2)	0	...	37	(83.8)	1	(100.0)	0	...
Michigan	148	16	(93.8)	0	...	46	(93.5)	57	(84.2)	1	(100.0)	27	(92.6)	0	...	1	(100.0)
Minnesota	123	10	(100.0)	3	(66.7)	35	(82.9)	65	(90.8)	0	...	9	(77.8)	1	(100.0)	0	...
Mississippi	106	8	(87.5)	1	(100.0)	5	(60.0)	65	(92.3)	0	...	27	(96.3)	0	...	0	...
Missouri	96	11	(72.7)	1	(100.0)	24	(87.5)	27	(85.2)	3	(100.0)	28	(71.4)	0	...	2	(100.0)
Montana	6	0	...	5	(100.0)	0	...	0	...	0	...	1	(100.0)	0	...	0	...
Nebraska	25	11	(90.9)	0	...	4	(100.0)	6	(100.0)	0	...	3	(100.0)	1	(0.0)	0	...
Nevada	108	28	(78.6)	1	(100.0)	39	(89.7)	21	(85.7)	2	(100.0)	16	(87.5)	1	(100.0)	0	...
New Hampshire	8	2	(100.0)	0	...	2	(100.0)	3	(100.0)	0	...	1	(0.0)	0	...	0	...
New Jersey	344	103	(91.3)	0	...	134	(88.8)	73	(91.8)	0	...	34	(94.1)	0	...	0	...

Table 59. (Cont'd) Completion of Tuberculosis Therapy (COT) Cases and Percentages[1] by Hispanic Ethnicity and Non-Hispanic Race: Reporting Areas, 2010[2]

Reporting Area	Total Cases[3]	Hispanic[4]		Non-Hispanic													
				American Indian or Alaska Native		Asian		Black		Native Hawaiian or Other Pacific Islander		White		Multiple Race		Unknown or Missing	
		No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)
New Mexico	31	16	(87.5)	8	(87.5)	2	(100.0)	2	(100.0)	0	...	3	(100.0)	0	...	0	...
New York State[5]	215	72	(86.1)	0	...	65	(95.4)	40	(90.0)	0	...	37	(89.2)	1	(100.0)	0	...
New York City	616	173	(90.8)	0	...	244	(93.4)	140	(93.6)	1	(100.0)	44	(90.9)	7	(100.0)	7	(100.0)
North Carolina	264	50	(94.0)	5	(100.0)	50	(94.0)	98	(94.9)	4	(100.0)	51	(94.1)	6	(100.0)	0	...
North Dakota	8	0	...	2	(0.0)	2	(100.0)	4	(100.0)	0	...	0	...	0	...	0	...
Ohio	157	19	(84.2)	0	...	22	(77.3)	71	(95.8)	0	...	44	(84.1)	0	...	1	(100.0)
Oklahoma	77	13	(92.3)	10	(100.0)	13	(92.3)	11	(90.9)	6	(83.3)	22	(95.5)	2	(100.0)	0	...
Oregon	74	18	(100.0)	1	(100.0)	21	(100.0)	13	(92.3)	3	(100.0)	18	(100.0)	0	...	0	...
Pennsylvania	191	19	(94.7)	0	...	66	(86.4)	64	(78.1)	0	...	40	(92.5)	2	(100.0)	0	...
Rhode Island	23	8	(62.5)	0	...	7	(57.1)	4	(75.0)	0	...	4	(75.0)	0	...	0	...
South Carolina	132	22	(81.8)	0	...	8	(75.0)	71	(91.5)	8	(100.0)	23	(100.0)	0	...	0	...
South Dakota	12	0	...	10	(90.0)	0	...	1	(100.0)	0	...	1	(100.0)	0	...	0	...
Tennessee	156	35	(91.4)	0	...	21	(90.5)	48	(93.8)	0	...	51	(88.2)	1	(100.0)	0	...
Texas	1,117	544	(87.7)	4	(75.0)	176	(83.0)	220	(75.9)	3	(100.0)	169	(81.1)	1	(100.0)	0	...
Utah	16	7	(100.0)	0	...	2	(100.0)	5	(100.0)	0	...	2	(50.0)	0	...	0	...
Vermont	5	0	...	0	...	0	...	2	(100.0)	0	...	3	(100.0)	0	...	0	...
Virginia	242	42	(92.9)	0	...	93	(81.7)	77	(90.9)	0	...	29	(89.7)	1	(100.0)	0	...
Washington	205	26	(92.3)	8	(100.0)	92	(84.8)	34	(85.3)	15	(86.7)	25	(76.0)	3	(100.0)	2	(100.0)
West Virginia	12	0	...	0	...	0	...	4	(100.0)	0	...	8	(100.0)	0	...	0	...
Wisconsin	45	9	(77.8)	1	(0.0)	13	(92.3)	13	(84.6)	0	...	9	(100.0)	0	...	0	...
Wyoming	6	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
American Samoa[6]	3	0	...	0	...	1	(0.0)	0	...	2	(50.0)	0	...	0	...	0	...
Fed. States of Micronesia[6]	162	1	(100.0)	0	...	0	...	0	...	155	(91.6)	0	...	2	(100.0)	4	(100.0)
Guam[6]	87	1	(100.0)	0	...	24	(100.0)	0	...	58	(91.4)	2	(100.0)	0	...	2	(100.0)
Marshall Islands[6]	191	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
N. Mariana Islands[6]	25	0	...	0	...	8	(87.5)	0	...	12	(100.0)	0	...	0	...	5	(100.0)
Puerto Rico[6]	70	67	(92.5)	0	...	0	...	0	...	0	...	2	(50.0)	0	...	1	(100.0)
Republic of Palau[6]	15	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
U.S. Virgin Islands[6]	.	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--

[1] Percentages shown only for reporting areas with information reported for ≥90% of cases, and indicate the percentage of those who completed therapy within 1 year.
[2] Most recent year for which data are available.
[3] Therapy < 1 year indicated in persons alive at diagnosis with an initial regimen of one or more drugs prescribed, who did not die during therapy. Excludes persons with initial isolate rifampin resistant, or patient with meningeal disease, or pediatric patient (aged <15) with miliary disease or positive blood culture, and those who moved out of country during treatment.
[4] Persons of Hispanic or Latino origin may be of any race.
[5] Excludes New York City.
[6] Not included in U.S. totals.
Note: Case counts and percentage for race categories do not include persons of Hispanic ethnicity. Ellipses indicate data not available. See Technical Notes for description of Completion of Therapy calculation (page 9).