Table 7. Tuberculosis Cases and Percentages by Case Verification Criterion and Site of Disease: United States, 1993–2012

Year	Total Cases	Verification Criterion[1]										Site of Disease[5]			
		Positive Culture		Positive NAA[2]		Positive Smear		Clinical Case Definition		Provider Diagnosis		Pulmonary[3]		Extra-pulmonary[4]	
		No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)
1993	25103	20307	(81)	0	(0)	185	(1)	3088	(12)	1523	(6)	21154	(84)	3940	(16)
1994	24205	19506	(81)	0	(0)	189	(1)	2916	(12)	1594	(7)	20318	(84)	3885	(16)
1995	22727	18266	(80)	0	(0)	189	(1)	2748	(12)	1524	(7)	18887	(83)	3835	(17)
1996	21210	17154	(81)	0	(0)	131	(1)	2607	(12)	1318	(6)	17387	(82)	3814	(18)
1997	19751	15979	(81)	0	(0)	155	(1)	2411	(12)	1206	(6)	16239	(82)	3509	(18)
1998	18287	14790	(81)	0	(0)	155	(1)	2253	(12)	1089	(6)	14801	(81)	3484	(19)
1999	17499	13994	(80)	0	(0)	172	(1)	2103	(12)	1230	(7)	14065	(80)	3431	(20)
2000	16309	13013	(80)	0	(0)	148	(1)	1951	(12)	1197	(7)	13086	(80)	3211	(20)
2001	15945	12750	(80)	0	(0)	123	(1)	1886	(12)	1186	(7)	12724	(80)	3217	(20)
2002	15055	11974	(80)	0	(0)	104	(1)	1822	(12)	1155	(8)	11901	(79)	3148	(21)
2003	14835	11683	(79)	0	(0)	116	(1)	1783	(12)	1253	(8)	11805	(80)	3020	(20)
2004	14498	11326	(78)	0	(0)	80	(1)	1824	(13)	1268	(9)	11523	(80)	2971	(20)
2005	14061	10955	(78)	0	(0)	96	(1)	1797	(13)	1213	(9)	11122	(79)	2933	(21)
2006	13727	10744	(78)	0	(0)	93	(1)	1629	(12)	1261	(9)	10852	(79)	2872	(21)
2007	13282	10426	(78)	0	(0)	69	(1)	1496	(11)	1291	(10)	10592	(80)	2687	(20)
2008	12895	10024	(78)	18	(0)	60	(0)	1548	(12)	1245	(10)	10238	(79)	2653	(21)
2009	11520	8886	(77)	56	(0)	74	(1)	1776	(15)	728	(6)	9010	(78)	2494	(22)
2010	11163	8454	(76)	109	(1)	71	(1)	1875	(17)	654	(6)	8728	(78)	2426	(22)
2011	10517	8084	(77)	129	(1)	57	(1)	1680	(16)	567	(5)	8330	(79)	2182	(21)
2012	9945	7597	(76)	151	(2)	43	(0)	1611	(16)	543	(5)	7824	(79)	2100	(21)

[1] Based on the public health surveillance case definition for tuberculosis; see Appendix A .
[2] Nucleic Acid Amplification test
[3] Includes all cases among persons with pulmonary as the only site of disease, and persons with both pulmonary and extrapulmonary sites of disease.
[4] Includes cases among persons with extrapulmonary TB disease only.
[5] Excludes missing and unknowns.
Note: See Technical Notes.
Data for all years updated through June 10, 2013.