

**Table 27. Tuberculosis Cases and Percentages by Reason Tuberculosis Therapy Stopped and Type of Move: United States, 2011**

Type of Move	Total Cases	Completed Therapy		Adverse Event		Lost		Refused		Died		Other <sup>1</sup>		Unknown	
	No.	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)
Moved in state <sup>2</sup>	336	258	(76.8)	2	(0.6)	7	(2.1)	5	(1.5)	24	(7.1)	34	(10.1)	6	(1.8)
Moved out of state <sup>3</sup>	284	213	(75.0)	0	(0.0)	17	(6.0)	2	(0.7)	6	(2.1)	38	(13.4)	8	(2.8)
Moved out of country <sup>4</sup>	397	165	(41.6)	0	(0.0)	22	(5.5)	0	(0.0)	7	(1.8)	193	(48.6)	10	(2.5)
Did not move <sup>5</sup>	9134	8211	(89.9)	26	(0.3)	72	(0.8)	61	(0.7)	645	(7.1)	28	(0.3)	91	(1.0)

<sup>1</sup> Therapy was discontinued for a known reason other than those listed (e.g. patient moved outside the U.S., or patient moved from state A to state B, and though state A notified state B, state B never followed up).

<sup>2</sup> Includes patients who were alive at diagnosis, started on treatment, and moved in state.

<sup>3</sup> Includes patients who were alive at diagnosis, started on treatment, and moved out of state.

<sup>4</sup> Includes patients who were alive at diagnosis, started on treatment, and moved out of the country; transnational referrals were provided for 264 (66.7%) TB patients who moved out of the country.

<sup>5</sup> Includes patients who were alive at diagnosis, started on treatment, and did not indicate having moved.

**Note:** There may be differences in the way jurisdictions determine treatment completion for patients who moved out of the country; some reporting jurisdictions may be classifying all patients who moved out of the country as 'other' for reason therapy stopped.