

Table 28. Deaths Prior to Tuberculosis Diagnosis or During Tuberculosis Therapy by Age Group: United States, 2011

Age Group	Total			Dead at Diagnosis						Died During Therapy ¹									
	Total Deaths Reported	Deaths Related to TB Disease or Therapy ²		Total Dead at Diagnosis	TB a Cause of Death		TB Not a Cause of Death		Unknown		Total Died During Therapy	Related to TB ³			Unrelated to TB		Unknown		
	No.	No.	(%)	No.	No.	(%)	No.	(%)	No.	(%)	No.	No.	(%)	No.	(%)	No.	(%)	No.	(%)
Total	928	337	(36.3)	244	88	(36.1)	104	(42.6)	52	(21.3)	684	249	(36.4)	307	(44.9)	128	(18.7)		
0-4	5	4	(80.0)	3	2	(66.7)	1	(33.3)	0	(0.0)	2	2	(100.0)	0	(0.0)	0	(0.0)	0	(0.0)
5-14	1	1	(100.0)	1	1	(100.0)	0	(0.0)	0	(0.0)	0	0	...	0	...	0	...	0	...
15-24	13	6	(46.2)	5	2	(40.0)	2	(40.0)	1	(20.0)	8	4	(50.0)	2	(25.0)	2	(25.0)	2	(25.0)
25-44	99	49	(49.5)	33	14	(42.4)	10	(30.3)	9	(27.3)	66	35	(53.0)	23	(34.8)	8	(12.1)	8	(12.1)
45-64	298	103	(34.6)	76	26	(34.2)	34	(44.7)	16	(21.1)	222	77	(34.7)	104	(46.8)	41	(18.5)	41	(18.5)
65+	512	174	(34.0)	126	43	(34.1)	57	(45.2)	26	(20.6)	386	131	(33.9)	178	(46.1)	77	(19.9)	77	(19.9)

¹ Among patients alive at diagnosis. Excludes 16 patients who died during therapy but did not start on therapy or unknown whether or not therapy was started.

² Includes patients who were dead at diagnosis or died during therapy, for which TB or TB therapy was indicated as a cause of death.

³ Six patient deaths during therapy were related to TB therapy.

Note: Ellipses indicate data not available.