

Table 15. Genotyped Tuberculosis Cases with *Mycobacterium bovis*¹ by Origin of Birth: United States, 2004–2014

Year	Total Genotyped Cases	<i>Mycobacterium bovis</i> cases					
		Total		U.S.-born		Foreign-born	
	No.	No. ²	(%)	No.	(%) ³	No.	(%) ³
2004	5,956	73	(1.2)	23	(31.5)	50	(68.5)
2005	7,500	81	(1.1)	23	(28.4)	58	(71.6)
2006	7,528	116	(1.5)	25	(21.6)	90	(77.6)
2007	8,431	112	(1.3)	17	(15.2)	94	(83.9)
2008	8,180	129	(1.6)	29	(22.5)	100	(77.5)
2009	7,714	113	(1.5)	27	(23.9)	86	(76.1)
2010	7,749	108	(1.4)	20	(18.5)	88	(81.5)
2011	7,616	117	(1.5)	30	(25.6)	87	(74.4)
2012	7,226	109	(1.5)	18	(16.5)	91	(83.5)
2013	7,044	96	(1.4)	22	(22.9)	74	(77.1)
2014	6,887	107	(1.6)	24	(22.4)	83	(77.6)

¹ *M. bovis* cases were defined predominantly by spoligotyping results with missing spacers 3, 9, 16, and 39-43. Data exclude 107 cases of Bacillus Calmette-Guérin (BCG) *M. bovis*, which have x, y or z in the second MIRU position. Please see technical notes for details regarding changes in the *M. bovis* definition, which result in a 13.9% reduction in the average of annual *M. bovis* case counts (n=105) compared to average annual counts that would have been reported without these changes (n=122).

² This column reports all genotyped *M. bovis* cases, including those where origin of birth is unknown.

³ Denominator is all *M. bovis* cases.