

Table 28. Deaths Among Reported Tuberculosis Cases, by Age Group: United States, 2013

Age group	Total			Dead at diagnosis						Died after diagnosis									
	Total deaths reported	Deaths related to TB disease or therapy ¹		Total dead at TB diagnosis		TB a cause of death		TB not a cause of death		Cause of death unknown/missing		Total died during therapy		Related to TB therapy/disease ²		Unrelated to TB therapy/disease		Cause of death unknown/missing	
	No.	No.	(%)	No.	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	No.	(%)	No.	(%)	No.	(%)
Total	807	285	(35.3)	211	60	(28.4)	111	(52.6)	40	(19.0)	596	225	(37.8)	288	(48.3)	83	(13.9)		
0–4	5	4	(80.0)	3	2	(66.7)	1	(33.3)	0	(0)	2	2	(100.0)	0	(0)	0	(0)		
5–14	0	0	...	0	0	...	0	...	0	...	0	0	...	0	...	0	...		
15–24	8	5	(62.5)	2	2	(100.0)	0	(0)	0	(0)	6	3	(50.0)	2	(33.3)	1	(16.7)		
25–44	61	30	(49.2)	24	13	(54.2)	5	(20.8)	6	(25.0)	37	17	(45.9)	15	(40.5)	5	(13.5)		
45–64	234	91	(38.9)	71	23	(32.4)	38	(53.5)	10	(14.1)	163	68	(41.7)	79	(48.5)	16	(9.8)		
≥65	499	155	(31.1)	111	20	(18.0)	67	(60.4)	24	(21.6)	388	135	(34.8)	192	(49.5)	61	(15.7)		

¹Includes patients who were dead at diagnosis or died during therapy, for which TB or TB therapy was indicated as a cause of death.

²Eight patient deaths during therapy were related to TB therapy.

Note: Ellipses (...) indicate that the percentage cannot be calculated and the denominator is 0.