

**Table 46. Tuberculosis Cases and Percentages, by Noninjecting Drug Use,<sup>1</sup> Ages ≥15 Years: Reporting Areas, 2015**

Reporting area	Total cases	Cases with information on noninjecting drug use		Cases reporting noninjecting drug use	
		No.	(%)	No.	(%)
<b>United States</b>	<b>9,115</b>	<b>9,015</b>	<b>(98.9)</b>	<b>653</b>	<b>(7.2)</b>
Alabama	116	116	(100.0)	17	(14.7)
Alaska	55	52	(94.5)	12	(23.1)
Arizona	184	181	(98.4)	23	(12.7)
Arkansas	78	78	(100.0)	8	(10.3)
California	2,065	2,033	(98.5)	111	(5.5)
Colorado	66	66	(100.0)	1	(1.5)
Connecticut	70	70	(100.0)	2	(2.9)
Delaware	19	19	(100.0)	0	(0)
District of Columbia	31	31	(100.0)	1	(3.2)
Florida	581	574	(98.8)	53	(9.2)
Georgia	306	300	(98.0)	28	(9.3)
Hawaii	119	119	(100.0)	3	(2.5)
Idaho	9	9	(100.0)	0	(0)
Illinois	330	329	(99.7)	24	(7.3)
Indiana	112	112	(100.0)	14	(12.5)
Iowa	36	36	(100.0)	0	(0)
Kansas	35	35	(100.0)	4	(11.4)
Kentucky	64	64	(100.0)	9	(14.1)
Louisiana	114	111	(97.4)	14	(12.6)
Maine	16	16	(100.0)	0	(0)
Maryland	169	168	(99.4)	5	(3.0)
Massachusetts	185	184	(99.5)	5	(2.7)
Michigan	128	127	(99.2)	11	(8.7)
Minnesota	140	140	(100.0)	2	(1.4)
Mississippi	71	70	(98.6)	14	(20.0)
Missouri	84	83	(98.8)	4	(4.8)
Montana	9	9	(100.0)	1	(11.1)
Nebraska	31	27	(87.1)	1	(3.7)
Nevada	84	84	(100.0)	4	(4.8)
New Hampshire	12	12	(100.0)	0	(0)
New Jersey	316	313	(99.1)	9	(2.9)
New Mexico	46	43	(93.5)	4	(9.3)
New York State <sup>2</sup>	173	169	(97.7)	2	(1.2)
New York City	560	557	(99.5)	31	(5.6)
North Carolina	189	189	(100.0)	33	(17.5)
North Dakota	9	8	(88.9)	0	(0)
Ohio	138	138	(100.0)	7	(5.1)
Oklahoma	62	61	(98.4)	7	(11.5)
Oregon	72	71	(98.6)	7	(9.9)
Pennsylvania	196	194	(99.0)	9	(4.6)
Rhode Island	30	27	(90.0)	2	(7.4)
South Carolina	98	96	(98.0)	10	(10.4)
South Dakota	16	15	(93.8)	2	(13.3)
Tennessee	125	124	(99.2)	14	(11.3)
Texas	1,250	1,250	(100.0)	131	(10.5)
Utah	30	30	(100.0)	0	(0)
Vermont	7	7	(100.0)	1	(14.3)
Virginia	202	202	(100.0)	4	(2.0)
Washington	196	185	(94.4)	4	(2.2)
West Virginia	9	9	(100.0)	0	(0)
Wisconsin	68	68	(100.0)	5	(7.4)
Wyoming	4	4	(100.0)	0	(0)
American Samoa <sup>3</sup>	4	4	(100.0)	0	(0)
Fed. States of Micronesia <sup>3</sup>	82	82	(100.0)	3	(3.7)
Guam <sup>3</sup>	60	58	(96.7)	0	(0)
Marshall Islands <sup>3</sup>	108	108	(100.0)	0	(0)
N. Mariana Islands <sup>3</sup>	24	24	(100.0)	0	(0)
Puerto Rico <sup>3</sup>	52	52	(100.0)	12	(23.1)
Republic of Palau <sup>3</sup>	13	13	(100.0)	0	(0)
U.S. Virgin Islands <sup>3</sup>	0	0	...	0	...

<sup>1</sup>Noninjecting drug use within past 12 months of TB diagnosis. Percentage based on 52 reporting areas (50 states, New York City, and the District of Columbia). Counts and percentages shown only for reporting areas with information reported for ≥75% of cases.

<sup>2</sup>Excludes New York City.

<sup>3</sup>Not included in U.S. totals.

**Note:** Ellipses (...) indicate that the percentage cannot be calculated, and the denominator is 0.