

Table 49. Tuberculosis Cases and Percentages, by Initial Drug Regimen: Reporting Areas, 2015

Reporting area	Total Cases	Cases in persons alive at diagnosis	Cases with information on initial drug regimen ¹		Percentage of cases in persons with initial drug regimen ^{2,3}		
			No.	(%)	IR	IRZ	IRZE ³
United States	9,557	9,344	9,274	(99.3)	(0.5)	(2.0)	(85.1)
Alabama	119	117	116	(99.1)	(0)	(2.6)	(81.0)
Alaska	68	63	61	(96.8)	(1.6)	(6.6)	(73.8)
Arizona	198	193	193	(100.0)	(0.5)	(1.6)	(93.8)
Arkansas	90	87	87	(100.0)	(0)	(4.6)	(87.4)
California	2,133	2,091	2,068	(98.9)	(0.1)	(0.9)	(90.2)
Colorado	73	72	72	(100.0)	(0)	(8.3)	(66.7)
Connecticut	70	70	70	(100.0)	(0)	(0)	(18.6)
Delaware	22	21	20	(95.2)	(0)	(0)	(90.0)
District of Columbia	33	30	30	(100.0)	(0)	(0)	(93.3)
Florida	602	583	583	(100.0)	(0.2)	(2.9)	(88.7)
Georgia	324	316	314	(99.4)	(0)	(1.3)	(33.8)
Hawaii	127	126	126	(100.0)	(0)	(3.2)	(92.9)
Idaho	11	11	11	(100.0)	(0)	(0)	(90.9)
Illinois	343	336	333	(99.1)	(1.2)	(3.9)	(86.2)
Indiana	116	114	114	(100.0)	(0)	(0.9)	(94.7)
Iowa	38	38	38	(100.0)	(0)	(2.6)	(97.4)
Kansas	36	35	35	(100.0)	(0)	(0)	(85.7)
Kentucky	67	67	67	(100.0)	(0)	(0)	(89.6)
Louisiana	119	117	115	(98.3)	(0)	(2.6)	(93.9)
Maine	18	17	17	(100.0)	(5.9)	(5.9)	(76.5)
Maryland	176	174	171	(98.3)	(0.6)	(4.1)	(88.3)
Massachusetts	192	190	187	(98.4)	(2.7)	(0.5)	(70.1)
Michigan	131	130	129	(99.2)	(0.8)	(3.9)	(58.1)
Minnesota	150	148	146	(98.6)	(0)	(4.8)	(87.0)
Mississippi	74	72	70	(97.2)	(2.9)	(5.7)	(78.6)
Missouri	92	90	90	(100.0)	(0)	(1.1)	(81.1)
Montana	9	9	8	(88.9)	(0)	(0)	(87.5)
Nebraska	33	33	32	(97.0)	(0)	(3.1)	(71.9)
Nevada	85	82	82	(100.0)	(1.2)	(0)	(98.8)
New Hampshire	13	12	12	(100.0)	(0)	(8.3)	(83.3)
New Jersey	326	316	312	(98.7)	(2.6)	(2.6)	(80.1)
New Mexico	47	42	41	(97.6)	(0)	(4.9)	(90.2)
New York State ⁴	188	186	184	(98.9)	(0)	(5.4)	(84.8)
New York City	577	566	566	(100.0)	(0.4)	(1.6)	(85.3)
North Carolina	199	193	192	(99.5)	(0)	(0.5)	(82.3)
North Dakota	9	9	9	(100.0)	(0)	(11.1)	(77.8)
Ohio	143	139	139	(100.0)	(1.4)	(3.6)	(86.3)
Oklahoma	67	64	63	(98.4)	(6.3)	(3.2)	(77.8)
Oregon	76	75	75	(100.0)	(0)	(1.3)	(90.7)
Pennsylvania	200	197	195	(99.0)	(0)	(0)	(84.1)
Rhode Island	30	29	28	(96.6)	(0)	(0)	(89.3)
South Carolina	104	100	100	(100.0)	(1.0)	(1.0)	(88.0)
South Dakota	17	17	17	(100.0)	(0)	(5.9)	(70.6)
Tennessee	131	128	128	(100.0)	(0)	(1.6)	(86.7)
Texas	1,334	1,311	1,304	(99.5)	(0.4)	(1.6)	(92.6)
Utah	37	37	37	(100.0)	(0)	(5.4)	(83.8)
Vermont	7	7	7	(100.0)	(0)	(14.3)	(71.4)
Virginia	212	208	208	(100.0)	(0.5)	(0)	(97.1)
Washington	208	198	194	(98.0)	(0)	(1.0)	(80.4)
West Virginia	10	8	8	(100.0)	(0)	(12.5)	(75.0)
Wisconsin	69	66	66	(100.0)	(0)	(7.6)	(84.8)
Wyoming	4	4	4	(100.0)	(0)	(0)	(100.0)
American Samoa ⁵	4	4	4	(100.0)	(0)	(0)	(50.0)
Fed. States of Micronesia ⁵	103	103	101	(98.1)	(0)	(1.0)	(96.0)
Guam ⁵	76	74	73	(98.6)	(0)	(0)	(91.8)
Marshall Islands ⁵	137	132	132	(100.0)	(0)	(0)	(94.7)
N. Mariana Islands ⁵	27	27	27	(100.0)	(0)	(0)	(18.5)
Puerto Rico ⁵	52	49	48	(98.0)	(0)	(0)	(100.0)
Republic of Palau ⁵	14	14	14	(100.0)	(0)	(0)	(100.0)
U.S. Virgin Islands ⁵	0	0	0

¹Includes persons who were alive at diagnosis and started on one or more drug.

²Overall U.S. percentage based on 52 reporting areas (50 states, New York City, and the District of Columbia). Counts and percentages shown only for reporting areas with information reported for ≥75% of cases.

³I, isoniazid; R, rifampin; Z, pyrazinamide; E, ethambutol. Cases with other drugs prescribed in addition to these regimens are excluded.

⁴Excludes New York City.

⁵Not included in U.S. totals.

Note: Excluding cases with no information on drug regimen, 45 (0.48%) persons were not started on any drugs, 11 (0.12%) were started on one drug, and 1,146 (12.30%) had an initial multiple drug regimen other than IR, IRZ, or IRZE.

Ellipses (...) indicate that the percentage cannot be calculated, and the denominator is 0.