

Table 60. Tuberculosis Cases and Percentages, by Completion of Tuberculosis Therapy (COT): Reporting Areas, 2013¹

Reporting area	Total cases	Therapy ≤1 year indicated ^{2,3,4}			Therapy >1 year indicated ^{3,5}	
		No.	COT ≤1 year(%)	COT(%)	No.	COT(%)
United States	9,550	7,774	(89.6)	(95.9)	567	(92.8)
Alabama	108	94	(95.7)	(98.9)	1	(100.0)
Alaska	71	67	(92.5)	(98.5)	2	(100.0)
Arizona	184	131	(86.3)	(95.4)	8	(100.0)
Arkansas	72	61	(72.1)	(82.0)	3	(100.0)
California	2,164	1758	(88.5)	(95.5)	102	(89.2)
Colorado	74	58	(94.8)	(98.3)	4	(100.0)
Connecticut	62	49	(85.7)	(91.8)	5	(100.0)
Delaware	18	17	—	—	1	—
District of Columbia	37	27	(85.2)	(100.0)	4	(100.0)
Florida	651	547	(93.6)	(97.6)	33	(97.0)
Georgia	339	290	(88.6)	(94.1)	17	(100.0)
Hawaii	115	102	(90.2)	(100.0)	7	(100.0)
Idaho	11	9	(88.9)	(100.0)	1	(100.0)
Illinois	327	267	(89.5)	(97.0)	14	(100.0)
Indiana	94	77	(96.1)	(100.0)	7	(85.7)
Iowa	47	41	(87.8)	(97.6)	2	(100.0)
Kansas	36	30	(100.0)	(100.0)	2	(50.0)
Kentucky	59	47	(91.5)	(97.9)	3	(100.0)
Louisiana	139	120	(74.2)	(87.5)	9	(77.8)
Maine	15	12	(91.7)	(100.0)	0	...
Maryland	175	149	(91.3)	(100.0)	9	(100.0)
Massachusetts	201	149	(91.9)	(93.3)	24	(87.5)
Michigan	141	107	(95.3)	(98.1)	13	(100.0)
Minnesota	151	125	(94.4)	(98.4)	18	(88.9)
Mississippi	65	42	(90.5)	(97.6)	4	(100.0)
Missouri	104	84	—	—	7	—
Montana	6	4	—	—	0	...
Nebraska	21	20	(95.0)	(95.0)	0	...
Nevada	91	78	(96.2)	(100.0)	4	(100.0)
New Hampshire	15	12	(83.3)	(91.7)	1	(100.0)
New Jersey	319	252	(92.1)	(97.2)	30	(86.7)
New Mexico	50	34	(94.1)	(97.1)	3	(100.0)
New York State ⁶	215	172	(89.0)	(97.1)	23	(91.3)
New York City	650	544	(95.2)	(97.6)	55	(96.4)
North Carolina	216	178	(97.8)	(99.4)	18	(100.0)
North Dakota	12	8	(75.0)	(100.0)	1	(100.0)
Ohio	148	118	(83.9)	(95.8)	10	(100.0)
Oklahoma	70	58	(93.1)	(100.0)	2	(100.0)
Oregon	73	61	(86.9)	(96.7)	5	(80.0)
Pennsylvania	214	176	(76.1)	(93.2)	9	(100.0)
Rhode Island	27	23	(95.7)	(95.7)	2	(100.0)
South Carolina	112	92	(84.8)	(91.3)	8	(100.0)
South Dakota	9	6	(83.3)	(100.0)	1	(0)
Tennessee	143	118	(96.6)	(100.0)	4	(100.0)
Texas	1,209	951	(88.2)	(94.4)	64	(93.8)
Utah	33	28	(100.0)	(100.0)	1	(100.0)
Vermont	5	4	(100.0)	(100.0)	1	(0)
Virginia	179	148	(94.6)	(98.6)	10	(100.0)
Washington	210	178	(84.8)	(93.8)	9	(77.8)
West Virginia	13	9	(100.0)	(100.0)	1	(0)
Wisconsin	50	42	(90.5)	(100.0)	5	(80.0)
Wyoming	0	0	—	—	0	...
American Samoa ⁷	2	2	(100.0)	(100.0)	0	...
Fed. States of Micronesia ⁷	128	117	(93.2)	(94.0)	4	(100.0)
Guam ⁷	48	44	(95.5)	(100.0)	1	(100.0)
Marshall Islands ⁷	153	137	(90.5)	(93.4)	3	(100.0)
N. Mariana Islands ⁷	31	0	—	—	0	...
Puerto Rico ⁷	49	37	(91.9)	(97.3)	1	(100.0)
Republic of Palau ⁷	7	7	(100.0)	(100.0)	0	...
U.S. Virgin Islands ⁷	2	2	—	—	0	...

¹Most recent year for which data are available.

²Initial isolate susceptible to rifampin (n = 5,755) or susceptibility unknown (n = 111); culture negative (n = 1,647); culture status unknown (n = 261).

³Number of cases in persons alive at diagnosis, with an initial regimen of one or more drugs prescribed. Percentage for U.S. based on 52 reporting areas (50 states, New York City, and the District of Columbia). Percentages shown only for reporting areas with information reported for ≥90% of cases.

⁴Therapy ≤1 year indicated in persons alive at diagnosis with an initial regimen of one or more drugs prescribed, who did not die within one year of initiating therapy. Excludes persons with initial isolate rifampin resistant, or patient with bone and joint disease, meningeal disease or disease of the central nervous system, or pediatric patient (age <15) with military disease or positive blood culture or a positive nucleic acid amplification test on a blood specimen, and those who moved out of the country within one year of initiating treatment.

⁵Initial isolate rifampin resistant, or patient with meningeal disease or bone and joint disease, or disease of the central nervous system, or pediatric patient (age <15) with military disease or positive blood culture or a positive nucleic acid amplification test on a blood specimen, and those who did not move out of the country or die during treatment.

⁶Excludes New York City.

⁷Not included in U.S. totals.

Note: Ellipses (...) indicate that the percentage cannot be calculated, and the denominator is 0. See Technical Notes for description of completion of therapy calculation.