

Table 12. Tuberculosis Cases and Percentages, by Reason Tuberculosis Therapy Was Stopped: United States, 1993–2013

| Year | Total cases ¹ | Completed therapy | | Adverse event | | Moved ² | | Lost | | Refused | | Died ³ | | Unknown ⁴ | |
|------|--------------------------|-------------------|--------|---------------|-------|--------------------|-------|-------|-------|---------|-------|-------------------|--------|----------------------|-------|
| | No. | No. | (%) | No. | (%) | No. | (%) | No. | (%) | No. | (%) | No. | (%) | No. | (%) |
| 1993 | 23,740 | 18,043 | (76.0) | 0 | (0) | 1,120 | (4.7) | 1,086 | (4.6) | 223 | (0.9) | 3,053 | (12.9) | 215 | (0.9) |
| 1994 | 23,052 | 17,764 | (77.1) | 0 | (0) | 1,194 | (5.2) | 740 | (3.2) | 183 | (0.8) | 2,743 | (11.9) | 428 | (1.9) |
| 1995 | 21,705 | 17,306 | (79.7) | 0 | (0) | 969 | (4.5) | 570 | (2.6) | 155 | (0.7) | 2,396 | (11.0) | 309 | (1.4) |
| 1996 | 20,298 | 16,528 | (81.4) | 0 | (0) | 783 | (3.9) | 525 | (2.6) | 156 | (0.8) | 1,998 | (9.8) | 308 | (1.5) |
| 1997 | 18,930 | 15,673 | (82.8) | 0 | (0) | 667 | (3.5) | 444 | (2.3) | 119 | (0.6) | 1,755 | (9.3) | 272 | (1.4) |
| 1998 | 17,583 | 14,766 | (84.0) | 0 | (0) | 533 | (3.0) | 411 | (2.3) | 104 | (0.6) | 1,579 | (9.0) | 190 | (1.1) |
| 1999 | 16,861 | 14,234 | (84.4) | 0 | (0) | 456 | (2.7) | 359 | (2.1) | 104 | (0.6) | 1,437 | (8.5) | 271 | (1.6) |
| 2000 | 15,784 | 13,407 | (84.9) | 0 | (0) | 406 | (2.6) | 397 | (2.5) | 112 | (0.7) | 1,294 | (8.2) | 168 | (1.1) |
| 2001 | 15,409 | 13,242 | (85.9) | 0 | (0) | 378 | (2.5) | 402 | (2.6) | 99 | (0.6) | 1,121 | (7.3) | 167 | (1.1) |
| 2002 | 14,564 | 12,482 | (85.7) | 0 | (0) | 336 | (2.3) | 412 | (2.8) | 87 | (0.6) | 1,080 | (7.4) | 167 | (1.1) |
| 2003 | 14,379 | 12,418 | (86.4) | 0 | (0) | 313 | (2.2) | 390 | (2.7) | 84 | (0.6) | 994 | (6.9) | 180 | (1.3) |
| 2004 | 14,080 | 12,118 | (86.1) | 0 | (0) | 337 | (2.4) | 370 | (2.6) | 82 | (0.6) | 975 | (6.9) | 198 | (1.4) |
| 2005 | 13,673 | 11,727 | (85.8) | 1 | (0) | 323 | (2.4) | 338 | (2.5) | 90 | (0.7) | 985 | (7.2) | 209 | (1.5) |
| 2006 | 13,317 | 11,541 | (86.7) | 0 | (0) | 292 | (2.2) | 358 | (2.7) | 79 | (0.6) | 939 | (7.1) | 108 | (0.8) |
| 2007 | 12,907 | 11,348 | (87.9) | 0 | (0) | 241 | (1.9) | 327 | (2.5) | 73 | (0.6) | 819 | (6.3) | 99 | (0.8) |
| 2008 | 12,552 | 10,887 | (86.7) | 7 | (0.1) | 256 | (2.0) | 329 | (2.6) | 78 | (0.6) | 843 | (6.7) | 152 | (1.2) |
| 2009 | 11,185 | 9,832 | (87.9) | 22 | (0.2) | 96 | (0.9) | 165 | (1.5) | 82 | (0.7) | 682 | (6.1) | 306 | (2.7) |
| 2010 | 10,835 | 9,543 | (88.1) | 29 | (0.3) | — | — | 159 | (1.5) | 64 | (0.6) | 657 | (6.1) | 383 | (3.5) |
| 2011 | 10,209 | 8,982 | (88.0) | 28 | (0.3) | — | — | 127 | (1.2) | 69 | (0.7) | 688 | (6.7) | 315 | (3.1) |
| 2012 | 9,674 | 8,502 | (87.9) | 31 | (0.3) | — | — | 123 | (1.3) | 57 | (0.6) | 607 | (6.3) | 354 | (3.7) |
| 2013 | 9,266 | 8,127 | (87.7) | 40 | (0.4) | — | — | 97 | (1.0) | 66 | (0.7) | 579 | (6.2) | 357 | (3.9) |

¹Includes all cases in persons reported as alive at diagnosis and taking one or more TB drugs.

²In 2009 the “moved” response option was removed from the RVCT’s reason therapy was stopped variable; see Technical Notes for details.

³Died = died of any cause (not only TB).

⁴Includes cases in persons reporting reason therapy stopped = other, missing, unknown, or moved (from 2010).

Note: Data for all years are updated through June 9, 2016.

Data complete to 2013. See Technical Notes for details.