

Table 15. Genotyped Tuberculosis Cases with *Mycobacterium bovis*¹ by Origin of Birth: United States, 2004–2015

Year	Total genotyped cases	<i>Mycobacterium bovis</i> cases					
		Total		U.S.-born		Foreign-born	
	No.	No. ²	(%)	No.	(%) ³	No.	(%) ³
2004	5,954	73	(1.2)	23	(31.5)	50	(68.5)
2005	7,500	81	(1.1)	23	(28.4)	58	(71.6)
2006	7,529	116	(1.5)	25	(21.6)	90	(77.6)
2007	8,427	113	(1.3)	17	(15.0)	95	(84.1)
2008	8,179	129	(1.6)	29	(22.5)	100	(77.5)
2009	7,718	113	(1.5)	27	(23.9)	86	(76.1)
2010	7,749	108	(1.4)	20	(18.5)	88	(81.5)
2011	7,617	117	(1.5)	30	(25.6)	87	(74.4)
2012	7,234	109	(1.5)	18	(16.5)	91	(83.5)
2013	7,049	86	(1.2)	17	(19.8)	69	(80.2)
2014	6,981	107	(1.5)	24	(22.4)	83	(77.6)
2015	7,123	123	(1.7)	30	(24.4)	93	(75.6)

¹*M. bovis* cases were defined predominantly by spoligotyping results with missing spacers 3, 9, 16, and 39–43. Data exclude cases of Bacillus Calmette-Guérin (BCG) *M. bovis*, which have x, y or z in the second MIRU position.

²This column reports all genotyped *M. bovis* cases, including those where origin of birth is unknown.

³Denominator is all *M. bovis* cases.