

**Table 24. Tuberculosis Cases and Clusters, by Cluster Size<sup>1</sup>: United States, 2013–2015**

Cluster Size	Clusters		Clustered cases <sup>2</sup>	
	No.	(%) <sup>3</sup>	No.	(%) <sup>4</sup>
<b>Total</b>	<b>1,464</b>	<b>(100.0)</b>	<b>4,529</b>	<b>(100.0)</b>
2-case cluster	958	(65.4)	1,916	(42.3)
3-case cluster	233	(15.9)	699	(15.4)
4-case cluster	115	(7.9)	460	(10.2)
5-case cluster	48	(3.3)	240	(5.3)
6-case cluster	31	(2.1)	186	(4.1)
7-case cluster	14	(1.0)	98	(2.2)
8-case cluster	17	(1.2)	136	(3.0)
9-case cluster	10	(0.7)	90	(2.0)
≥10-case cluster	38	(2.6)	704	(15.5)

<sup>1</sup>Clusters have two or more cases with matching spoligotype and 24-locus mycobacterial interspersed repetitive unit-variable number tandem repeat type (GENType) within a county during the specified 3-year time period.

<sup>2</sup>Cases with matching spoligotype and 24-locus mycobacterial interspersed repetitive unit-variable number tandem repeat type (GENType) are members of a cluster within a county during the specified 3-year time period.

<sup>3</sup>Denominator is total number of clusters.

<sup>4</sup>Denominator is total number of cases.

See Surveillance Slide #35.