

**Table 27. Tuberculosis Cases and Percentages by Reason Tuberculosis Therapy Was Stopped and Type of Move: United States, 2013**

Type of move	Total cases	Completed therapy		Adverse event		Lost		Refused		Died		Other <sup>4</sup>		Unknown	
	No.	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)
Moved in state <sup>1</sup>	320	268	(83.8)	1	(0.3)	8	(2.5)	1	(0.3)	14	(4.4)	27	(8.4)	1	(0.3)
Moved out of state <sup>2</sup>	293	239	(81.6)	0	(0)	17	(5.8)	3	(1.0)	5	(1.7)	25	(8.5)	4	(1.4)
Moved out of country <sup>3</sup>	367	147	(40.1)	3	(0.8)	21	(5.7)	4	(1.1)	7	(1.9)	167	(45.5)	17	(4.6)
Did not move	8,033	7,310	(91.0)	35	(0.4)	52	(0.6)	54	(0.7)	536	(6.7)	26	(0.3)	20	(0.2)
Unknown if moved	273	168	(61.5)	1	(0.4)	1	(0.4)	4	(1.5)	18	(6.6)	6	(2.2)	75	(27.5)

<sup>1</sup>Includes patients who were alive at diagnosis, started on treatment, and moved in state.

<sup>2</sup>Includes patients who were alive at diagnosis, started on treatment, and moved out of state.

<sup>3</sup>Includes patients who were alive at diagnosis, started on treatment, and moved out of the country; transnational referrals were provided for 230 (62.7%) TB patients who moved out of the country.

<sup>4</sup>Therapy was discontinued for a known reason other than those listed (e.g., patient moved outside the United States, or patient moved from state A to state B, and although state A notified state B, state B never followed up).

**Note:** There may be differences in the way jurisdictions determine treatment completion for patients who moved out of the country; some reporting jurisdictions may be classifying all patients who moved out of the country as 'other' for reason therapy stopped.

Moving in and out of state or country is not mutually exclusive. Two patients moved twice, and one patient moved but did not indicate moving in state, out of state or out of country.