

Table 12. Tuberculosis Cases and Percentages, by Reason Tuberculosis Therapy Was Stopped: United States, 1993–2014

Year	Total cases ¹	Completed therapy		Adverse event		Moved ²		Lost		Refused		Died ³		Unknown ⁴	
	No.	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)
1993	23,740	18,043	(76.0)	0	(0)	1,120	(4.7)	1,086	(4.6)	223	(0.9)	3,053	(12.9)	215	(0.9)
1994	23,052	17,764	(77.1)	0	(0)	1,194	(5.2)	740	(3.2)	183	(0.8)	2,743	(11.9)	428	(1.9)
1995	21,705	17,306	(79.7)	0	(0)	969	(4.5)	570	(2.6)	155	(0.7)	2,396	(11.0)	309	(1.4)
1996	20,298	16,528	(81.4)	0	(0)	783	(3.9)	525	(2.6)	156	(0.8)	1,998	(9.8)	308	(1.5)
1997	18,930	15,673	(82.8)	0	(0)	667	(3.5)	444	(2.3)	119	(0.6)	1,755	(9.3)	272	(1.4)
1998	17,583	14,766	(84.0)	0	(0)	533	(3.0)	411	(2.3)	104	(0.6)	1,579	(9.0)	190	(1.1)
1999	16,861	14,234	(84.4)	0	(0)	456	(2.7)	359	(2.1)	104	(0.6)	1,437	(8.5)	271	(1.6)
2000	15,784	13,407	(84.9)	0	(0)	406	(2.6)	397	(2.5)	112	(0.7)	1,294	(8.2)	168	(1.1)
2001	15,409	13,242	(85.9)	0	(0)	378	(2.5)	402	(2.6)	99	(0.6)	1,121	(7.3)	167	(1.1)
2002	14,564	12,482	(85.7)	0	(0)	336	(2.3)	412	(2.8)	87	(0.6)	1,080	(7.4)	167	(1.1)
2003	14,379	12,418	(86.4)	0	(0)	313	(2.2)	390	(2.7)	84	(0.6)	994	(6.9)	180	(1.3)
2004	14,080	12,118	(86.1)	0	(0)	337	(2.4)	370	(2.6)	82	(0.6)	975	(6.9)	198	(1.4)
2005	13,677	11,727	(85.7)	1	(0)	323	(2.4)	340	(2.5)	90	(0.7)	986	(7.2)	210	(1.5)
2006	13,317	11,541	(86.7)	0	(0)	292	(2.2)	358	(2.7)	79	(0.6)	939	(7.1)	108	(0.8)
2007	12,906	11,347	(87.9)	0	(0)	241	(1.9)	327	(2.5)	73	(0.6)	819	(6.3)	99	(0.8)
2008	12,550	10,887	(86.7)	7	(0.1)	256	(2.0)	329	(2.6)	77	(0.6)	843	(6.7)	151	(1.2)
2009	11,182	9,829	(87.9)	22	(0.2)	96	(0.9)	165	(1.5)	82	(0.7)	682	(6.1)	306	(2.7)
2010	10,834	9,541	(88.1)	29	(0.3)	—	—	159	(1.5)	64	(0.6)	658	(6.1)	383	(3.5)
2011	10,210	8,988	(88.0)	28	(0.3)	—	—	127	(1.2)	70	(0.7)	688	(6.7)	309	(3.0)
2012	9,674	8,513	(88.0)	31	(0.3)	—	—	123	(1.3)	58	(0.6)	610	(6.3)	339	(3.5)
2013	9,282	8,173	(88.1)	40	(0.4)	—	—	99	(1.1)	68	(0.7)	584	(6.3)	318	(3.4)
2014	9,163	8,119	(88.6)	18	(0.2)	—	—	83	(0.9)	83	(0.9)	568	(6.2)	292	(3.2)

¹Includes all cases in persons reported as alive at diagnosis and taking one or more TB drugs.

²In 2009 the “moved” response option was removed from the RVCT’s reason therapy was stopped variable; see Technical Notes for details.

³Died = died of any cause (not only TB).

⁴Includes cases in persons reporting reason therapy stopped = other, missing, unknown, or moved (from 2010).

Note: Data for all years are updated through June 21, 2017.

Data complete to 2014. See Technical Notes for details.