

Table 27. Deaths Among Reported Tuberculosis Cases, by Age Group: United States, 2014

Age group	Total			Dead at diagnosis						Died after diagnosis							
	Total deaths reported	Deaths related to TB disease or therapy ¹		Total dead at TB diagnosis	TB a cause of death		TB not a cause of death		Cause of death unknown/missing		Total died during therapy	Related to TB therapy/disease ²		Unrelated to TB therapy/disease		Cause of death unknown/missing	
	No.	No.	(%)	No.	No.	(%)	No.	(%)	No.	(%)	No.	No.	(%)	No.	(%)	No.	(%)
Total	774	287	(37.1)	194	62	(32.0)	99	(51.0)	33	(17.0)	580	225	(38.8)	253	(43.6)	102	(17.6)
0–4	4	3	(75.0)	2	1	(50.0)	1	(50.0)	0	(0)	2	2	(100.0)	0	(0)	0	(0)
5–14	0	0	...	0	0	...	0	...	0	...	0	0	...	0	...	0	...
15–24	10	5	(50.0)	1	1	(100.0)	0	(0)	0	(0)	9	4	(44.4)	5	(55.6)	0	(0)
25–44	68	32	(47.1)	18	8	(44.4)	6	(33.3)	4	(22.2)	50	24	(48.0)	21	(42.0)	5	(10.0)
45–64	220	86	(39.1)	53	24	(45.3)	20	(37.7)	9	(17.0)	167	62	(37.1)	73	(43.7)	32	(19.2)
≥65	472	161	(34.1)	120	28	(23.3)	72	(60.0)	20	(16.7)	352	133	(37.8)	154	(43.8)	65	(18.5)

¹Includes patients who were dead at diagnosis or died during therapy, for which TB or TB therapy was indicated as a cause of death.

²Three patient deaths during therapy were related to TB therapy.

Note: Ellipses (...) indicate that the percentage cannot be calculated and the denominator is 0.