

Table 58. Tuberculosis Cases and Percentages, by Completion of Tuberculosis Therapy (COT): Reporting Areas, 2014¹

| Reporting area | Total cases | Therapy ≤1 year indicated ^{2,3,4} | | | Therapy >1 year indicated ^{3,5} | |
|--|--------------|--|----------------|---------------|--|---------------|
| | | No. | COT ≤1 year(%) | COT(%) | No. | COT(%) |
| United States | 9,398 | 7720 | (90.1) | (96.6) | 526 | (94.1) |
| Alabama | 133 | 112 | (93.8) | (99.1) | 6 | (100.0) |
| Alaska | 62 | 54 | (94.4) | (100.0) | 3 | (66.7) |
| Arizona | 193 | 130 | (86.9) | (96.2) | 11 | (100.0) |
| Arkansas | 93 | 80 | (76.3) | (91.3) | 4 | (100.0) |
| California | 2,130 | 1745 | (88.3) | (96.6) | 126 | (94.4) |
| Colorado | 64 | 52 | (96.2) | (100.0) | 4 | (100.0) |
| Connecticut | 60 | 51 | (98.0) | (100.0) | 6 | (100.0) |
| Delaware | 21 | 20 | (85.0) | (95.0) | 0 | ... |
| District of Columbia | 32 | 27 | (92.6) | (100.0) | 1 | (100.0) |
| Florida | 595 | 507 | (92.9) | (97.0) | 21 | (95.2) |
| Georgia | 334 | 282 | (89.4) | (96.1) | 11 | (100.0) |
| Hawaii | 136 | 121 | (98.3) | (99.2) | 6 | (83.3) |
| Idaho | 11 | 8 | (75.0) | (87.5) | 1 | (100.0) |
| Illinois | 319 | 262 | (89.7) | (97.7) | 22 | (95.5) |
| Indiana | 108 | 91 | (92.3) | (97.8) | 9 | (100.0) |
| Iowa | 54 | 43 | (90.7) | (95.3) | 4 | (100.0) |
| Kansas | 40 | 35 | (94.3) | (100.0) | 1 | (100.0) |
| Kentucky | 80 | 62 | (95.2) | (98.4) | 5 | (100.0) |
| Louisiana | 121 | 103 | (79.6) | (93.2) | 3 | (100.0) |
| Maine | 14 | 14 | (78.6) | (85.7) | 0 | ... |
| Maryland | 198 | 163 | (89.6) | (98.2) | 11 | (90.9) |
| Massachusetts | 199 | 163 | (89.0) | (96.9) | 22 | (100.0) |
| Michigan | 105 | 87 | (95.4) | (96.6) | 5 | (100.0) |
| Minnesota | 147 | 122 | (92.6) | (98.4) | 15 | (93.3) |
| Mississippi | 74 | 60 | (81.7) | (91.7) | 1 | (100.0) |
| Missouri | 79 | 65 | (69.2) | (80.0) | 5 | (100.0) |
| Montana | 6 | 3 | (100.0) | (100.0) | 0 | ... |
| Nebraska | 38 | 35 | (68.6) | (68.6) | 0 | ... |
| Nevada | 74 | 66 | (93.9) | (97.0) | 2 | (100.0) |
| New Hampshire | 11 | 10 | (100.0) | (100.0) | 1 | (100.0) |
| New Jersey | 307 | 244 | (92.6) | (97.5) | 23 | (82.6) |
| New Mexico | 50 | 35 | (94.3) | (97.1) | 1 | (100.0) |
| New York State ⁶ | 202 | 167 | (92.2) | (97.0) | 8 | (100.0) |
| New York City | 582 | 476 | (95.4) | (98.3) | 39 | (92.3) |
| North Carolina | 195 | 159 | (97.5) | (100.0) | 15 | (100.0) |
| North Dakota | 15 | 12 | (58.3) | (83.3) | 3 | (100.0) |
| Ohio | 156 | 121 | (90.9) | (96.7) | 12 | (83.3) |
| Oklahoma | 59 | 53 | (94.3) | (98.1) | 2 | (100.0) |
| Oregon | 77 | 58 | (94.8) | (98.3) | 2 | (100.0) |
| Pennsylvania | 208 | 166 | (80.1) | (94.6) | 13 | (92.3) |
| Rhode Island | 21 | 17 | (88.2) | (88.2) | 3 | (100.0) |
| South Carolina | 79 | 65 | (95.4) | (98.5) | 6 | (83.3) |
| South Dakota | 8 | 5 | (80.0) | (100.0) | 1 | (100.0) |
| Tennessee | 151 | 128 | (95.3) | (100.0) | 7 | (85.7) |
| Texas | 1,269 | 1025 | (87.7) | (95.0) | 60 | (93.3) |
| Utah | 31 | 23 | (100.0) | (100.0) | 4 | (75.0) |
| Vermont | 2 | 2 | (100.0) | (100.0) | 0 | ... |
| Virginia | 198 | 170 | (95.9) | (100.0) | 10 | (100.0) |
| Washington | 194 | 169 | (91.1) | (95.9) | 7 | (100.0) |
| West Virginia | 13 | 12 | (91.7) | (100.0) | 1 | (100.0) |
| Wisconsin | 48 | 38 | (89.5) | (100.0) | 3 | (66.7) |
| Wyoming | 2 | 2 | (100.0) | (100.0) | 0 | ... |
| American Samoa ⁷ | 1 | 0 | ... | ... | 1 | (100.0) |
| Fed. States of Micronesia ⁷ | 170 | 160 | (92.5) | (93.1) | 2 | (100.0) |
| Guam ⁷ | 56 | 51 | (94.1) | (96.1) | 2 | (50.0) |
| Marshall Islands ⁷ | 152 | 137 | (92.0) | (94.2) | 2 | (100.0) |
| N. Mariana Islands ⁷ | 25 | 10 | (100.0) | (100.0) | 1 | (100.0) |
| Puerto Rico ⁷ | 44 | 32 | (90.6) | (93.8) | 1 | (100.0) |
| Republic of Palau ⁷ | 14 | 7 | (100.0) | (100.0) | 2 | (50.0) |
| U.S. Virgin Islands ⁷ | 0 | — | — | — | — | ... |

¹Most recent year for which data are available.

²Initial isolate susceptible to rifampin (n = 5,746) or susceptibility unknown (n = 90); culture negative (n = 1,622); culture status unknown (n = 272).

³Number of cases in persons alive at diagnosis, with an initial regimen of one or more drugs prescribed. Percentage for U.S. based on 52 reporting areas (50 states, New York City, and the District of Columbia).

⁴Therapy ≤1 year indicated in persons alive at diagnosis with an initial regimen of one or more drugs prescribed, who did not die within one year of initiating therapy. Excludes persons with initial isolate rifampin resistant, or patient with bone and joint disease, meningeal disease or disease of the central nervous system, or pediatric patient (age <15) with miliary disease or positive blood culture or a positive nucleic acid amplification test on a blood specimen, and those who moved out of the country within one year of initiating treatment.

⁵Initial isolate rifampin resistant, or patient with meningeal disease or bone and joint disease, or disease of the central nervous system, or pediatric patient (age <15) with miliary disease or positive blood culture or a positive nucleic acid amplification test on a blood specimen, and those who did not move out of the country or die during treatment.

⁶Excludes New York City.

⁷Not included in U.S. totals.

Note: Ellipses (...) indicate that the percentage cannot be calculated, and the denominator is 0.

See Technical Notes for description of completion of therapy calculation.