



# TOBACCO USE

## Healthy People 2020 Leading Health Indicators (LHIs)

Goal: Reduce illness, disability, and death related to tobacco use and secondhand smoke exposure.

### TU-1.1 Reduce cigarette smoking by adults

BASELINE  
20.6%  
2008

17.9%  
2013



of **adults** aged 18 years and older were current cigarette smokers in 2013  
*(age adjusted to the year 2000 standard population)*

*Data Source: National Health Interview Survey*

TARGET  
12%  
YEAR 2020



### TU-2.2 Reduce use of cigarette by adolescents (past month)

BASELINE  
19.5%  
2008

15.7%  
2013



of **adolescents** in grades 9 through 12 smoked cigarettes in the past 30 days in 2013

*Data Source: Youth Risk Behavior Surveillance System*

TARGET  
16%  
YEAR 2020



### TU-11.1 Reduce the proportion of children aged 3 to 11 years exposed to secondhand smoke

BASELINE  
52.2%  
2005-2008

41.3%  
2009-2012



of **children** aged 3 to 11 years were exposed to secondhand smoke in 2009-2012

*Data Source: National Health and Nutrition Examination Survey*

TARGET  
47%  
YEAR 2020



For more information, please visit <http://www.cdc.gov/tobacco/hp2020>

Email Questions to: [hp2020tobacco@cdc.gov](mailto:hp2020tobacco@cdc.gov)



USE YOUR  
SMARTPHONE  
TO SCAN THIS  
QR CODE



Centers for Disease  
Control and Prevention  
National Center for Chronic  
Disease Prevention and  
Health Promotion