List of Tables and Figures

Chapter 1

Introduction and Summary of Conclusions	
Table 1.	U.S. population distribution, by race/ ethnicity and Hispanic origin, and percentage change, 1980–1990 <i>8</i>
Table 2.	Selected demographic characteristics for the U.S. population, by race/ethnicity, 1990 9
Table 3.	Estimated percentage distribution of the U.S. population, by race/ethnicity and Hispanic origin, 1990–2050 <i>10</i>

Chapter 2

Patterns of Tobacco Use Among Four Racial/Ethnic Minority Groups

- Table 1.Percentage of adult African Americans
who reported being current cigarette
smokers, overall and by gender, age, and
education, National Health Interview
Surveys, United States, 1978–1995
aggregate data 22
- Table 2.Percentage of adult African American
smokers who reported smoking <15,
15-24, or ≥ 25 cigarettes per day, overall
and by gender, age, and education,
National Health Interview Surveys,
United States, 1978–1995 aggregate
data 24
- Table 3.Percentage of adult African American
ever smokers who have quit, overall and
by gender, age, and education, National
Health Interview Surveys, United States,
1978–1995 aggregate data
- Table 4.Current cigarette smoking status among
persons who reported that they were
daily smokers 1 year before being
surveyed, Current Population Survey
National Cancer Institute Supplement,
1992–1993 27

- Table 5.Percentage of African American women
of reproductive age who reported being
current cigarette smokers, overall and by
education, National Health Interview
Surveys, United States, 1978–1995
aggregate data 28
- Table 6.Percentage of live-born infants' mothers
who reported smoking during preg-
nancy, by year and race/ethnicity, U.S.
final natality statistics, 1989–1995
- Table 7.Trends in the percentage of high school
seniors who were previous-month
smokers, by race/ethnicity and gender,
Monitoring the Future surveys, United
States, 1976–1979, 1980–1984, 1985–1989,
1990–1994 32
- Table 8.Percentage of African Americans and
whites 20–34 years of age who reported
being current cigarette smokers, by age
group and gender, National Health
Interview Surveys, United States,
1978–1995 aggregate data 33
- Table 9.Percentage of African Americans and
whites 20–24 years of age who reported
being current cigarette smokers, by
education and gender, National Health
Interview Surveys, United States,
1978–1995 aggregate data 34
- Table 10.Percentage of African American and
white high school seniors who reported
recently using or not using cigarettes
and other selected substances, Monitor-
ing the Future surveys, United States,
1976–1994 aggregate data 37
- Table 11.Cumulative percentages of recalled age
at which a respondent first tried a
cigarette and began smoking daily,
among African American, Hispanic, and
white men and women aged 30–39,
National Household Surveys on Drug
Abuse, United States, 1994–1995

- Table 12.Percentage of African American and
white high school seniors who reported
previous-month and heavy smoking, by
selected variables, Monitoring the
Future surveys, United States,
1976–1994 42
- Table 13. Percentage of American Indian and Alaska Native adults who reported being current cigarette smokers, overall and by gender, age, and education, National Health Interview Surveys, United States, 1978–1995 aggregate data 46
- Table 14.Percentage of adult American Indian
and Alaska Native smokers who re-
ported smoking <15, 15–24, or ≥ 25
cigarettes per day, overall and by
gender, age, and education, National
Health Interview Surveys, United States,
1978–1995 aggregate data 47
- Table 15.Percentage of adult American Indian
and Alaska Native ever smokers who
have quit, overall and by gender, age,
and education, National Health Inter-
view Surveys, United States, 1978–1995
aggregate data 48
- Table 16.Percentage of American Indian and
Alaska Native women of reproductive
age who reported being current cigarette
smokers, overall and by education,
National Health Interview Surveys,
United States, 1978–1995 aggregate
data 49
- Table 17.Percentage of American Indian and
Alaska Native adults who reported
being current cigarette smokers, overall
and by region/state, gender, age, and
education, Behavioral Risk Factor
Surveillance System, 1988–1992
aggregate data 50
- Table 18.Percentage of adult American Indian
and Alaska Native smokers who re-
ported smoking <15, 15–24, or ≥ 25
cigarettes per day, overall and by
region/state, gender, age, and educa-
tion, Behavioral Risk Factor Surveillance
System, 1988–1992 aggregate data 52

- Table 19.Percentage of adult American Indian
and Alaska Native smokers who re-
ported they quit smoking, overall and
by region/state, gender, age, and
education, Behavioral Risk Factor
Surveillance System, 1988–1992
aggregate data 54
- Table 20.Percentage of American Indian and
Alaska Native women of reproductive
age who reported being current cigarette
smokers, overall and by region/state,
Behavioral Risk Factor Surveillance
System, 1988–1992 aggregate data 56
- Table 21.Percentage of adult Asian Americans
and Pacific Islanders who reported
being current cigarette smokers, overall
and by gender, age, and education,
National Health Interview Surveys,
United States, 1978–1995 aggregate
data 57
- Table 22.Percentage of adult Asian American and
Pacific Islander smokers who reported
smoking <15, 15–24, or \geq 25 cigarettes
per day, overall and by gender, age, and
education, National Health Interview
Surveys, United States, 1978–1995
aggregate data 58
- Table 23.Percentage of adult Asian American and
Pacific Islander ever smokers who have
quit, overall and by gender, age, and
education, National Health Interview
Surveys, United States, 1978–1995
aggregate data 59
- Table 24.Percentage of adult Asian American and
Pacific Islander women of reproductive
age who reported being current cigarette
smokers, overall and by education,
National Health Interview Surveys,
United States, 1978–1995 aggregate
data 60
- Table 25.Percentage of Chinese and Vietnamese
men who reported they smoke and the
number of cigarettes they smoke per
day, by age, education, annual house-
hold income, and level of acculturation,
Behavioral Risk Factor Surveillance
System, California, 1990 and 1991
aggregate data 62

- Table 26.Percentage of adult Asian Americans
and Pacific Islanders who reported
being current smokers, overall and by
gender, age, and education, Screener
Survey, California, 1990 and 1991
aggregate data 63
- Table 27.Summary of selected findings on the
percentage of Asian American and
Pacific Islander adults who smoke,
overall and by gender, 1975–199564
- Table 28.Percentage of adult Hispanics who
reported being current cigarette smok-
ers, overall and by gender, age, and
education, National Health Interview
Surveys, United States, 1978–1995
aggregate data 67
- Table 29. Percentage of adult Hispanic smokers who reported smoking <15, 15–24, or ≥25 cigarettes per day, overall and by gender, age, and education, National Health Interview Surveys, United States, 1978–1995 aggregate data 68
- Table 30.Percentage of adult Hispanic ever
smokers who have quit, overall and by
gender, age, and education, National
Health Interview Surveys, United States,
1978–1995 aggregate data 70
- Table 31.Percentage of Hispanic women of
reproductive age who reported being
current cigarette smokers, overall and by
education, National Health Interview
Surveys, United States, 1978–1995
aggregate data 71
- Table 32.Percentage of self-reported cigarette
brand use among adult current cigarette
smokers, overall and by race/ethnicity
and gender, National Health Interview
Surveys (NHIS) 1978–1980 combined,
Adult Use of Tobacco Survey (AUTS)
1986, and NHIS 1987
- Table 33.Percentage of self-reported cigarette
brand use among adolescent current
cigarette smokers, by race/ethnicity,
Teenage Attitudes and Practices Survey
(TAPS), 1989 and 1993

- Table 34.Relationship between smoking status
and race/ethnicity among adults, before
and after controlling for education,
National Health Interview Surveys,
United States, 1987, 1988, 1990, and 1991
aggregate data
- Table 35.Percentage of all adults and nonsmokers
who reported levels of exposure to
environmental tobacco smoke in the
home, by race/ethnicity and gender,
National Health Interview Surveys,
United States, 1991–1993 aggregate
data 87
- Table 36.Age-adjusted prevalence of current
cigarette smoking among adults, overall
and by race/ethnicity and gender,
National Health Interview Surveys,
United States, 1994 and 1995 aggregate
data 88
- Table 37.Cigarette smoking status and number of
cigarettes smoked per day among
adults, overall and by race/ethnicity and
gender, National Health Interview
Surveys, United States, 1987, 1988, 1990,
and 1991 aggregate data
- Table 38.Percentage of adults who reported using
cigars, pipes, chewing tobacco, snuff, or
any form of tobacco, overall and by
race/ethnicity and gender, National
Health Interview Surveys, United States,
1987 and 1991 aggregate data
- Table 39.Percentage of white adults who reported
being current cigarette smokers, overall
and by gender, age, and education,
National Health Interview Surveys,
United States, 1965–1995
- Table 40.Percentage of adult white smokers who
reported smoking <15, 15–24, and \geq 25
cigarettes per day, overall and by
gender, age, and education, National
Health Interview Surveys, United States,
1965–1995 100
- Table 41.Percentage of adult white ever smokers
who have quit, overall and by gender,
age, and education, National Health
Interview Surveys, United States,
1965–1995 104

- Table 42.Percentage of white women of reproduc-
tive age who reported being current
cigarette smokers, overall and by
education, National Health Interview
Surveys, United States, 1965–1995
- Table 43.Percentage of white adults who reported
being current cigarette smokers, overall
and by gender, age, and education,
National Health Interview Surveys,
United States, 1978–1995 aggregate
data 107
- Table 44. Percentage of adult white smokers who reported smoking <15, 15–24, ≥25 cigarettes per day, overall and by gender, age, and education, National Health Interview Surveys, United States, 1978–1995 aggregate data 108
- Table 45.Percentage of adult white ever smokers
who have quit, overall and by gender,
age, and education, National Health
Interview Surveys, United States,
1978–1995 aggregate data 110
- Table 46.Percentage of white women of reproduc-
tive age who reported being current
cigarette smokers, overall and by
education, National Health Interview
Surveys, United States, 1978–1995
aggregate data 111
- Table 47.Percentage of adult African Americans
who reported being current cigarette
smokers, overall and by gender, age, and
education, National Health Interview
Surveys, United States, 1965–1995
- Table 48.Percentage of adult African American
smokers who reported smoking <15,
15–24, or ≥25 cigarettes per day, overall
and by gender, age, and education,
National Health Interview Surveys,
United States, 1965–1995
- Table 49.Percentage of adult African American
ever smokers who have quit, overall and
by gender, age, and education, National
Health Interview Surveys, United States,
1965–1995
- Table 50.Percentage of African American women
of reproductive age who reported being
current cigarette smokers, overall and by
education, National Health Interview
Surveys, United States, 1965–1995120

- Table 51.Sample sizes for birth cohorts, by
gender, race/ethnicity, and education,
National Health Interview Surveys,
1978–1980, 1987 and 1988 combined,
and Hispanic Health and Nutrition
Examination Survey, 1982–1984122
- Table 52.Comparison of current smoking preva-
lence between reconstructed estimates
from National Health Interview Surveys
(NHISs), 1987 and 1988 combined, NHIS
cross-sectional survey estimates, and
Gallup poll estimates 124
- Figure 1. Trends in the prevalence of cigarette smoking among African American and white men and women, National Health Interview Surveys, United States, 1965–1995 23
- Figure 2. Trends in daily smoking among African American and white high school seniors, by gender, United States, 1977–1996 30

Figure 3. Trends in daily smoking among African American, Hispanic, and white high school seniors, United States, 1977–1996 *31*

Figure 4. Trends in smoking among African Americans and whites aged 20–34 years, United States, 1978–1995 35

Figure 5. Use of cigarettes and alcohol among African American and white high school seniors, United States, 1976–1979 and 1990–1994 38

- Figure 6. Use of cigarettes and illicit drugs among African American and white high school seniors, United States, 1976–1979 and 1990–1994 39
- Figure 7. Cigarette smoking prevalence among successive birth cohorts of African American men, by education, National Health Interview Surveys, United States, 1978–1980, 1987, and 1988 75
- Figure 8. Cigarette smoking prevalence among successive birth cohorts of African American women, by education, National Health Interview Surveys, United States, 1978–1980, 1987, and 1988 76

- Figure 9. Cigarette smoking prevalence among successive birth cohorts of Hispanic men, by education, Hispanic Health and Nutrition Examination Survey, 1982– 1984 77
- Figure 10. Cigarette smoking prevalence among successive birth cohorts of Hispanic women, by education, Hispanic Health and Nutrition Examination Survey, 1982–1984 77
- Figure 11. Reconstructed prevalence of smoking among African American adults aged 20–29 years, by gender and education, National Health Interview Surveys, United States, 1910–1988 *78*
- Figure 12. Reconstructed prevalence of smoking among Hispanic adults aged 20–29 years, by gender and education, Hispanic Health and Nutrition Examination Surveys, 1920–1984 79
- Figure 13. Trends in the age-adjusted prevalence of current cigarette smoking among African American, American Indian and Alaska Native, Asian American and Pacific Islander, Hispanic, and white adults, National Health Interview Surveys, United States, 1978–1995 aggregate data 88
- Figure 14. Trends in the age-adjusted prevalence of current cigarette smoking among African American, American Indian and Alaska Native, Asian American and Pacific Islander, Hispanic, and white men, National Health Interview Surveys, United States, 1978–1995 aggregate data 89
- Figure 15. Trends in the age-adjusted prevalence of current cigarette smoking among African American, American Indian and Alaska Native, Asian American and Pacific Islander, Hispanic, and white women, National Health Interview Surveys, United States, 1978–1995 aggregate data 89
- Figure 16. Comparison of smoking prevalence estimates from selected U.S. surveys, 1910–1991 123

Chapter 3 Health Consequences of Tobacco Use Among Four Racial/Ethnic Minority Groups

- Table 1.Death rates per 100,000 U.S. residents for
malignant diseases of the respiratory
system, by race/ethnicity and gender,
United States, 1950–1995, selected
years 139
- Table 2.Age-adjusted death rates for selected
smoking-related causes of death, by
race/ethnicity and gender, United States,
1992–1994 141
- Table 3.Death rates for lung cancer among
American Indians and Alaska Natives,
by Indian Health Service (IHS) area,
1984–19881984–1988146
- Table 4.Age-adjusted death rates for selected
smoking-related causes of death among
Mexican Americans, Puerto Rican
Americans, and Cuban Americans,
United States, 1992–1994
- Table 5.Odds ratios for the risk of lung cancer,
by gender, race/ethnicity, and smoking
status, case-control study, New Mexico,
1980–19821980–1982149
- Table 6.Age-adjusted incidence and death rates
for selected smoking-related cancers, by
race/ethnicity and gender, National
Cancer Institute's Surveillance, Epidemi-
ology, and End Results (SEER) Program,
1988–1992
- Table 7.Age-adjusted cancer death rates among
American Indians and Alaska Natives at
all 12 Indian Health Service areas,
United States, 1984–1988155
- Table 8.Odds ratios and 95% confidence inter-
vals for the risk of oral cancer associated
with cigarette smoking, by race/
ethnicity and smoking status, 1984–
19851985156
- Table 9.Odds ratios for the risk of urinary
bladder cancer associated with smoking,
by gender, race/ethnicity, and smoking
status 157
- Table 10.Rates of selected infant outcomes, by
mother's race/ethnicity, United
States 168

- Table 11.Risk of sudden infant death syndrome
associated with smoking, by race/
ethnicity, selected studies, United
States 170
- Table 12.Exposure to household smoke among
children 5 years of age and younger
and percentage distribution, by level
of exposure since birth and selected
characteristics, United States, 1988173
- Table 13.Criteria for drug dependence176
- Table 14.American Psychiatric Association
diagnostic criteria for substance
dependence 177
- Table 15.Human pharmacology of nicotine178
- Table 16.Incidence of nicotine withdrawal
symptoms, United States 179
- Table 17. Percentage of adult smokers who reported that they smoked their first cigarette within 10 minutes and within 30 minutes of awakening, by race/ ethnicity and number of cigarettes smoked per day, National Health Interview Survey, United States, 1987 182
- Table 18.Percentage of men and women who
considered smoking a habit or addiction,
overall and by smoking status, Current
Population Survey, United States, 1992–
1993 184
- Figure 1. Incidence of cancer of the lung and bronchus, by race/ethnicity and gender, National Cancer Institute's Surveillance, Epidemiology, and End Results (SEER) Program, 1973–1994 138
- Figure 2. Age-adjusted lung cancer death rates among American Indian and Alaska Native men in selected states compared with rates among all U.S. men, 1968–1987 144
- Figure 3. Age-adjusted lung cancer death rates among American Indian and Alaska Native women in selected states compared with rates among all U.S. women, 1968–1987 145

- Figure 4. SEER cancer incidence and U.S. death rates, 1988–1992, ratio of African American rate to white rate for all ages, by cancer site 154
- Figure 5. Serum cotinine levels by number of cigarettes smoked daily for African Americans, Mexican Americans, and whites, National Health and Nutrition Examination Survey, United States, 1988–1991 *180*

Chapter 4

Factors That Influence Tobacco Use Among Four Racial/Ethnic Minority Groups

Table 1.High school seniors' perceptions
about the risks associated with cigarette
smoking, Monitoring the Future sur-
veys, United States, 1980–1989230

Chapter 5

Tobacco Control and Education Efforts Among Members of Four Racial/Ethnic Minority Groups

- Table 1.Percentage of adult smokers who would
like to stop smoking, by race/ethnicity
and gender, National Health Interview
Survey, United States, 1993 261
- Table 2.Adults' beliefs about the health effects of
smoking, by race/ethnicity, gender, and
smoking status, National Health Inter-
view Survey, United States, 1992264
- Table 3.Adults' beliefs about minors' ease in
purchasing cigarettes and other tobacco
products, by race/ethnicity, smoking
status, and gender, Current Population
Survey, United States, 1992–1993
- Table 4.Public support for and beliefs about
policies regarding tobacco access and
marketing, by selected characteristics,
Robert Wood Johnson Foundation Youth
Access Survey, 1994 270
- Table 5.Percentage of adult smokers who have
received advice to quit smoking from
either a medical doctor or a dentist, by
race/ethnicity and gender, Current
Population Survey, United States, 1992–
1993 285

- Table 6.Percentage of adults who reported that
no one is allowed to smoke anywhere
inside the home, by race/ethnicity,
smoking status, and gender, Current
Population Survey, United States, 1992–
1993 288
- Table 7.Percentage of adults who think that
smoking should be allowed in some
areas or not allowed at all in selected
public locations, by race/ethnicity and
smoking status, Current Population
Survey, United States, 1992–1993
290
- Table 8.Percentage of adults who think that the
advertising of tobacco products should
be always allowed or not allowed at all,
by race/ethnicity, smoking status, and
gender, Current Population Survey,
United States, 1992–1993 294

- Table 9.Percentage of Californians who support
curtailment of tobacco advertising and
promotion efforts, by race/ethnicity,
1992 295
- Table 10.Percentage of adults who think that
giving away free tobacco samples
should be always allowed or not al-
lowed at all, by race/ethnicity, smoking
status, and gender, Current Population
Survey, United States, 1992–1993
296
- Table 11.Public beliefs about and support for
policies related to nicotine and tobacco
product regulation, Robert Wood
Johnson Foundation Youth Access
Survey, 1994
- Figure 1. Billboard used by the California Department of Health Services in targeting Hispanics to quit smoking 261