

**TABLE. Tobacco use screening, and cessation assistance during outpatient visits to office-based physicians among adults aged  $\geq 18$  years, by selected demographic characteristics — National Ambulatory Medical Care Survey, United States, 2010**

Characteristic	Visits with tobacco screening* (n = 17,015)		Visits with current tobacco use <sup>†</sup> (n = 3,056)		Visits with current tobacco use with tobacco counseling <sup>§</sup> (n = 570)		Visits with current tobacco use with tobacco counseling and/or cessation medication <sup>¶</sup> (n = 627)	
	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
<b>Total (n = 25,757)</b>	65.6	(62.1, 69.1)	16.9	(15.5, 18.2)	19.1	(16.3, 22.0)	20.7	(17.9, 23.5)
<b>Age group (years)</b>								
18–24	69.8	(64.8, 74.7)	18.9	(15.7, 22.2)	18.6	(11.0, 26.2)	18.8	(11.1, 26.5)
25–44	68.5	(64.7, 72.2)	20.4	(18.2, 22.6)	17.7	(13.5, 21.9)	19.5	(15.1, 23.9)
45–64	64.7	(60.8, 68.5)	20.1	(18.1, 22.1)	21.1	(17.2, 25.1)	23.2	(19.4, 27.0)
$\geq 65$	63.5	(59.1, 67.8)	9.5	(8.2, 10.9)	17.1	(11.2, 23.1)	17.7	(11.8, 23.7)
<b>Sex</b>								
Male	64.4	(60.7, 68.1)	20.7	(18.9, 22.6)	19.7	(16.5, 22.8)	21.1	(17.9, 24.2)
Female	66.4	(62.8, 70.0)	14.4	(13.0, 15.8)	18.7	(14.6, 22.7)	20.4	(16.4, 24.4)
<b>Race/ethnicity</b>								
White, non-Hispanic	67.0	(63.5, 70.5)	17.5	(16.0, 19.0)	18.2	(15.1, 21.2)	20.0	(17.0, 23.0)
Black, non-Hispanic	62.5	(56.0, 69.0)	18.8	(16.0, 21.6)	25.6	(15.9, 35.2)	25.7	(16.1, 35.4)
Hispanic	60.1	(51.9, 68.3)	10.6	(8.5, 12.7)	20.2	(11.8, 28.5)	20.9	(12.7, 29.0)
Other Race/Multiple Race, Non-Hispanic	60.2	(46.4, 74.0)	11.6	(7.9, 15.3)	**	**	**	**
<b>††Poverty%</b>								
Less than 5%	65.2	(60.8, 69.6)	13.0	(11.6, 14.5)	19.4	(12.5, 26.3)	22.0	(14.7, 29.2)
5.00% - 9.99%	66.8	(62.3, 71.2)	16.0	(13.9, 18.1)	20.9	(16.4, 25.4)	22.6	(18.1, 27.2)
10.00% - 19.99%	67.6	(63.4, 71.7)	19.6	(17.5, 21.7)	19.2	(15.0, 23.5)	20.1	(15.8, 24.4)
20% or more	60.6	(54.0, 67.2)	19.3	(16.0, 22.5)	16.2	(12.0, 20.5)	17.9	(13.7, 22.1)

**Abbreviations:** CI = confidence interval

\* Visits during which current tobacco use (currently smoke cigarettes or cigars or use snuff or chewing tobacco) or no current use was recorded. Denominator includes current tobacco use, no current use, unknown, and blanks.

<sup>†</sup> Visits during which current tobacco use (smoking cigarettes or cigars or using snuff or chewing tobacco) was recorded.

<sup>§</sup> Tobacco counseling refers to any information provided that related to tobacco use in any form, including cigarettes, cigars, snuff, and chewing tobacco, and on exposure to tobacco in the form of secondhand smoke, smoking cessation, and prevention of tobacco use, as well as referrals to other health-care providers for smoking cessation programs.

<sup>¶</sup> Cessation medications include nicotine replacement therapy (nicotine patch, gum, lozenge, nasal spray, and inhaler), bupropion, and varenicline.

\*\* Data not shown because sample size is  $< 30$ , or the relative standard error of the estimate is  $> 30\%$ .

<sup>††</sup> Percent of poverty in patient's zip code.