

- Warren CW, Jones NR, Chauvin J, Peruga A. Tobacco use and cessation counseling: cross-country data from the Global Health Professions Student Survey (GHPSS), 2005-2007. *Tobacco Control*. 2008; 17: 238-247.
- Sirichotiratana N, Sovann S, Aditama TY, Krishnan M, Kyaing NN, Miguel-Baquilod M, Hai PT, Sinha DN, Warren CW, Jones NR. Prevalence of tobacco use among students aged 13-15 years in Association of Southeast Asian Nations (ASEAN) member states, 2000-06. *Tobacco Control Online*. 31 July 2008.
- Sinha DN, Gupta PC, Reddy KS, Prasad VM, Rahman K, Warren CW, Jones NR, Asma S. Linking Global Youth Tobacco Survey (GYTS) 2003 and 2006 Data to Tobacco Control Policy in India. *Journal of School Health*. 2008; 78(7): 368-373.
- Rachiotis G, Muula AS, Rudatsiikira E, Siziya S, Kyrlesis A, Gourgoulis K, Hadjichristodoulou C. Factors associated with adolescent cigarette smoking in Greece: Results from a cross sectional study (GYTS Study). *BMC Public Health (Online)*. 2008; 8: 313.
- Sirichotiratana N, Techatrakakdi C, Rahman K, Warren CW, Jones NR, Asma S, Lee J. Linking Global Youth Tobacco Survey (GYTS) Data to the WHO Framework Convention on Tobacco Control: the case for Thailand. (*BMC Public Health Online Supplement*) (In press)
- Zavaleta A, Salas M, Peruga A, Curi A, Warren CW, Jones NR, Asma S. Linking Global Youth Tobacco Survey (GYTS) Data to the WHO Framework Convention on Tobacco Control: the case for Peru. (*BMC Public Health Online Supplement*) (In press)
- Erguder T, Çakır B, Aslan D, Warren CW, Jones NR, Asma S. Linking Global Youth Tobacco Survey (GYTS) Data to the tobacco control policies in Turkey. (*BMC Public Health Online Supplement*) (In press)
- El-Awa F, Warren CW, Jones NR. Changes in tobacco use among 13-15 year olds between 1999 and 2005: findings from the Eastern Mediterranean Region. (*Eastern Mediterranean Health Journal*) (In press)
- Sovinova H, Csemy L, Prochazka B, Kottnerova S. Smoking attributable mortality in the Czech Republic: a comparison of two estimates. *Journal of Public Health*. 2008; 16: 37-42.
- Baska T, Pudule I, Tilgale N, Warren CW, Lee J, Lea V, Jones NR. Smoking tobacco in waterpipes among adolescents in Europe: the case of Latvia and Slovakia. (*Tobacco Control*) (In press)
- Christophi CA, Kolokotroni-Zacharopoulou O, Alpert HR, Warren CW, Jones NR, Demokritou P, Connolly GN. Prevalence and main determinants of tobacco use in Cyprus youth through the use of the GYTS. *BMC Public Health (In press)*
- Sinha DN, Gewaifel G, Pandey MR, Khan SA, Gunasekara PW, El-Awa F, Warren CW, Jones NR. Prevalence of tobacco use among students aged 13-15 years in South Asian Association for Regional Cooperation (SAARC) Member Countries, 2003-2006. (*Indian Journal of Public Health*) (Under review)
- Sovinova H, Csemy L, Warren CW, Lee J, Lea V. Changes in tobacco use among 13-15 year olds in Czech Republic: 2002-07. (*Central European Journal of Public Health*) (Under review)
- Saade G, Jaoude SA, Soweid RA, Warren CW, Jones NR. Alarming changes in patterns of tobacco use: results from 2005 GYTS in Lebanon. (*Eastern Mediterranean Health Journal*) (Under review)
- Chen PL, Huang W, Chuang Y, Jones NR, Warren CW. Secondhand smoke exposure and knowledge and attitudes concerning tobacco use among Junior High and Senior High School students in Taiwan. (*Asian Pacific Journal of Public Health*) (Under review)
- Baska T, Warren CW, Baskova M, Jones NR, Lea V, Lee J. Prevalence of youth cigarette smoking and factors influencing use among 25 European countries – findings from the Global Youth Tobacco Survey (GYTS). (*International Journal of Public Health*) (Under review)
- Roa R, Franklin-Peroune R, Warren CW, Lee J, Lea V, Goding, A, Jones NR, Asma S. Changes in tobacco use among 13-15 year olds in Panama, 2002 and 2008. (Submitted *MMWR*).
- Maki Barto S, El Awa F, Fouad H, Warren CW, Lee J, Lea V, Goding A, Asma S, McKenna M. Tobacco use among students aged 13-15 years – Baghdad, 2008. (Submitted *MMWR*).
- Khader A, Shaheen Y, Turki Y, El Awa F, Fouad H, Warren CW, Jones NR, Lea V. Tobacco use among UNRWA students aged 13-15. (In progress)
- Warren CW, Jones NR, Lea V, Lee J. Change in tobacco use among 13-15 year olds between 1999 and 2008: Findings from the Global Youth Tobacco Survey. (In progress)
- San Martin V, Franklin-Peroune R, Warren CW, Lee J, Lea V. Linking Global Youth Tobacco Survey (GYTS) 2003 and 2008 data to tobacco control policy in Asuncion, Paraguay. (In progress)
- ban on tobacco  
 advertising, promotion and sponsorship 20, 23, 48  
 reported by health professions students 71  
 smoking 29, 43  
 by school personnel 79  
 by students 78  
 in training-school buildings 71  
*see also* smoke-free laws
- bidis *see* other tobacco use
- Bloomberg Initiative to Reduce Tobacco Use 76, 85
- boys  
 other tobacco use 38-39  
 smoking 34-35  
*see also* youth
- Canadian Public Health Association (CPHA) 21, 65
- Centers for Disease Control and Prevention (CDC) 21, 32, 65  
*Youth Tobacco Cessation: A Guide for Making Informed Decisions* 45
- cessation services 20, 23, 44-45, 60-61, 64, 72-73  
 materials 61  
 training 60
- cigarette smoking  
 age comparison 36  
 by boys 34-35, 86-95  
 by girls 36-37, 86-95  
 by school personnel 56-57  
 comparison with other tobacco use 39, 40  
 gender comparison 34, 35, 37, 38  
 price sensitivity 20, 50-51
- data on tobacco use 22, 32, 33, 86-95
- dentistry students 64  
 exposure to second-hand smoke 70  
 reporting school ban 71  
 tobacco use 68  
 tobacco use counseling 72-73
- environmental tobacco smoke (ETS) *see* second-hand smoke
- European Community 20
- Framework Convention on Tobacco Control (FCTC) 20-21, 22, 50, 85  
 article from 43, 44, 46, 49, 51, 59, 60, 70, 72  
 country status 20-21, 86-95
- girls  
 other tobacco use 40-41  
 smoking 36-37  
*comparison with adults* 36  
*see also* youth
- Global Adult Tobacco Survey (GATS) 29, 74-77, 85  
 countries surveyed 76-77, 86-95  
 survey method 76-77
- Global Health Professions Student Survey (GHPSS) 28, 62-73, 85  
 countries surveyed 64-65, 66-67, 86-95  
 survey method 66-67
- Global School Personnel Survey (GSPS) 27, 52-61, 84  
 countries surveyed 54-55, 86-95  
 survey method 54-55
- Global Tobacco Surveillance System (GTSS) 21  
 data dissemination 80-81  
 survey method 26, 28, 32-33  
 workshops 26-27, 28-29
- Global Youth Tobacco Survey (GYTS) 26, 30-51, 84  
 countries surveyed 32-33, 86-95  
 survey method 32-33  
 topics covered 32
- government policy on tobacco control 20-21, 44-45, 46-47, 48
- health professions students 64-73  
 exposure to second-hand smoke 70  
 reporting school ban 71  
 tobacco use 68  
 tobacco use counseling 72-73
- illicit trade in tobacco products 20
- labeling of tobacco products 20, 23  
 warnings 46-47
- medicine students 65  
 exposure to second-hand smoke 70  
 reporting school ban 71  
 tobacco use 68  
 tobacco use counseling 72-73
- MPOWER 22-23, 47, 60, 71, 85  
 definition of public places 43  
 strategies 20, 22, 23, 50

- nursing students 66
  - exposure to second-hand smoke 70
  - reporting school ban 71
  - tobacco use 69
  - tobacco use counseling 72–73
- other tobacco use 38–41
  - by boys 38–39, 86–95
  - by girls 40–41, 86–95
  - by school personnel 57
  - comparison with cigarette smoking 39, 40
- packet warnings 46–47
- pharmacy students 67
  - exposure to second-hand smoke 70
  - reporting school ban 71
  - tobacco use 69
  - tobacco use counseling 72–73
- pipes *see* other tobacco use
- population data 86–95
- prohibition of sales to minors 20
- public awareness of dangers of tobacco use 20, 46
- quitting *see* cessation services
- schools
  - policies on smoking 55, 58–59
  - tobacco use by personnel 56–57
  - smoking in 43, 58–59
- school personnel
  - ban on smoking in school 59
  - cigarette smoking by 56–57, 58–59
  - gender comparison 56–57
  - other tobacco use by 56–57
  - training on tobacco use prevention 60–61
- school students
  - ban on smoking 58
- second-hand smoke (SHS)
  - 42–43, 47, 70–71
  - health effects of 42, 70
  - in home 42, 43
  - in public places 42, 70–71
  - in work place 43
- shisha *see* other tobacco use
- smoke-free
  - areas 71
  - laws 23, 43
- smokeless tobacco *see* other tobacco use
- smoking
  - at work 20, 23, 43
  - health effects of 46, 60, 64
  - in public places 20, 23, 42
  - in schools 58–59
  - in training schools 70–71
- susceptibility to initiate smoking
  - boys 35
  - girls 36
- taxes, tobacco 20, 23, 50–51
  - use to fund cessation services 44
- teachers *see* school personnel
- tobacco industries 71
  - promotion by 48–49, 60
- tobacco-control
  - expenditure 23
  - milestones 84–85
- water pipes *see* other tobacco use
- World Health Organization (WHO) 20–21, 22–23, 32, 43, 44, 46, 49, 51, 84–85
- WHO *Report on the Global Tobacco Epidemic, 2008* 20, 22, 44, 85 *see also* MPOWER
- World Bank, tobacco taxes 50
- youth
  - cessation counseling 72–73
  - desire to quit 44–45
  - tobacco purchasing patterns 50
  - tobacco use 30–51
  - views on smoking ban 43