

References:

1. C Velicer. ACIP Presentation. Post-licensure Safety Study of Quadrivalent Human Papillomavirus Vaccine among 189,629 Females Presented at October 2011 ACIP Meeting.
2. Ahmed F, Temte JL, Campos-Outcalt D, Schünemann HJ, for the ACIP Evidence Based Recommendations Work Group (EBRWG). Methods for developing evidence-based recommendations by the Advisory Committee on Immunization Practices (ACIP) of the U.S. Centers for Disease Control and Prevention (CDC). *Vaccine* 2011;29:9171-76.
3. Bohlke K, Davis RL, DeStefano F, Marcy SM, Braun MM, Thompson RS; Vaccine Safety Datalink Team. Epidemiology of anaphylaxis among children and adolescents enrolled in a health maintenance organization. *J Allergy Clin Immunol* 2004;113:536-42.
4. Chesson HW, Ekwueme DU, Saraiya M, Dunne EF, Markowitz LE. The cost-effectiveness of male HPV vaccination in the United States. *Vaccine* 2011;29:8443-50.
5. Food and Drug Administration, Product approval information – licensing action, package insert: Gardasil (quadrivalent human papillomavirus types 6, 11, 16 and 18 HPV), Merck & Co. Whitehouse Station, NJ: Food and Drug Administration; 2009.
6. Gee J, Naleway A, Shui I, et al. Monitoring the safety of quadrivalent human papillomavirus vaccine: findings from the Vaccine Safety Datalink. *Vaccine* 2011;29:8279-84.
7. Giuliano AR, Palefsky JM, Goldstone S, et al. Efficacy of quadrivalent HPV vaccine against HPV infection and disease in males. *N Engl J Med* 2011;364:401-11.
8. Palefsky JM, Giuliano AR, Goldstone S, et al. HPV vaccine against anal HPV infection and anal intraepithelial neoplasia. *N Engl J Med* 2011;365:1576-85.