

Janssen COVID-19 Vaccine (Johnson & Johnson)

Ages: 18 years of age and older

Use for: Single dose from a multidose vial. COVID-19 vaccines are NOT interchangeable.

Route: Intramuscular (IM) injection

Beyond-Use Time: Use within 6 hours after the vial is first punctured if held between 2°C and 8°C (36°F and 46°F) or use within 2 hours after the vial is first punctured if held at room temperature (maximally 25°C or 77°F).



Janssen COVID-19 Vaccine (Johnson & Johnson)

Ages: 18 years of age and older

Use for: Single dose from a multidose vial. COVID-19 vaccines are NOT interchangeable.

Route: Intramuscular (IM) injection

Beyond-Use Time: Use within 6 hours after the vial is first punctured if held between 2°C and 8°C (36°F and 46°F) or use within 2 hours after the vial is first punctured if held at room temperature (maximally 25°C or 77°F).



Janssen COVID-19 Vaccine (Johnson & Johnson)

Ages: 18 years of age and older

Use for: Single dose from a multidose vial. COVID-19 vaccines are NOT interchangeable.

Route: Intramuscular (IM) injection

Beyond-Use Time: Use within 6 hours after the vial is first punctured if held between 2°C and 8°C (36°F and 46°F) or use within 2 hours after the vial is first punctured if held at room temperature (maximally 25°C or 77°F).



Janssen COVID-19 Vaccine (Johnson & Johnson)

Ages: 18 years of age and older

Use for: Single dose from a multidose vial. COVID-19 vaccines are NOT interchangeable.

Route: Intramuscular (IM) injection

Beyond-Use Time: Use within 6 hours after the vial is first punctured if held between 2°C and 8°C (36°F and 46°F) or use within 2 hours after the vial is first punctured if held at room temperature (maximally 25°C or 77°F).



Janssen COVID-19 Vaccine (Johnson & Johnson)

Ages: 18 years of age and older

Use for: Single dose from a multidose vial. COVID-19 vaccines are NOT interchangeable.

Route: Intramuscular (IM) injection

Beyond-Use Time: Use within 6 hours after the vial is first punctured if held between 2°C and 8°C (36°F and 46°F) or use within 2 hours after the vial is first punctured if held at room temperature (maximally 25°C or 77°F).



Janssen COVID-19 Vaccine (Johnson & Johnson)

Ages: 18 years of age and older

Use for: Single dose from a multidose vial. COVID-19 vaccines are NOT interchangeable.

Route: Intramuscular (IM) injection

Beyond-Use Time: Use within 6 hours after the vial is first punctured if held between 2°C and 8°C (36°F and 46°F) or use within 2 hours after the vial is first punctured if held at room temperature (maximally 25°C or 77°F).

