

## Janssen COVID-19 Vaccine (Johnson & Johnson)

**Ages:** 18 years of age and older

**Use for:** Single dose from a multidose vial.

**Route:** Intramuscular (IM) injection

**Beyond-Use Time:** Use within 6 hours after the vial is first punctured if held between 2°C and 8°C (36°F and 46°F) or use within 2 hours after the vial is first punctured if held at room temperature (maximally 25°C or 77°F).



## Janssen COVID-19 Vaccine (Johnson & Johnson)

**Ages:** 18 years of age and older

**Use for:** Single dose from a multidose vial.

**Route:** Intramuscular (IM) injection

**Beyond-Use Time:** Use within 6 hours after the vial is first punctured if held between 2°C and 8°C (36°F and 46°F) or use within 2 hours after the vial is first punctured if held at room temperature (maximally 25°C or 77°F).



## Janssen COVID-19 Vaccine (Johnson & Johnson)

**Ages:** 18 years of age and older

**Use for:** Single dose from a multidose vial.

**Route:** Intramuscular (IM) injection

**Beyond-Use Time:** Use within 6 hours after the vial is first punctured if held between 2°C and 8°C (36°F and 46°F) or use within 2 hours after the vial is first punctured if held at room temperature (maximally 25°C or 77°F).



## Janssen COVID-19 Vaccine (Johnson & Johnson)

**Ages:** 18 years of age and older

**Use for:** Single dose from a multidose vial.

**Route:** Intramuscular (IM) injection

**Beyond-Use Time:** Use within 6 hours after the vial is first punctured if held between 2°C and 8°C (36°F and 46°F) or use within 2 hours after the vial is first punctured if held at room temperature (maximally 25°C or 77°F).



## Janssen COVID-19 Vaccine (Johnson & Johnson)

**Ages:** 18 years of age and older

**Use for:** Single dose from a multidose vial.

**Route:** Intramuscular (IM) injection

**Beyond-Use Time:** Use within 6 hours after the vial is first punctured if held between 2°C and 8°C (36°F and 46°F) or use within 2 hours after the vial is first punctured if held at room temperature (maximally 25°C or 77°F).



## Janssen COVID-19 Vaccine (Johnson & Johnson)

**Ages:** 18 years of age and older

**Use for:** Single dose from a multidose vial.

**Route:** Intramuscular (IM) injection

**Beyond-Use Time:** Use within 6 hours after the vial is first punctured if held between 2°C and 8°C (36°F and 46°F) or use within 2 hours after the vial is first punctured if held at room temperature (maximally 25°C or 77°F).

