

Healthcare Provider Office Information

We believe that getting vaccinated is a critical step in protecting your health. Vaccines can help prevent common diseases that can be serious and costly for you or your loved ones.

Each year, thousands of adults in America suffer serious health problems (and some even die) from diseases they could be vaccinated against like whooping cough, hepatitis A and B, flu, pneumococcal disease, and shingles. Older adults and those with chronic health conditions are at increased risk for complication from certain diseases.

Together, let's take an active role in helping you and your loved ones stay healthy. To learn more about vaccines for adults, visit www.cdc.gov/vaccines/adults or use the Adult Vaccine Self-Assessment Tool at <https://www2.cdc.gov/nip/adultimmsched/> to find out which vaccines you may need.

All adults need a seasonal flu (influenza) vaccine every year. Look at the table below to see other vaccines you may need.

Please take a moment to fill out the questionnaire below to help us determine which vaccines may be recommended for you based on your specific health status, age, and lifestyle. Keep in mind that this list may not include every vaccine you need.

Check all that apply to you	Let's discuss these recommended vaccines
<input type="checkbox"/> I am 19 years or older	<ul style="list-style-type: none"> • Tetanus, diphtheria, and whooping cough (Tdap) vaccine for all adults who have never received Tdap vaccine • Tetanus and diphtheria (Td) vaccine or Tdap vaccine every 10 years <div style="border: 1px solid black; padding: 2px; text-align: center; margin-top: 5px;"> PREGNANT WOMEN SHOULD GET A TDAP VACCINE DURING EACH PREGNANCY </div>
<input type="checkbox"/> I am 50 years or older	<ul style="list-style-type: none"> • Shingles vaccine
<input type="checkbox"/> I am 65 years or older	<ul style="list-style-type: none"> • Pneumococcal polysaccharide vaccine • May discuss and decide, with clinician, to receive pneumococcal conjugate vaccine
<input type="checkbox"/> I didn't receive the human papillomavirus (HPV) vaccine series as a child	<ul style="list-style-type: none"> • HPV vaccine series (3 dose series) <ul style="list-style-type: none"> ◦ Females and males age 26 or younger
<input type="checkbox"/> I was born in the U.S. in 1957 or after or don't have immunity against measles, mumps, and rubella	<ul style="list-style-type: none"> • Measles, mumps, rubella (MMR) vaccine* (one dose)
<input type="checkbox"/> I was born in the U.S. in 1980 or after and don't have immunity against chickenpox	<ul style="list-style-type: none"> • Varicella "chickenpox" vaccine*
<input type="checkbox"/> I am a healthcare worker	<ul style="list-style-type: none"> • Hepatitis B vaccine series • Measles, mumps, rubella (MMR) vaccine* • Varicella "chickenpox" vaccine*
<input type="checkbox"/> I have heart disease, asthma or chronic lung disease	<ul style="list-style-type: none"> • Pneumococcal polysaccharide vaccine



U.S. Department of Health and Human Services
Centers for Disease Control and Prevention

Flip page to continue questionnaire

Healthcare Provider Office Information

<input type="checkbox"/> I have type 1 or type 2 diabetes	<ul style="list-style-type: none"> • Hepatitis B vaccine series • Pneumococcal polysaccharide vaccine
<input type="checkbox"/> I have a weakened immune system	<ul style="list-style-type: none"> • <i>Haemophilus influenzae</i> type b (Hib) vaccine (post-hematopoietic stem cell transplant only) • HPV vaccine series (if 22-26 years of age and not previously vaccinated) • Both types of meningococcal vaccines (meningococcal conjugate and serogroup B) (If you have complement component deficiency or taking complement inhibitor such as Soliris® or Ultomiris®) • Both types of pneumococcal vaccines (one dose of conjugate first, then one dose of polysaccharide >8 weeks later)
<input type="checkbox"/> I have HIV	<ul style="list-style-type: none"> • Hepatitis B vaccine series • HPV vaccine series (if 22-26 year of age and not previously vaccinated) • Meningococcal conjugate vaccine • Both types of pneumococcal vaccines (one dose of conjugate first, then one dose of polysaccharide >8 weeks later)
<input type="checkbox"/> I have chronic liver disease	<ul style="list-style-type: none"> • Hepatitis A vaccine series • Hepatitis B vaccine series • Pneumococcal polysaccharide vaccine
<input type="checkbox"/> I do not have a spleen or my spleen does not work well	<ul style="list-style-type: none"> • <i>Haemophilus influenzae</i> type b (Hib) vaccine (post-hematopoietic stem cell transplant only) • Both types of meningococcal vaccines (meningococcal conjugate and serogroup B) • Both types of pneumococcal vaccines (conjugate plus polysaccharide, get the conjugate vaccine first)
<input type="checkbox"/> I am a man who has sex with men	<ul style="list-style-type: none"> • Hepatitis A vaccine series • Hepatitis B vaccine series • HPV vaccine series (if 22-26 of age and not previously vaccinated)
<input type="checkbox"/> I am a laboratory worker and may be routinely exposed to isolates of <i>Neisseria meningitidis</i> , or specimens potentially containing hepatitis A or hepatitis B virus	<ul style="list-style-type: none"> • Hepatitis A vaccine series • Hepatitis B vaccine series • Both types of meningococcal vaccines (meningococcal conjugate and serogroup B)
<input type="checkbox"/> I am a college freshman living in a residence hall	<ul style="list-style-type: none"> • Measles, mumps, rubella (MMR) vaccine* • Meningococcal conjugate vaccine
<input type="checkbox"/> I am planning to travel out of the U.S.	<ul style="list-style-type: none"> • Talk to your healthcare professional to learn which vaccines you may need based on locations of travel.

*This is a live vaccine and should not be given to people who have a very weakened immune system, including those with a CD4 count less than 200, or to pregnant women.



**U.S. Department of
Health and Human Services**
Centers for Disease
Control and Prevention