


Immunizations and Developmental Milestones for Your Child from Birth Through 6 Years Old

Child's Name _____

Birth Date _____

		Birth	1 MONTH	2 MONTHS	4 MONTHS	6 MONTHS
Recommended Immunizations	Hepatitis B	<input type="radio"/> HepB	<input type="radio"/> HepB ¹			<input type="radio"/> HepB
	Rotavirus			<input type="radio"/> RV	<input type="radio"/> RV	<input type="radio"/> RV ²
	Diphtheria, Tetanus, Pertussis			<input type="radio"/> DTaP	<input type="radio"/> DTaP	<input type="radio"/> DTaP
	<i>Haemophilus influenzae</i> type b			<input type="radio"/> Hib	<input type="radio"/> Hib	<input type="radio"/> Hib
	Pneumococcal			<input type="radio"/> PCV	<input type="radio"/> PCV	<input type="radio"/> PCV
	Inactivated Poliovirus			<input type="radio"/> IPV	<input type="radio"/> IPV	<input type="radio"/> IPV
	Influenza (Flu)					<input type="radio"/> Influenza, first dose ³ <input type="radio"/> second dose
Milestones*	<p>Milestones should be achieved by the age indicated.</p> <p>Talk to your child's doctor about age-appropriate milestones if your child was born prematurely.</p> <ul style="list-style-type: none"> <input type="radio"/> Recognizes caregiver's voice <input type="radio"/> Turns head toward breast or bottle <input type="radio"/> Communicates through body language, fussing or crying, alert and engaged <input type="radio"/> Startles to loud sounds 	<ul style="list-style-type: none"> <input type="radio"/> Starts to smile <input type="radio"/> Raises head when on tummy <input type="radio"/> Calms down when rocked, cradled or sung to <input type="radio"/> Pays attention to faces 	<ul style="list-style-type: none"> <input type="radio"/> Begins to smile at people <input type="radio"/> Coos, makes gurgling sounds <input type="radio"/> Begins to follow things with eyes <input type="radio"/> Can hold head up 	<ul style="list-style-type: none"> <input type="radio"/> Babbles with expression <input type="radio"/> Likes to play with people <input type="radio"/> Reaches for toy with one hand <input type="radio"/> Brings hands to mouth <input type="radio"/> Responds to affection <input type="radio"/> Holds head steady, unsupported 	<ul style="list-style-type: none"> <input type="radio"/> Knows familiar faces <input type="radio"/> Responds to own name <input type="radio"/> Brings things to mouth <input type="radio"/> Rolls over in both directions <input type="radio"/> Strings vowels together when babbling ("ah", "eh", "oh") 	
Growth		_____	_____	_____	_____	_____
		WEIGHT / PERCENTILE	WEIGHT / PERCENTILE	WEIGHT / PERCENTILE	WEIGHT / PERCENTILE	WEIGHT / PERCENTILE
		_____	_____	_____	_____	_____
	LENGTH / PERCENTILE	LENGTH / PERCENTILE	LENGTH / PERCENTILE	LENGTH / PERCENTILE	LENGTH / PERCENTILE	
	_____	_____	_____	_____	_____	
	HEAD CIRCUMFERENCE	HEAD CIRCUMFERENCE	HEAD CIRCUMFERENCE	HEAD CIRCUMFERENCE	HEAD CIRCUMFERENCE	

continues on back page

 Shaded boxes indicate the vaccine can be given during shown age range.

VISIT DATE

VISIT DATE

VISIT DATE

VISIT DATE

VISIT DATE

¹ The second dose of HepB may be given either at the 1 month or 2 month visit.

² A third dose of rotavirus vaccine is only needed for RotaTeq.

³ Two doses given at least four weeks apart are recommended for children aged 6 months through 8 years of age who are getting a flu vaccine for the first time and for some other children in this age group.

* Milestones adapted from *Caring for your baby and young child: Birth to age 5*, Fifth Edition, edited by Steven Shelov and Tanya Remer Altmann © 1991, 1993, 1998, 2004, 2009 by the American Academy of Pediatrics and *Bright Futures: Guidelines for health supervision of infants, children, and adolescents*, Third Edition, edited by Joseph Hagan, Jr., Judith S. Shaw, and Paula M. Duncan, 2008, Elk Grove Village, IL: American Academy of Pediatrics.

This is not an exhaustive list of milestones from 0-6 years. See more at www.cdc.gov/milestones

If your child has any medical conditions that put him at risk for infections or is traveling outside the United States, talk to your child's doctor about additional vaccines that he may need.



U.S. Department of Health and Human Services
Centers for Disease Control and Prevention



American Academy of Pediatrics



DEDICATED TO THE HEALTH OF ALL CHILDREN™

Immunizations and Developmental Milestones for Your Child from Birth Through 6 Years Old

Child's Name _____

Birth Date _____

		12 MONTHS	15 MONTHS	18 MONTHS	19 23 MONTHS	2-3 YEARS	4-6 YEARS
Recommended Immunizations	Hepatitis B	<input type="radio"/> HepB (Final dose administered between 6 and 18 months)					
	Diphtheria, Tetanus, Pertussis		<input type="radio"/> DTaP				<input type="radio"/> DTaP
	Haemophilus influenzae type b	<input type="radio"/> Hib					
	Pneumococcal	<input type="radio"/> PCV					
	Inactivated Poliovirus	<input type="radio"/> IPV (Third dose administered between 6 and 18 months)					<input type="radio"/> IPV
	Influenza (Flu)	<input type="radio"/> Influenza , first dose ² <input type="radio"/> second dose (if needed)				Age 2 3 <input type="radio"/> <input type="radio"/> Influenza , first dose ² <input type="radio"/> <input type="radio"/> second dose (if needed)	Age 4 5 6 <input type="radio"/> <input type="radio"/> <input type="radio"/> Influenza , first dose ² <input type="radio"/> <input type="radio"/> <input type="radio"/> second dose (if needed)
	Measles, Mumps, Rubella	<input type="radio"/> MMR					<input type="radio"/> MMR
	Varicella	<input type="radio"/> Varicella					<input type="radio"/> Varicella
Hepatitis A	<input type="radio"/> <input type="radio"/> Hep A³						
Milestones*	Milestones should be achieved by the age indicated. Talk to your child's doctor about age-appropriate milestones if your child was born prematurely.	<input type="radio"/> Cries when mom or dad leaves <input type="radio"/> Says "mama" and "dada" <input type="radio"/> Copies gestures (for example, waves "bye bye") <input type="radio"/> May stand alone <input type="radio"/> Looks at right picture or thing when named	<input type="radio"/> Imitates what you are doing <input type="radio"/> Drinks from a cup <input type="radio"/> Scribbles on his own <input type="radio"/> Walks well <input type="radio"/> Says a couple of words other than "mama" and "dada"	<input type="radio"/> Points to show others something interesting <input type="radio"/> Says several single words <input type="radio"/> Points to one body part <input type="radio"/> May walk up steps and run	<input type="radio"/> Plays mainly beside other children <input type="radio"/> Follows two-step commands <input type="radio"/> Plays simple make-believe games <input type="radio"/> Throws ball overhand	<input type="radio"/> Can name most familiar things <input type="radio"/> Shows affection for friends without prompting <input type="radio"/> Turns book pages one at a time <input type="radio"/> Kicks a ball	<input type="radio"/> Speaks very clearly <input type="radio"/> Tells stories <input type="radio"/> Can print some letters or numbers <input type="radio"/> Hops; may be able to skip <input type="radio"/> Enjoys playing with other children
	Growth	At each well child visit, enter date, length, weight, and percentile information to keep track of your child's progress.	WEIGHT / PERCENTILE	WEIGHT / PERCENTILE	WEIGHT / PERCENTILE	WEIGHT / PERCENTILE	WEIGHT
		LENGTH / PERCENTILE	LENGTH / PERCENTILE	LENGTH / PERCENTILE	LENGTH / PERCENTILE	HEIGHT	HEIGHT
		HEAD CIRCUMFERENCE	HEAD CIRCUMFERENCE	HEAD CIRCUMFERENCE	HEAD CIRCUMFERENCE	BMI	BMI

Shaded boxes indicate the vaccine can be given during shown age range.

VISIT DATE

VISIT DATE

VISIT DATE

VISIT DATE

VISIT DATE

VISIT DATE

² Two doses given at least four weeks apart are recommended for children aged 6 months through 8 years of age who are getting a flu vaccine for the first time and for some other children in this age group.

³ Two doses of HepA vaccine are needed for lasting protection. The first dose of HepA vaccine should be given between 12 months and 23 months of age. The second dose should be given 6 months after the first dose. HepA vaccination may be given to any child 12 months and older to protect against HepA. Children and adolescents who did not receive the HepA vaccine and are at high-risk, should be vaccinated against HepA.

* Milestones adapted from AAP *Bright Futures: Guidelines for Health Supervision of Infants, Children, and Adolescents* Third Edition.

This is not an exhaustive list of milestones from 0-6 years. See more at www.cdc.gov/Milestones and download the Milestone Tracker App at www.cdc.gov/MilestoneTracker

If your child has any medical conditions that put him at risk for infections or is traveling outside the United States, talk to your child's doctor about additional vaccines that he may need.



U.S. Department of Health and Human Services
Centers for Disease Control and Prevention



American Academy of Pediatrics



DEDICATED TO THE HEALTH OF ALL CHILDREN™