



# Only Administer Abrysvo (Pfizer) Vaccine to Pregnant People



Two respiratory syncytial virus (RSV) vaccine products are available for use in the United States.



## Strategies to Help Prevent Errors



- Order and stock vaccine products that fit best with your patient population. Avoid stocking both products, if possible.



- If both RSV vaccine products are stocked, label the Arexvy (GSK) vaccine “Do NOT administer to pregnant people.”



- Educate staff on vaccine recommendations. If both RSV products are stocked, train staff about the differences in preparation and indications.



- Follow medication administration best practices – read and check the vaccine product label at least 3 times and ask another staff member to confirm that it is the correct vaccine product for the patient.



- If referring pregnant people to another vaccine provider, tell the provider to administer Abrysvo (Pfizer) vaccine and to confirm the vaccine product prior to administration.

## CDC Clinical Resources

[Healthcare Providers: RSV Vaccination for Pregnant People | CDC](#)

[RSV Vaccine Information Statement | CDC](#)

[Adult Immunization Schedule – Healthcare Providers | CDC](#)

[Use of the Pfizer Respiratory Syncytial Virus Vaccine During Pregnancy for the Prevention of Respiratory Syncytial Virus–Associated Lower Respiratory Tract Disease in Infants: Recommendations of the Advisory Committee on Immunization Practices – United States, 2023 | MMWR \(cdc.gov\)](#)

[Vaccine Administration: Preventing Vaccine Administration Errors \(cdc.gov\)](#)