

EBOLA BASICS

Ebola virus is **not** spread through

- **Casual contact**
- **Air**
- **Water**
- **Food grown or legally purchased in the U.S.**

How do you get the Ebola virus?

Direct contact with

- 1 Body fluids of a person who is sick with or has died from Ebola** (blood, vomit, urine, feces, sweat, semen, spit, other fluids)
- 2 Objects contaminated with the virus** (needles, medical equipment)
- 3 Infected fruit bats or primates** (apes and monkeys)
- 4 Possibly from contact with semen from a man who has recovered from Ebola** (by having oral, vaginal, or anal sex)

Early Symptoms

Symptoms can appear from 2 to 21 days after exposure.

- **Fever**
- **Stomach pain**
- **Headache**
- **Unexplained bleeding or bruising**
- **Diarrhea**
- **Muscle pain**
- **Vomiting**

When is someone able to spread the disease to others?

A person is only contagious after Ebola symptoms begin.



MONTH						
S	M	T	W	T	F	S
		1	2	3	4	5
6	7	8	9	10	11	12
13	14	15	16	17	18	19
20	21	22	23	24	25	26
27	28	29	30	31		

After 21 days, if an exposed person does not develop symptoms, they will not become sick with Ebola.

It is possible that Ebola can be spread through semen from men who have survived.



U.S. Department of Health and Human Services
Centers for Disease Control and Prevention

[cdc.gov/ebola](https://www.cdc.gov/ebola)

CS250531-H