

Preventing Sexual Violence

What is sexual violence?

Sexual violence is sexual activity when consent is not obtained or freely given. It is a serious public health problem in the United States that profoundly impacts lifelong health, opportunity, and well-being. Sexual violence impacts every community and affects people of all genders, sexual orientations, and ages. Anyone can experience or perpetrate sexual violence. The perpetrator of sexual violence is usually someone the victim knows, such as a friend, current or former intimate partner, coworker, neighbor, or family member.¹ Sexual violence can occur in person, online, or through technology, such as posting or sharing sexual pictures of someone without their consent, or non-consensual sexting.

How big is the problem?

Sexual violence affects millions of people each year in the United States. Researchers know the numbers underestimate this problem as many cases are unreported. Survivors may be ashamed, embarrassed, or afraid to tell the police, friends, or family about the violence. Survivors may also keep quiet because they have been threatened with further harm if they tell anyone or do not think anyone will help them. The data shows:

- **Sexual violence is common.** Over half of women and almost 1 in 3 men have experienced sexual violence involving physical contact during their lifetimes. One in 4 women and about 1 in 26 men have experienced completed or attempted rape. About 1 in 9 men were made to penetrate someone during his lifetime. Additionally, 1 in 3 women and about 1 in 9 men experienced sexual harassment in a public place.²
- **Sexual violence starts early.** More than 4 in 5 female rape survivors reported that they were first raped before age 25 and almost half were first raped as a minor (i.e., before age 18). Nearly 8 in 10 male rape survivors reported that they were made to penetrate someone before age 25 and about 4 in 10 were first made to penetrate as a minor.²
- **Sexual violence disproportionately affects some groups.** Women and racial and ethnic minority groups experience a higher burden of sexual violence. For example, more than 2 in 5 non-Hispanic American Indian or Alaska Native and non-Hispanic multiracial women were raped in their lifetime.²
- **Sexual violence is costly.** Recent estimates put the lifetime cost of rape at \$122,461 per victim, including medical costs, lost productivity, criminal justice activities, and other costs.³

Over **half** of women have experienced sexual violence involving physical contact during her lifetime.



Almost **1 in 3** men have experienced sexual violence involving physical contact during his lifetime.



Estimated Lifetime
Cost of Rape



Centers for Disease
Control and Prevention
National Center for Injury
Prevention and Control

What are the consequences?

Sexual violence consequences are physical, like bruising and genital injuries, sexually transmitted infections, and pregnancy (for women)² and psychological, such as depression, anxiety, and suicidal thoughts.⁴

The consequences may be chronic. Survivors may suffer from post-traumatic stress disorder, experience re-occurring reproductive, gastrointestinal, cardiovascular and sexual health problems.⁴


Sexual violence is also linked to negative health behaviors. Survivors are more likely to smoke, abuse alcohol, use drugs, and engage in risky sexual activity.⁵

The trauma from sexual violence may impact a survivor's employment in terms of time off from work, diminished performance, job loss, or inability to work. These disrupt earning power and have a long-term effect on the economic well-being of survivors and their families. Coping and completing everyday tasks after victimization can be challenging. Survivors may have difficulty maintaining personal relationships, returning to work or school, and regaining a sense of normalcy.⁴

Additionally, sexual violence is connected to other forms of violence. For example, girls who have been sexually abused are more likely to experience additional sexual violence and other violence types and become victims of intimate partner violence in adulthood.⁶ Bullying perpetration in early middle school is associated with sexual harassment perpetration in high school.⁷


How can we prevent sexual violence?

Certain factors may increase or decrease the risk for perpetrating or experiencing sexual violence. To prevent sexual violence, we must understand and address the factors that put people at risk for or protect them from violence.⁶ We must also understand how historical trauma and structural inequities impact health.² CDC developed *STOP SV: A Technical Package to Prevent Sexual Violence* to help communities use the best available evidence to prevent sexual violence. This resource is available in English and Spanish and can impact individual behaviors and relationship, family, school, community, and societal factors that influence risk and protective factors for violence. Different violence types are connected and often share root causes. Sexual violence is linked to other violence types through shared risk and protective factors. Addressing and preventing one violence type may help prevent other violence types.



STOP SV: A Technical Package to Prevent Sexual Violence

A **technical package** is a collection of strategies based on the best available evidence to prevent or reduce public health problems. The **strategy** lays out the direction and actions to prevent sexual violence. The **approach** includes the specific ways to advance the strategy through programs, policies and practices. The **evidence** for each of the approaches in preventing sexual violence and associated risk factors is also included.



References

- Basile KC, Smith SG, Breiding MJ, Black M C, & Mahendra, R. (2014). Sexual violence surveillance: uniform definitions and recommended data elements, Version 2.0. Atlanta (GA): National Center for Injury Prevention and Control, Centers for Disease Control and Prevention.
- Basile KC, Smith SG, Kresnow M, Khatiwada S, & Leemis RW. (2022). The National Intimate Partner and Sexual Violence Survey: 2016/2017 Report on Sexual Violence. Atlanta, GA: National Center for Injury Prevention and Control, Centers for Disease Control and Prevention.
- Peterson C, DeGue S, Florence C, Lokey C. (2017). Lifetime Economic Burden of Rape in the United States. *American Journal of Preventive Medicine* 52(6): 691-701.
- Basile KC and Smith SG. (2011). Sexual Violence Victimization of Women: Prevalence, Characteristics, and the Role of Public Health and Prevention. *American Journal of Lifestyle Medicine* (5): 407-417.
- Basile KC, Clayton HB, Rostad WL, & Leemis RW. (2020). Sexual violence victimization of youth and health risk behaviors. *American Journal of Preventive Medicine*, 58(4), 570-579.
- Preventing Multiple Forms of Violence: A Strategic Vision for Connecting the Dots. (2016). Atlanta, GA: National Center for Injury Prevention and Control, Centers for Disease Control and Prevention.
- Espelage DL, Basile KC, Hamburger ME. (2012). Bullying perpetration and subsequent sexual violence perpetration among middle school students. *Journal of Adolescent Health* 50(1): 60-65.