

Vision Health Disparities in the United States by Race/Ethnicity, Education, and Economic Status: Findings From Two Nationally Representative Surveys

XINZHI ZHANG, MARY FRANCES COTCH, ASEL RYSKULOVA, SUSAN A. PRIMO, PARVATHY NAIR, CHIU-FANG CHOU, LINDA S. GEISS, LAWRENCE E. BARKER, AMANDA F. ELLIOTT, JOHN E. CREWS, AND JINAN B. SAADDINE

- **PURPOSE:** To assess vision health disparities in the United States by race/ethnicity, education, and economic status.
- **DESIGN:** Cross-sectional, nationally representative samples.
- **METHODS:** We used national survey data from the National Health and Nutrition Examination Survey (NHANES) and the National Health Interview Survey (NHIS). Main outcome measures included, from NHANES, age-related eye diseases (ie, age-related macular degeneration [AMD], cataract, diabetic retinopathy [DR], glaucoma) and from NHIS, eye care use (ie, eye doctor visits and cannot afford eyeglasses when needed) among those with self-reported visual impairment. The estimates were age- and sex-standardized to the 2000 US Census population. Linear trends in the estimates were assessed by weighted least squares regression.
- **RESULTS:** Non-Hispanic whites had a higher prevalence of AMD and cataract surgery than non-Hispanic blacks, but a lower prevalence of DR and glaucoma (all $P < .001$ in NHANES 2005-2008). From 1999 to 2008, individuals with less education (ie, <high school vs >high school) and lower income (poverty income ratio [PIR] <1.00 vs ≥ 4.00) were consistently less likely to have had an eye care visit in the past 12 months compared with their counterparts (all $P < .05$). During this period, inability to afford needed eyeglasses increased among non-Hispanic whites and Hispanics (trend $P = .004$ and $P = .007$; respectively), those with high school education (trend $P = .036$), and those with PIR 1.00-1.99 (trend $P < .001$).

Accepted for publication Aug 26, 2011.

From the Division of Diabetes Translation, National Center for Chronic Disease Prevention and Health Promotion, Centers for Disease Control and Prevention, Atlanta, Georgia (X.Z., P.N., C.F.C., L.S.G., L.E.B., A.F.E., J.E.C., J.B.S.); Division of Epidemiology and Clinical Applications, National Eye Institute, National Institutes of Health, Bethesda, Maryland (M.F.C.); National Center for Health Statistics, Centers for Disease Control and Prevention, Hyattsville, Maryland (A.R.); and Emory Eye Center, Emory University School of Medicine, Atlanta, Georgia (S.A.P.).

Inquiries to Xinzhi Zhang, Vision Health Initiative, Division of Diabetes Translation, National Center for Chronic Disease Prevention and Health Promotion, Centers for Disease Control and Prevention, 4770 Buford Hwy, N.E. (K-10), Atlanta, GA 30341-3727; e-mail: XZhang4@cdc.gov

- **CONCLUSIONS:** Observed vision health disparities suggest a need for educational and innovative interventions among socioeconomically disadvantaged groups. (Am J Ophthalmol 2012;154:S53-S62. © 2012 Published by Elsevier Inc.)

VISION LOSS IS A SERIOUS PUBLIC HEALTH CHALLENGE today, and global efforts to reduce avoidable blindness by 2020 are underway.^{1,2} In the United States, age-related eye diseases (ie, age-related macular degeneration [AMD], cataract, diabetic retinopathy [DR], glaucoma) and other eye conditions are common causes of vision loss.³⁻⁷ More than 3.3 million Americans 40 years or older are either legally blind (0.9 million, 0.8%) or have low vision (ie, less than 20/40 best-corrected vision in the better-seeing eye; 2.4 million, 2.0%) caused mostly by age-related eye diseases.⁸ By 2020, the number of people with eye diseases and associated vision loss could increase by 50% or more because of the aging of the US population.³⁻⁷

Blindness and low vision often affect people's ability to read, walk, drive, or conduct other daily activities.^{9,10} Poor vision also results in an increased risk for falls and injuries,^{11,12} social isolation, depression,^{13,14} loss of productivity,^{15,16} and, ultimately, an increased risk for morbidity or premature death.¹⁷⁻²⁰ Moreover, many eye diseases are asymptomatic in the early stages, when treatment may forestall or delay disease onset and prevent vision loss. Given the rapidly aging US population, the growing obesity epidemic, and the increasing prevalence of diabetes, evidence is mounting that vision loss will become an even greater problem for the United States in the future.

Vision objectives are included in *Healthy People 2010* and *Healthy People 2020*, a national disease prevention initiative that identifies opportunities to improve the health of all Americans.^{21,22} One of the overarching goals of *Healthy People 2010/2020* is to eliminate health disparities, including differences that occur by sex, race/ethnicity, education, income, disability, geographic location, or sexual orientation. Addressing vision health disparities would help enhance public health programs and close the gaps to improve the nation's vision and eye health.^{23,24} A

national and state surveillance system using national, state, and local surveys; birth and death certificates; and Medicare, Medicaid, and managed care data helps the nation's policy makers analyze, interpret, and report on vision loss, risk factors, care practices, morbidity, and mortality.

This report describes the use of national data to assess the prevalence of major age-related eye diseases and eye care utilization; examines disparities by race/ethnicity, education, and economic status; explores the strengths and weaknesses of using existing national data sets to identify vision health disparities; and helps future surveillance activities to guide public health priorities and policies.

METHODS

• **DATA SOURCES:** *National Health and Nutrition Examination Survey.* The National Health and Nutrition Examination Survey (NHANES) (www.cdc.gov/nchs/nhanes.htm) is a national representative survey conducted by the National Center for Health Statistics, the Centers for Disease Control and Prevention (CDC). The NHANES was conducted during 1960-1962 (NHES I), 1963-1965 (NHES II), 1966-1970 (NHES III), 1971-1974 (NHANES I), 1976-1980 (NHANES II), 1982-1984 (HHANES), and 1988-1994 (NHANES III). Since 1999, the NHANES has been conducted continuously. The NHANES survey protocols were approved by a human subjects review board, and written informed consent was obtained from all participants.

The NHANES data consist of samples of the US noninstitutionalized civilian population, which were obtained by using a stratified multistage probability design with planned oversampling of certain age and minority groups. Participants were interviewed at home to obtain sociodemographic, medical, and family information; those who reported to the Mobile Examination Center received a detailed medical examination. The NHANES III also included a retinal photography protocol, which was used to determine the presence and severity of AMD and DR. Beginning in 1999, visual acuity testing was included. The ophthalmic component of digital retinal images and frequency-doubling technology (FDT) for participants aged 40 or older was added to the 2005-2008 NHANES. Details about the methodology of these 2 new technologies have been described elsewhere.^{25,26}

National Health Interview Survey. Conducted annually since 1957 by the National Center for Health Statistics, the National Health Interview Survey (NHIS) (<http://www.cdc.gov/nchs/nhis.htm>) uses a stratified, cross-sectional, multistage probability sample to derive estimates for the civilian, noninstitutionalized population. A major redesign occurs every 10 to 15 years. Collected through an in-person household interview, the NHIS gathers data on illnesses, injuries, activity limitation, chronic conditions,

health insurance coverage, use of health care, and other health topics. Protocols were approved by a human subjects review board, and written informed consent was obtained from all participants. One adult and 1 child are randomly selected from each family for a detailed health profile. Basic health and demographic information are obtained by interviewing all family members. Self-reported visual function questions were added to NHIS starting in 1986. An eye care use question was added starting in 1997. In 2002 and 2008, additional questions were included in NHIS as a vision health supplement to assess and monitor progress of the Healthy People 2010 and 2020 vision objectives.

• **MEASUREMENTS:** We used data from NHANES III (1988-1994, n = 5704) and NHANES 2005-2008 (n = 8208) to assess eye disease disparity among US adults aged 40 years or older. The NHANES III response rate for the interview sample aged 40 years or older was 79% and 67% for the examined sample. The NHANES 2005-2008 response rate for the interview sample aged 40 years or older was 71% and 69% for the examined sample. As defined previously by the University of Wisconsin's Ocular Epidemiologic Reading Center, Madison,²⁶ participants with diabetes (defined as self-report of a previous diagnosis of the disease by a clinician excluding gestational diabetes mellitus or hemoglobin A1c of 6.5% or greater) were considered to have DR (NHANES III and NHANES 2005-2008) if retinal images revealed the presence of 1 or more retinal microaneurysms or retinal blot hemorrhages with or without more severe lesions. By the Wisconsin Reading Center definition,²⁷ participants were considered to have AMD on the basis of the drusen size, type, and area; increased retinal pigment; retinal pigment epithelial depigmentation; pure geographic atrophy; and signs of exudative macular degeneration found in their retinal image. One eye was photographed in NHANES III and both in NHANES 2005-2008. To be comparable to estimates from NHANES III, we estimated the prevalences of DR and AMD in NHANES 2005-2008 by using the right eye if the last digit of the participant identification number was even and the left eye if it was odd. Cataract and glaucoma were defined on the basis of the participant self-report in NHANES III and NHANES 2005-2008. As a proxy, the presence of cataract was declared if the participant answered affirmatively to the question, "Have you ever had a cataract operation?" Likewise, we define glaucoma as answering yes to the question, "Have you ever been told by an eye doctor that you have glaucoma, sometimes called high pressure in your eyes?"

We used 10 consecutive years (1999-2008) of NHIS data to evaluate eye care use by race/ethnicity, education, and economic status among the US civilian, noninstitutionalized population aged 18 years or older. The final response rate for the adult sample person component of the NHIS ranged from 62.6% in 2008 to 74.3% in 2002. Eye

care use variables included respondent self-report of having visited an eye doctor in the past 12 months and being unable to afford eyeglasses when needed. These measures were chosen because eye doctor visits capture health care use, whereas inability to afford eyeglasses addresses financial barriers and the effect of reduced visual acuity caused by uncorrected refractive error on vision-related quality of life.²⁴ The analyses of annual eye care use indicators were restricted to those who reported visual impairment (n = 29 175) because annual eye care visits for those only needing new eyeglass prescriptions but no vision function loss (ie, without visual impairment) is not considered cost-effective.²⁸ Respondents were classified as having visual impairment if they answered yes to the question, "Do you have any trouble seeing, even when wearing glasses or contact lenses?"

In this study, we specifically focused on racial/ethnic and socioeconomic (educational attainment, income level) disparities in the major eye diseases and eye care use. Because of the survey design, we divided racial/ethnic status into non-Hispanic white, non-Hispanic black, and Mexican American in NHANES and non-Hispanic white, non-Hispanic black, and Hispanic in NHIS. Educational attainment was categorized as less than high school, high school graduate, and more than high school. Income status was measured by using the poverty-to-income ratio (PIR), an index of total income assessed in relation to federal poverty thresholds that vary by family size and composition.²⁹ A PIR score of 1.00 is defined as the official federal poverty threshold level. We categorized PIR as less than 1.00, 1.00 to 1.99, 2.00 to 3.99, and 4.00 or more.

• **STATISTICAL ANALYSIS:** The data were analyzed using SAS 9.1.3 (SAS Institute, Cary, North Carolina, USA) and SUDAAN 9.0 (Research Triangle Institute, Research Triangle Park, North Carolina, USA). All analyses accounted for the complex multistage sampling, clustering, and stratification design implemented by the NHANES and NHIS. Taylor linearization was used to estimate variances.³⁰ All prevalence estimates were age- and sex-standardized to the 2000 US Census population.³¹ Estimates with a relative standard error more than 30% were considered not reliable and therefore not reported. Linear trends in the prevalence estimates were assessed by weighted least squares regression. P values less than .05 were considered statistically significant.

RESULTS

• **PREVALENCE OF EYE DISEASES BY RACE/ETHNICITY, EDUCATIONAL ATTAINMENT, AND ECONOMIC STATUS:** The observed racial/ethnic differences of DR among non-Hispanic whites and non-Hispanic blacks remained during the decade (Table 1). The prevalence of DR was significantly higher among non-Hispanic blacks than among

non-Hispanic whites in NHANES III and NHANES 2005-2008 (4.9% vs 1.9%, 8.7% vs 3.2%; both $P < .001$). Some significant differences in age- and sex-standardized prevalence of AMD or cataract surgery by race/ethnicity were also present. For example, the prevalence of AMD was higher among non-Hispanic whites than among non-Hispanic blacks in NHANES III and NHANES 2005-2008 (12.9% vs 9.9%, 8.3% vs 3.1%; $P = .017$ and $P < .001$, respectively). In addition, non-Hispanic whites reported a higher prevalence of cataract surgery than non-Hispanic blacks in NHANES III and NHANES 2005-2008 (19.3% vs 16.4%, 18.4% vs 13.5%; $P = .018$ and $P < .001$, respectively). Non-Hispanic blacks had a significantly higher age- and sex-standardized prevalence of glaucoma in NHANES 2005-2008 compared to non-Hispanic white (11.5% vs 6.9%; $P < .001$) and Mexican-American (11.5% vs 6.5%; $P = .006$) participants.

In both NHANES III and NHANES 2005-2008, the age- and sex-standardized prevalence of DR was significantly higher among those with less than a high school education than among those with more than a high school education in NHANES III and NHANES 2005-2008 (2.8% vs 1.1%, 6.9% vs 2.7%; both $P = .001$). No significant difference by educational attainment was observed in the prevalence of AMD, glaucoma, or cataract surgery for either time.

Although no significant differences for income were observed in NHANES III, the age- and sex-standardized prevalence of DR in NHANES 2005-2008 was significantly higher among individuals at the lowest income level ($PIR < 1.00$) compared to those at the highest income levels ($PIR \geq 4.00$) (5.1% vs 2.2%; $P = .030$). The age- and sex-standardized prevalence of AMD in NHANES III was also higher among those at the lowest income level than those at the highest income levels (17.9% vs 11.5%; $P = .030$), although the difference was not significant in NHANES 2005-2008 (10.4% vs 6.8%; $P = .057$). No significant difference in the prevalence of glaucoma or cataract surgery by income level was observed in NHANES III or NHANES 2005-2008.

• **EYE CARE USE BY RACE/ETHNICITY, EDUCATIONAL ATTAINMENT, AND ECONOMIC STATUS:** Among adults with reported visual impairment, no increasing or decreasing trend of visiting an eye doctor in the past 12 months was found by race/ethnicity, education attainment, or income status from 1999 to 2008 (Table 2). For several years, we observed some racial/ethnic differences in eye doctor visits. For example, in 2008, non-Hispanic whites were more likely to visit an eye doctor than Hispanics (52.6% vs 36.9%; $P < .001$). Moreover, from 1999 to 2008, individuals with less than a high school education were consistently less likely to have had an eye care visit in the past 12 months than those with more than a high school education (all $P < .01$). During this period (except

TABLE 1. Age- and Sex-Standardized Prevalence of Age-Related Eye Diseases Among Adults Aged 40 Years or Older, by Race/Ethnicity, Education, and Income, NHANES III and NHANES 2005-2008^a

| | Diabetic Retinopathy | | | Age-Related Macular Degeneration | | | Glaucoma | | | Cataract Surgery | | | | | | |
|-------------------------|----------------------|------------|------------------|----------------------------------|------|------------------|-------------------------|------------|------------------|------------------|------|------------------|-------|-------------|------|-------------|
| | NHANES III | | NHANES 2005-2008 | NHANES III | | NHANES 2005-2008 | NHANES III ^b | | NHANES 2005-2008 | NHANES III | | NHANES 2005-2008 | | | | |
| | % | 95% CI | % | 95% CI | % | 95% CI | % | 95% CI | % | 95% CI | % | 95% CI | | | | |
| Non-Hispanic white | 1.9 | (1.3-2.6) | 3.2 | (2.5-4.2) | 12.9 | (11.5-14.4) | 8.3 | (7.0-9.8) | N/A | N/A | 6.9 | (5.9-8.0) | 19.3 | (17.6-21.1) | 18.4 | (16.8-20.1) |
| Non-Hispanic black | 4.9 | (3.9-6.0) | 8.7 | (6.6-11.5) | 9.9 | (8.1-11.9) | 3.1 | (1.9-4.9) | N/A | N/A | 11.5 | (9.6-13.8) | 16.4 | (14.5-18.4) | 13.5 | (11.3-16.1) |
| Mexican American | 6.8 | (4.6-10.0) | 5.0 | (3.1-7.8) | 10.4 | (7.9-13.6) | 9.8 | (6.5-14.3) | N/A | N/A | 6.5 | (4.5-9.3) | 20.5 | (18.6-22.6) | 16.4 | (13.2-20.2) |
| P, white vs black | <.001 | | <.001 | | .017 | | <.001 | | N/A | N/A | N/A | | <.001 | | .018 | <.001 |
| P, white vs Mexican | .001 | | .136 | | .072 | | .449 | | N/A | N/A | N/A | .757 | | .283 | | .274 |
| P, black vs Mexican | .17 | | .068 | | .756 | | .003 | | N/A | N/A | N/A | .006 | | .002 | | .182 |
| < High school | 2.8 | (2.2-3.6) | 6.9 | (5.1-9.2) | 12.8 | (11.2-14.6) | 6.7 | (5.1-9.1) | N/A | N/A | 8.2 | (6.7-9.8) | 19.5 | (17.8-21.4) | 19.1 | (16.3-22.3) |
| High school | 3.0 | (2.0-4.4) | 3.5 | (2.3-5.2) | 11.7 | (9.8-14.0) | 7.9 | (5.8-10.5) | N/A | N/A | 7.0 | (5.0-9.7) | 17.8 | (14.7-21.3) | 17.1 | (14.7-19.9) |
| > High school | 1.1 | (0.6-2.0) | 2.7 | (2.1-3.6) | 13.6 | (11.5-16.0) | 8.3 | (6.7-10.2) | N/A | N/A | 7.1 | (5.7-8.7) | 20.0 | (17.6-22.7) | 17.5 | (15.7-19.4) |
| P, < HS vs > HS | .001 | | .001 | | .490 | | .210 | | N/A | N/A | N/A | .397 | | .763 | | .309 |
| PIR ^c < 1.00 | 2.4 | (1.6-3.5) | 5.1 | (3.4-7.8) | 17.9 | (13.8-23.0) | 10.4 | (7.6-14.1) | N/A | N/A | 7.4 | (5.9-9.1) | 17.2 | (14.2-20.6) | 16.7 | (13.7-20.3) |
| PIR 1.00-1.99 | 2.6 | (1.7-3.8) | 4.8 | (3.3-6.9) | 12.1 | (10.2-14.2) | 7.1 | (5.1-9.8) | N/A | N/A | 8.3 | (7.0-9.8) | 21.1 | (19.1-23.4) | 21.3 | (19.0-23.6) |
| PIR 2.00-3.99 | 2.2 | (1.3-3.6) | 3.8 | (2.5-5.7) | 13 | (10.9-15.5) | 8.3 | (6.9-10.1) | N/A | N/A | 7.8 | (5.6-10.8) | 17.5 | (15.1-20.1) | 18.5 | (16.4-20.8) |
| PIR ≥ 4.00 | 1.9 | (1.1-3.2) | 2.2 | (1.4-3.5) | 11.5 | (9.6-13.7) | 6.8 | (5.0-9.2) | N/A | N/A | 6.1 | (4.4-8.4) | 19.7 | (17.3-22.3) | 14.0 | (11.4-17.0) |
| PIR < 1.00 vs ≥ 4.00 | .405 | | .030 | | .030 | | .057 | | N/A | N/A | N/A | .394 | | .23 | | .116 |

CI = confidence interval; HS = high school; N/A = not applicable; NHANES = National Health and Nutrition Examination Survey; PIR = Poverty Income Ratio.

^an = 5704 in NHANES III (1988-94); n = 8202 in NHANES 2005-2008.

^bData related to glaucoma were not available from NHANES III.

^cPIR is an index of total income assessed in relation to federal poverty thresholds that vary by family size and composition. A PIR score of 1.00 is defined as the official federal poverty threshold level.

TABLE 2. Age- and Sex-Standardized Prevalence of Self-Reported Annual Eye-Care Visits Among Adults Aged 18 Years or Older With Self-Reported Visual Impairment, by Race/Ethnicity, Education, and Income, National Health Interview Survey, 1999-2008^a

| Race/Ethnicity | 1999 | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | Trend |
|------------------------|-----------------------|-----------------------|-----------------------|-----------------------|-----------------------|-----------------------|-----------------------|-----------------------|-----------------------|-----------------------|----------------------|
| | % | % | % | % | % | % | % | % | % | % | P Value ^b |
| Non-Hispanic white | 51.1 (47.0 - 55.1) | 53.5 (49.8 - 57.2) | 51.4 (47.7 - 55.0) | 54.2 (50.3 - 58.0) | 52.7 (48.9 - 56.4) | 55.0 (51.4 - 58.6) | 54.5 (50.6 - 58.3) | 50.9 (46.2 - 55.5) | 51.7 (47.5 - 55.9) | 52.6 (47.8 - 57.4) | .807 |
| Non-Hispanic black | 47.8 (39.8 - 56.0) | 52.7 (43.5 - 61.8) | 48.1 (39.9 - 56.5) | 46.5 (37.8 - 55.4) | 43.4 (35.4 - 51.7) | 44.6 (37.3 - 52.1) | 43.0 (35.0 - 51.4) | 44.8 (36.6 - 53.3) | 48.4 (40.1 - 56.8) | 47.2 (37.8 - 56.9) | .272 |
| Hispanic | 39.5 (28.3 - 51.8) | 43.6 (36.3 - 51.1) | 44.5 (37.4 - 51.8) | 35.8 (26.6 - 46.2) | 42.7 (31.9 - 54.2) | 50.2 (42.7 - 57.8) | 43.1 (34.5 - 52.1) | 45.3 (36.2 - 54.8) | 43.5 (33.0 - 54.7) | 36.9 (30.1 - 44.2) | .630 |
| P, white vs black | .483 | .878 | .485 | .127 | .047 | .014 | .016 | .199 | .493 | .308 | |
| P, white vs Hispanic | .070 | .018 | .103 | .001 | .102 | .265 | .020 | .308 | .173 | <.001 | |
| P, black vs Hispanic | .261 | .142 | .518 | .116 | .926 | .286 | .991 | .931 | .485 | .101 | |
| Education | | | | | | | | | | | |
| < High school | 38.3 (31.9 - 45.1) | 43.5 (37.2 - 50.0) | 43.7 (38.2 - 49.4) | 38.2 (32.6 - 44.2) | 37.2 (31.8 - 43.0) | 42.5 (36.5 - 48.7) | 39.7 (34.0 - 45.6) | 42.6 (35.8 - 49.6) | 38.5 (32.1 - 45.4) | 35.4 (28.6 - 42.8) | .330 |
| High school | 52.9 (46.7 - 59.1) | 47.5 (41.9 - 53.2) | 44.5 (39.3 - 49.9) | 51.5 (45.3 - 57.7) | 52.8 (47.3 - 58.2) | 50.9 (44.9 - 56.8) | 53.1 (47.7 - 58.5) | 46.1 (40.2 - 52.0) | 52.6 (46.5 - 58.6) | 54.9 (48.2 - 61.5) | .312 |
| high school | 54.1 (49.0 - 59.0) | 63.1 (59.3 - 66.8) | 58.8 (54.1 - 63.4) | 60.9 (56.0 - 65.7) | 59.0 (54.5 - 63.4) | 60.9 (56.3 - 65.3) | 59.2 (54.3 - 64.0) | 55.8 (49.4 - 62.0) | 57.0 (51.2 - 62.6) | 59.0 (53.8 - 64.0) | .637 |
| P, <HS vs HS | <.001 | <.001 | <.001 | <.001 | <.001 | <.001 | <.001 | .006 | <.001 | <.001 | |
| PIR^c | | | | | | | | | | | |
| <1.00 | 44.2 (35.9 - 52.8) | 41.0 (32.7 - 49.9) | 47.1 (40.6 - 53.6) | 36.7 (28.3 - 46.0) | 39.5 (32.0 - 47.6) | 47.7 (40.4 - 55.2) | 39.1 (30.8 - 48.1) | 41.0 (32.5 - 50.1) | 38.7 (30.6 - 47.5) | 33.3 (23.6 - 44.7) | .147 |
| 1.00 - 1.99 | 44.8 (37.9 - 51.9) | 51.7 (45.6 - 57.7) | 43.0 (37.1 - 49.2) | 46.7 (40.1 - 53.5) | 48.3 (42.2 - 54.5) | 43.4 (36.8 - 50.3) | 44.8 (38.1 - 51.8) | 44.3 (37.1 - 51.7) | 47.2 (39.5 - 55.1) | 44.3 (37.1 - 51.9) | .393 |
| 2.00 - 3.99 | 49.7 (43.6 - 55.8) | 49.8 (44.6 - 55.0) | 51.4 (45.9 - 56.8) | 50.5 (44.9 - 56.0) | 51.9 (46.2 - 57.6) | 51.8 (45.3 - 58.2) | 55.9 (49.8 - 61.7) | 51.2 (42.9 - 59.4) | 49.3 (42.6 - 56.0) | 52.6 (44.5 - 60.6) | .393 |
| ≥4.00 | 55.2 (46.9 - 63.2) | 60.9 (55.5 - 66.1) | 58.5 (52.8 - 64.0) | 68.0 (61.8 - 73.6) | 57.5 (51.2 - 63.6) | 64.6 (57.5 - 71.1) | 60.0 (53.2 - 66.4) | 58.2 (49.0 - 66.9) | 62.5 (55.4 - 69.0) | 65.7 (57.9 - 72.7) | .342 |
| P, <1 vs ≥4 | .070 | <.001 | .009 | <.001 | .002 | .001 | <.001 | .009 | <.001 | <.001 | |

CI = confidence interval; HS = high school; PIR, Poverty Income Ratio.

^an = 29 175. We categorized adults who reported experiencing any trouble seeing despite wearing contact lens or glasses as having visual impairment.

^bLinear trend.

^cPIR is an index of total income assessed in relation to federal poverty thresholds that vary by family size and composition. A PIR score of 1.00 is defined as the official federal poverty threshold level.

year 1999), individuals with PIR of less than 1.00 were consistently less likely to have an eye care visit than those with PIR of 4.00 or more (all $P < .05$).

From 1999 to 2008, the inability to afford eyeglasses increased among non-Hispanic whites and Hispanics (trend $P = .004$ and $P = .007$; respectively), those with high school education (trend $P = .036$), and those with income PIR 1.00-1.99 (trend $P = .001$) (Table 3). We also observed some significant racial/ethnic differences in the inability to afford eyeglasses in several years. For example, in 2008, Hispanics were more likely to report being unable to afford eyeglasses when needed than non-Hispanic whites and non-Hispanic blacks (26.7% vs 16.0% and 15.3%; both $P = .020$). Individuals with less than a high school education were more likely to report being unable to afford eyeglasses when needed than those with more than a high school education in most years (all $P < .01$) except in year 2002 ($P = .075$) and year 2007 ($P = .069$). From 1999 to 2008, impoverished individuals (PIR < 1.00) were consistently more likely to report being unable to afford eyeglasses when needed compared to those at the highest income level (all $P < .001$).

DISCUSSION

• **VISION HEALTH DISPARITIES:** We report observed disparities by race/ethnicity, education, and economic status in the prevalence of the major eye diseases and the use of eye care services. We also found that these disparities existed previously and have persisted over a decade.

Our survey-based evidence of racial/ethnic differences for the effects of major eye diseases is consistent with previous literature.^{27,32-34} We found that the prevalence of DR and glaucoma among Americans 40 years of age or older is greater in non-Hispanic blacks than in non-Hispanic whites. We also found that non-Hispanic whites had a higher prevalence of AMD and cataract surgery than non-Hispanic blacks. Previous meta-analyses of findings from individual population-based studies within geographically circumscribed settings also reported similar findings.³⁻⁶ Furthermore, we found that DR prevalence was higher among the less educated and the impoverished, which underscores a critical need for services, particularly regular eye examinations.

Affordability, continuity, and regular sources of care, as well as physician advice, have been shown to be strongly associated with receipt of needed eye care services.^{35,36} Reviewing NHIS data from 1999 to 2008, we found that individuals with less education and lower income were both less likely to visit an eye doctor within the past year and less able to afford eyeglasses when needed. A previous study similarly found that those with more education are more likely to see an eye care professional compared to those less educated.³⁷ More than 50% of Medicare beneficiaries have not visited an eye care provider or received

an annual eye examination.³⁸ Lack of awareness about vision health is a major problem, especially among low-income, minority, and uninsured families who are at highest risk of not accessing vision screening programs.³⁹ Furthermore, increasing trends of inability to afford eyeglasses among those with a high school education and those with PIR of 1.00 to 1.99 suggest that public health interventions to improve vision and eye health should focus more on those at the relatively lower end (ie, \leq high school education or PIR < 2.00 vs high school education or PIR ≥ 2.00) of the socioeconomic spectrum.

• **STRENGTH AND WEAKNESS OF CURRENT NATIONAL DATA:** NHANES and NHIS data are widely used as major data sources to assess *Healthy People 2010* and *Healthy People 2020* vision health goals and objectives. They can provide national population-based estimates of age-related eye diseases and eye care use. Although all eye disease prevalence estimates in *Healthy People 2010/2020* come from the NHIS self-reported vision supplement questions, our estimates from clinical measures (DR and AMD) increased the reliability of the diagnosis and provided similar findings on vision health disparities by race/ethnicity, education, and economic status. Compared with the single-eye protocol used in NHANES III, the protocol begun in the 2005 NHANES of digital fundus images of 2 photographs for each eye has improved the ability to detect disease. Previous reports using data on both eyes from participants in NHANES 2005-2008 reflect more realistic estimates of the true eye disease prevalence. They indicate racial/ethnic disparities in DR²⁶ and AMD²⁷ and are consistent with what we report by using data from only 1 eye per participant. Combined with other population-based studies and state or local health surveys, an integrated national vision health surveillance system can provide baseline and trend data for vision health disparities and assist policy decision making.

Our study has several limitations. National surveys such as NHANES and NHIS do not include residents in long-term care facilities (eg, nursing homes), persons on active duty with the Armed Forces, prisoners, and US nationals living in foreign countries. A previous study has shown a higher prevalence of eye disease in nursing home residents than in community-dwelling older adults.⁴⁰ In addition, much of the available data is based on self-reports. More research on the reliability and validity of self-reported measures and ways to improve their collection is needed. The self-reported data used in this analysis are subject to many potential sources of bias, including recall bias, social desirability bias, and access to health care biases. The type and magnitude of these biases are difficult to evaluate. Thus, it is likely that our estimates of the rates of glaucoma, cataract, and use of eye care are negatively biased. Glaucoma is known to be persistently under-reported and under-diagnosed. Using self-reported cataract extraction surgery as a proxy for the presence of cataract provides a rough estimate of clinically significant cataract but underestimates

TABLE 3. Age- and Sex-Standardized Prevalence of Being Unable to Afford Eyeglasses When Needed Among Adults Aged 18 Years or Older With Self-Reported Visual Impairment, by Race/Ethnicity, Education, and Income, National Health Interview Survey, 1999-2008^a

| | 1999 | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | Trend |
|------------------------|-----------------------|-----------------------|-----------------------|-----------------------|-----------------------|-----------------------|-----------------------|-----------------------|-----------------------|-----------------------|----------------------|
| | % | % | % | % | % | % | % | % | % | % | P Value ^b |
| Race/Ethnicity | | | | | | | | | | | |
| Non-Hispanic white | 11.2 (9.5 - 13.1) | 15.2 (13.1 - 17.6) | 12.7 (10.8 - 14.9) | 12.7 (10.9 - 14.8) | 14.5 (12.3 - 16.9) | 15.1 (12.8 - 17.7) | 17.1 (14.6 - 19.8) | 16.1 (12.5 - 20.5) | 17.9 (15.1 - 21.1) | 16.0 (13.3 - 19.2) | .004 |
| Non-Hispanic black | 17.7 (12.6 - 24.3) | 26.7 (21.1 - 33.2) | 13.7 (9.1 - 20.2) | 17.4 (11.9 - 24.7) | 17.0 (12.1 - 23.3) | 22.5 (17.1 - 29.2) | 21.4 (15.6 - 28.5) | 16.6 (12.1 - 22.4) | 18.1 (12.9 - 24.9) | 15.3 (11.1 - 20.7) | .451 |
| Hispanic | 12.4 (8.9 - 17.0) | 9.7 (7.1 - 13.0) | 16.9 (11.8 - 23.6) | 19.0 (12.1 - 28.7) | 23.1 (17.0 - 30.6) | 21.7 (17.3 - 26.8) | 17.4 (13.4 - 22.3) | 23.2 (15.8 - 32.7) | 16.7 (10.8 - 25.0) | 26.7 (19.1 - 36.1) | .007 |
| P, white vs black | .035 | .001 | .743 | .162 | .417 | .026 | .231 | .876 | .941 | .803 | |
| P, white vs Hispanic | .588 | .003 | .194 | .145 | .019 | .016 | .898 | .124 | .775 | .020 | |
| P, black vs Hispanic | .151 | <.001 | .447 | .759 | .171 | .833 | .317 | .182 | .764 | .020 | |
| Education | | | | | | | | | | | |
| < High school | 18.7 (14.2 - 24.2) | 21.0 (16.5 - 26.4) | 16.3 (12.8 - 20.5) | 16.5 (12.6 - 21.3) | 22.7 (18.4 - 27.6) | 22.5 (18.5 - 27.0) | 24.9 (20.3 - 30.2) | 27.7 (20.7 - 35.9) | 19.4 (14.9 - 24.9) | 25.8 (19.6 - 33.2) | .071 |
| High school | 11.0 (8.9 - 13.5) | 15.4 (12.2 - 19.2) | 16.4 (13.1 - 20.4) | 14.0 (10.5 - 18.3) | 14.2 (11.6 - 17.2) | 13.8 (11.0 - 17.3) | 15.2 (12.0 - 19.0) | 13.9 (10.3 - 18.6) | 20.3 (16.2 - 25.3) | 17.4 (13.2 - 22.6) | .036 |
| High school | 10.7 (8.2 - 13.8) | 12.5 (10.2 - 15.2) | 9.8 (7.8 - 12.2) | 11.8 (9.5 - 14.6) | 12.0 (9.9 - 14.5) | 15.1 (12.1 - 18.6) | 14.8 (12.2 - 17.8) | 13.3 (10.4 - 16.8) | 13.9 (10.6 - 17.9) | 12.6 (10.1 - 15.6) | .058 |
| P, <HS vs HS | .007 | .003 | .003 | .075 | <.001 | .009 | .001 | <.001 | .069 | <.001 | |
| PIR^c | | | | | | | | | | | |
| <1.00 | 24.7 (18.4 - 32.4) | 22.0 (17.3 - 27.6) | 21.1 (16.1 - 27.2) | 24.2 (18.3 - 31.2) | 28.6 (22.6 - 35.5) | 24.7 (19.0 - 31.5) | 29.0 (23.2 - 35.6) | 24.0 (17.4 - 32.1) | 30.3 (23.1 - 38.7) | 23.7 (17.8 - 30.8) | .156 |
| 1.00 - 1.99 | 16.1 (12.7 - 20.4) | 20.0 (16.1 - 24.6) | 20.5 (16.4 - 25.3) | 18.0 (13.6 - 23.3) | 20.5 (16.1 - 25.7) | 21.9 (17.5 - 27.0) | 22.5 (18.0 - 27.7) | 22.6 (16.9 - 29.5) | 26.9 (21.0 - 33.7) | 28.6 (22.7 - 35.3) | <.001 |
| 2.00 - 3.99 | 9.5 (7.2 - 12.5) | 15.4 (11.9 - 19.6) | 11.5 (8.9 - 14.8) | 10.8 (8.3 - 13.9) | 13.6 (10.6 - 17.2) | 16.4 (12.9 - 20.6) | 15.5 (12.1 - 19.8) | 14.5 (10.2 - 20.0) | 13.4 (9.9 - 17.8) | 13.3 (9.9 - 17.5) | .094 |
| ≥4.00 | 6.3 (3.9 - 10.2) | 8.5 (5.8 - 12.2) | 5.0 (3.2 - 7.8) | 7.6 (5.3 - 10.8) | 5.5 (3.5 - 8.4) | 8.1 (5.5 - 11.8) | 8.9 (5.7 - 13.8) | 9.1 (4.7 - 16.9) | 4.8 (3.0 - 7.5) | 5.4 (3.7 - 7.8) | .381 |
| P, <1 vs ≥4 | <.001 | <.001 | <.001 | <.001 | <.001 | <.001 | <.001 | <.001 | <.001 | <.001 | |

CI = confidence interval; HS = high school; PIR, Poverty Income Ratio.

^an = 29 175. We categorized adults who reported experiencing any trouble seeing despite wearing contact lens or glasses as having visual impairment.

^bLinear trend.

^cPIR is an index of total income assessed in relation to federal poverty thresholds that vary by family size and composition. A PIR score of 1.00 is defined as the official federal poverty threshold level.

true disease burden among those with cataract who were undiagnosed or have not had their cataracts removed. By using the new clinical retinal imaging measures in the ophthalmology component of NHANES for AMD and DR, we were able to pick up individuals with AMD and DR who might otherwise not have known of their condition because of lack of access to and use of the health care system. Also, we estimated utilization among those with self-reported visual impairment, yet those who do not know they have impairment were missed.

Assessment of disease incidence is crucial to understanding the population effects of disease and the effectiveness of preventive efforts. However, national surveys rarely provide incidence information. NHANES and NHIS do not have more comprehensive examination or questionnaire data on eye diseases and use of eye care services because of the high cost attached to these surveys. In addition, because they are cross-sectional, causal inference cannot be obtained or inferred from national surveys. Finally, because of logistic constraints, national surveys often have only small sample sizes for certain ethnic groups (eg, Asian and Pacific Islanders and American Indians). National surveys frequently require that component protocols be targeted to specific diseases or conditions. Thus, a comprehensive picture of vision health with trends over time and across all ethnic groups within the United States is not currently feasible.

• **PUBLIC HEALTH IMPLICATIONS:** One of the core elements of a coordinated public health approach to improve the nation's vision health is implementing a national vision health surveillance and evaluation system.⁴¹ A national surveillance system assesses and monitors long-term outcomes of improved prevention and control of eye diseases and vision loss, increased access to eye care, better quality of life for the visually impaired, and enhanced vision health promotion throughout all life stages for all Americans. Surveillance data are used in many different contexts to estimate disease burden, recognize trends, detect gaps in coverage, and identify high-risk groups. Thereafter, strategies are developed and subsequently evaluated to effectively allocate limited health care resources

to reduce disparities. This is the model used for establishing and assessing progress on national vision health priorities in *Healthy People 2010* and *Healthy People 2020*.

National data (eg, NHANES and NHIS) have shown that vision health disparities exist. We need to first identify the underlying reasons for such disparities. Whenever possible, the efficacy, cost-effectiveness, and appropriate methodologies of screening should be established and implemented among the subpopulations most in need.⁴² Periodic, national point-prevalence data are needed with objective measurements of visual acuity and cause-specific disease. As technologies and treatments develop and demographics and the environment change, we need timely data and fast and efficient methods to detect new trends. In addition, people with eye disease likely have other health issues, particularly those of advancing age. A national surveillance system could ultimately improve the translation of vision research into a coordinated and personalized approach to monitor quality of care and advance public health prevention.

If vision data elements could be standardized and used across different surveys, more comprehensive analyses would be possible, and we would be better able to understand how to transfer knowledge about appropriate clinical care to the community to address disparities. For example, although an efficacious and cost-effective intervention is available for detecting diabetic retinopathy,^{43,44} challenges remain for getting those at risk screened and seen by a health care professional. There is also a need to assess whether there are socioeconomic gradients in the quality and outcome of eye care in the United States. An integrated surveillance system could identify and monitor the process of translating such public health interventions for communities to reduce existing visual health disparities. These efforts could help to improve overall health for all Americans.


In conclusion, there are vision health disparities among the major eye diseases and for vision health care in the United States. Disparities by race/ethnicity, education, and economic status show that the greatest need is for innovative interventions to reduce unnecessary vision loss among socioeconomically disadvantaged groups.

ALL AUTHORS HAVE COMPLETED AND SUBMITTED THE ICMJE FORM FOR DISCLOSURE OF POTENTIAL CONFLICTS OF INTEREST and none were reported. Publication of this article was supported by the National Center for Health Statistics (NCHS), Centers for Disease Control and Prevention (Atlanta, Georgia). Funding for the National Health and Nutrition Examination Survey (NHANES) ophthalmology component was provided by the Intra Agency Agreement 05FED47304 from the Division of Diabetes Translation, the Centers for Disease Control and Prevention. Funding for the vision component was provided by the National Eye Institute, National Institutes of Health, Intramural Research Program grant ZIAEY000402 (Bethesda, Maryland). The NCHS was involved in the design and conduct of the NHANES and NHIS study and in data collection. The Division of Diabetes Translation provided funding support for the ophthalmology component and was involved in the design and conduct of the study, in the collection, analysis, and interpretation of the data, and in the preparation, review, and approval of this article before submission. Involved in design (X.Z., M.F.C., A.R., J.B.S.) and conduct of the study (X.Z., M.F.C., A.R., S.A.P., J.B.S.); collection, management, analysis (X.Z., P.N., C.F.C.), and interpretation of the data (X.Z., M.F.C., A.R., S.A.P., P.N., C.F.C., L.S.G., L.E.B., A.F.E., J.E.C., J.B.S.); and preparation, review, or approval of the manuscript (X.Z., M.F.C., A.R., S.A.P., P.N., C.F.C., L.S.G., L.E.B., A.F.E., J.E.C., J.B.S.). Drs Zhang, Chou, and Nair had full access to all the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis. The NHANES and NHIS surveys are publicly available data and their protocols were approved by a human subjects review board, and written informed consent was obtained from all participants. The authors thank the NHANES and NHIS participants, without whom this study would not be possible. The findings and conclusions in this paper are those of the authors and do not necessarily represent the official position of the Centers for Disease Control and Prevention.

REFERENCES

- Saaddine JB, Narayan KM, Vinicor F. Vision loss: a public health problem? *Ophthalmology* 2003;110(2):253–254.
- World Health Organization. Global initiative for the elimination of avoidable blindness: Action plan 2006-2011: Available at: http://www.vision2020.org/documents/publications/Vision2020_report.pdf. Accessed: October 2, 2010.
- The Eye Diseases Prevalence Research Group. Prevalence of age-related macular degeneration in the United States. *Arch Ophthalmol* 2004;122(4):564–572.
- The Eye Diseases Prevalence Research Group. Prevalence of cataract and pseudophakia/aphakia among adults in the United States. *Arch Ophthalmol* 2004;122(4):487–494.
- The Eye Diseases Prevalence Research Group. Prevalence of open-angle glaucoma among adults in the United States. *Arch Ophthalmol* 2004;122(4):532–538.
- The Eye Diseases Prevalence Research Group. The prevalence of diabetic retinopathy among adults in the United States. *Arch Ophthalmol* 2004;122(4):552–563.
- The Eye Diseases Prevalence Research Group. The prevalence of refractive errors among adults in the United States, Western Europe, and Australia. *Arch Ophthalmol* 2004;122(4):495–505.
- The Eye Diseases Prevalence Research Group. Causes and prevalence of visual impairment among adults in the United States. *Arch Ophthalmol* 2004;122(4):477–485.
- Vu HT, Keeffe JE, McCarty CA, Taylor HR. Impact of unilateral and bilateral vision loss on quality of life. *Br J Ophthalmol* 2005;89(3):360–363.
- West SK, Munoz B, Rubin GS, et al. Function and visual impairment in a population-based study of older adults. The SEE project. Salisbury Eye Evaluation. *Invest Ophthalmol Vis Sci* 1997;38(1):72–82.
- Ivers RQ, Cumming RG, Mitchell P, Attebo K. Visual impairment and falls in older adults: the Blue Mountains Eye Study. *J Am Geriatr Soc* 1998;46(1):58–64.
- Ivers RQ, Norton R, Cumming RG, Butler M, Campbell AJ. Visual impairment and risk of hip fracture. *Am J Epidemiol* 2000;152(7):633–639.
- Bazargan M, Hamm-Baugh VP. The relationship between chronic illness and depression in a community of urban black elderly persons. *J Gerontol B Psychol Sci Soc Sci* 1995;50(2):S119–127.
- Rovner BW, Ganguli M. Depression and disability associated with impaired vision: the MoVies Project. *J Am Geriatr Soc* 1998;46(5):617–619.
- Fryback DG, Dasbach EJ, Klein R, et al. The Beaver Dam Health Outcomes Study: initial catalog of health-state quality factors. *Med Decis Making* 1993;13(2):89–102.
- Sharma S, Brown GC, Brown MM, et al. Converting visual acuity to utilities. *Can J Ophthalmol* 2000;35(5):267–272.
- Cugati S, Cumming RG, Smith W, Burlutsky G, Mitchell P, Wang JJ. Visual impairment, age-related macular degeneration, cataract, and long-term mortality: the Blue Mountains Eye Study. *Arch Ophthalmol* 2007;125(7):917–924.
- Klein R, Klein BE, Moss SE. Age-related eye disease and survival. The Beaver Dam Eye Study. *Arch Ophthalmol* 1995;113(3):333–339.
- Lee DJ, Gomez-Marin O, Lam BL, Zheng DD, Caban A. Visual impairment and morbidity in community-residing adults: the national health interview survey 1986-1996. *Ophthalmic Epidemiol* 2005;12(1):13–17.
- McCarty CA, Nanjan MB, Taylor HR. Vision impairment predicts 5 year mortality. *Br J Ophthalmol* 2001;85(3):322–326.
- Department of Health and Human Services. Healthy People 2010: Understanding and Improving Health. 2nd ed. Washington, DC: U.S. Government Printing Office; 2000.
- The Office of Disease Prevention and Health Promotion. Healthy People 2020: US Department of Health and Human Services. Available at: <http://www.healthypeople.gov/2020/default.aspx>. Accessed: January 25, 2011.
- Primo SA Jr, Wilson R, Hunt JW Jr, et al. Reducing visual health disparities in at-risk community health center populations. *J Public Health Manag Pract* 2009;15(6):529–534.
- Zhang X, Saaddine JB, Lee PP, et al. Eye care in the United States: do we deliver to high-risk people who can benefit most from it? *Arch Ophthalmol* 2007;125(3):411–418.
- Terry AL, Paulose-Ram R, Tilert TJ, et al. The methodology of visual field testing with frequency doubling technology in the National Health and Nutrition Examination Survey, 2005-2006. *Ophthalmic Epidemiol* 2010;17(6):411–421.
- Zhang X, Saaddine JB, Chou CF, et al. Prevalence of diabetic retinopathy in the United States, 2005-2008. *JAMA* 2010;304(6):649–656.
- Klein R, Chou CF, Klein BE, Zhang X, Meuer SM, Saaddine JB. Prevalence of age-related macular degeneration in the US population. *Arch Ophthalmol* 2011;129(1):75–80.
- U.S. Preventive Services Task Force. Screening for impaired visual acuity in older adults: U.S. Preventive Services Task Force recommendation statement. *Ann Intern Med* 2009;151(1):37–43, W10.
- U.S. Census Bureau. Population Division. Fertility & Family Statistics Branch. Current Population Survey: definitions and explanations. Available at: <http://www.census.gov/population/www/cps/cpsdef.html>. Accessed: September 8, 2010.
- Korn EL, Graubard BI. Analysis of Health Surveys. New York: Wiley; 1999.
- Klein RJ, Schoenborn CA. Age adjustment using the 2000 projected U.S. population. *Healthy People 2010. Stat Notes* 2001(20):1-10.
- Harris MI, Klein R, Cowie CC, Rowland M, Byrd-Holt DD. Is the risk of diabetic retinopathy greater in non-Hispanic blacks and Mexican Americans than in non-Hispanic whites with type 2 diabetes? A U.S. population study. *Diabetes Care* 1998;21(8):1230–1235.
- Tielsch JM, Sommer A, Katz J, Royall RM, Quigley HA, Javitt J. Racial variations in the prevalence of primary open-angle glaucoma. The Baltimore Eye Survey. *JAMA* 1991;266(3):369–374.
- West SK, Munoz B, Schein OD, Duncan DD, Rubin GS. Racial differences in lens opacities: the Salisbury Eye Evaluation (SEE) project. *Am J Epidemiol* 1998;148(11):1033–1039.
- Zhang X, Andersen R, Saaddine JB, Beckles GL, Duenas MR, Lee PP. Measuring access to eye care: a public health perspective. *Ophthalmic Epidemiol* 2008;15(6):418–425.
- Unzueta M, Globe D, Wu J, Paz S, Azen S, Varma R. Compliance with recommendations for follow-up care in Latinos: the Los Angeles Latino Eye Study. *Ethn Dis* 2004;14(2):285–291.

37. Orr P, Barron Y, Schein OD, Rubin GS, West SK. Eye care utilization by older Americans: the SEE Project. Salisbury Eye Evaluation. *Ophthalmology* 1999;106(5):904-909.
38. Ellwein LB, Urato CJ. Use of eye care and associated charges among the Medicare population: 1991-1998. *Arch Ophthalmol* 2002;120(6):804-811.
39. Bailey RN, Indian RW, Zhang X, Geiss LS, Duenas MR, Saaddine JB. Visual impairment and eye care among older adults - five States, 2005. *MMWR Morb Mortal Wkly Rep* 2006;55(49):1321-1325.
40. Mitchell P, Hayes P, Wang JJ. Visual impairment in nursing home residents: the Blue Mountains Eye Study. *Med J Aust* 1997;166(2):73-76.
41. Centers for Disease Control and Prevention. Vision Health Initiative. Improving the Nation's Vision Health: A Coordinated Public Health Approach. 2006. Available at: http://www.cdc.gov/visionhealth/pdf/improving_nations_vision_health.pdf. Accessed: February 1, 2010.
42. Baker RS, Bazargan M, Bazargan-Hejazi S, Calderon JL. Access to vision care in an urban low-income multiethnic population. *Ophthalmic Epidemiol* 2005;12(1):1-12.
43. Javitt JC, Aiello LP. Cost-effectiveness of detecting and treating diabetic retinopathy. *Ann Intern Med* 1996;124(1 Pt 2):164-169.
44. Singer DE, Nathan DM, Fogel HA, Schachat AP. Screening for diabetic retinopathy. *Ann Intern Med* 1992;116(8):660-671.



Biosketch

Xinzhi Zhang, MD, PhD, is an epidemiologist at the Centers for Disease Control and Prevention (CDC), Atlanta, Georgia. He received his medical degree from Peking Union Medical College and PhD in Health Services Administration from the University of Alabama at Birmingham (UAB), where he is also an adjunct professor. He has received awards from Agency for Healthcare Research and Quality, CDC, and American Public Health Association. Dr Zhang is an overseas fellow of the Royal Society of Medicine.