

Making Health Care Safer

Antibiotic Rx in Hospitals: Proceed with Caution



1 in 2

More than half of all hospital patients receive an antibiotic.



3x

Doctors in some hospitals prescribed 3 times as many antibiotics as doctors in other hospitals.



30%

Reducing the use of high-risk antibiotics by 30% can lower deadly diarrhea infections by 26%.

Antibiotics save lives, but poor prescribing practices are putting patients at unnecessary risk for preventable allergic reactions, super-resistant infections, and deadly diarrhea. Errors in prescribing decisions also contribute to antibiotic resistance, making these drugs less likely to work in the future.

To protect patients and preserve the power of antibiotics, hospital CEOs/medical officers can:


- ◇ Adopt an antibiotic stewardship program that includes, at a minimum, this checklist:
 1. **Leadership commitment:** Dedicate necessary human, financial, and IT resources.
 2. **Accountability:** Appoint a single leader responsible for program outcomes. Physicians have proven successful in this role.
 3. **Drug expertise:** Appoint a single pharmacist leader to support improved prescribing.
 4. **Act:** Take at least one prescribing improvement action, such as requiring reassessment within 48 hours, to check drug choice, dose, and duration.
 5. **Track:** Monitor prescribing and antibiotic resistance patterns.
 6. **Report:** Regularly report to staff prescribing and resistance patterns, and steps to improve.
 7. **Educate:** Offer education about antibiotic resistance and improving prescribing practices.
- ◇ Work with other health care facilities to prevent infections, transmission, and resistance.

→ See page 4

Want to learn more? Visit

www

www.cdc.gov/vitalsigns



Problem – Poor antibiotic prescribing harms patients

Antibiotic prescribing practices vary widely and errors are common.

- ◇ About half of patients receive an antibiotic for at least one day during the course of an average hospital stay.
- ◇ The most common types of infections for which hospital clinicians wrote antibiotic prescriptions were lung infections (22%), urinary tract infections (14%), and suspected infections caused by drug-resistant *Staphylococcus* bacteria, such as MRSA (17%).
- ◇ About 1 out of 3 times, prescribing practices to treat urinary tract infections and prescriptions for the critical and common drug vancomycin included a potential error – given without proper testing or evaluation, or given for too long.
- ◇ Doctors in some hospitals prescribed up to 3 times as many antibiotics as doctors in similar areas of other hospitals. This difference suggests the need to improve prescribing practices.

Poor prescribing puts patients at risk.

- ◇ Although antibiotics save lives (for example, in the prompt treatment of sepsis, a life-threatening infection throughout the body), they can also put patients at risk for a *Clostridium difficile* infection, deadly diarrhea that causes at least 250,000 infections and 14,000 deaths each year in hospitalized patients.
- ◇ Decreasing the use of antibiotics that most often lead to *C. difficile* infection by 30% (this is 5% of overall antibiotic use) could lead to 26% fewer of these deadly diarrheal infections. These antibiotics include fluoroquinolones, β -lactams with β -lactamase inhibitors, and extended-spectrum cephalosporins.
- ◇ Patients getting powerful antibiotics that treat a broad range of infections are up to 3 times more likely to get another infection from an even more resistant germ.

Every time antibiotics are prescribed:



Specific recommendations for common prescribing situations:



1. Order recommended cultures before antibiotics are given and start drugs promptly.



2. Make sure indication, dose, and expected duration are specified in the patient record.



3. Reassess within 48 hours and adjust Rx if necessary or stop Rx if indicated.



Rx for urinary tract infections

- Make sure that culture results represent true infection and not just colonization.
 - Assess patient for signs and symptoms of UTI.
 - Make sure that urinalysis is obtained with every urine culture.
- Treat for recommended length of time and ensure that planned post-discharge treatment takes into account the antibiotics given in the hospital.



Rx for pneumonia

- Make sure that symptoms truly represent pneumonia and not an alternate, non-infectious diagnosis.
- Treat for the recommended length of time and ensure that planned post-discharge treatment takes into account the antibiotics given in the hospital.




Rx for MRSA infections

- Verify that MRSA is growing in clinically relevant cultures. Do not use vancomycin to treat infections caused by methicillin-susceptible staph (and not MRSA).

Improving antibiotic prescribing in hospitals

Key moments for improving the cycle of antibiotic prescribing practices



What Can Be Done



The Federal government is

- ◇ Expanding the National Healthcare Safety Network to help hospitals track antibiotic use and resistance.
- ◇ Sharing prescribing improvement recommendations and tools with clinicians and administrators.
www.cdc.gov/getsmart/healthcare
- ◇ Supporting networks testing new prescribing improvement strategies.
- ◇ Helping hospitals and health departments create regional programs to improve antibiotic prescribing.
- ◇ Improving health care for veterans by launching antibiotic stewardship programs in Veteran's Health Administration hospitals.
- ◇ Providing incentives for development of new antibiotics.



State and local health departments can

- ◇ Gain an understanding of antibiotic stewardship activities in the state or area.
- ◇ Facilitate efforts to improve antibiotic prescribing and prevent antibiotic resistance.
- ◇ Provide educational tools to facilities to help prescribers improve practices.



Hospital CEOs/medical officers can

- ◇ Adopt an antibiotic stewardship program that includes, at a minimum, this checklist:
 - 1. Leadership commitment:** Dedicate necessary human, financial, and IT resources.
 - 2. Accountability:** Appoint a single leader responsible for program outcomes. Physicians have proven successful in this role.
 - 3. Drug expertise:** Appoint a single pharmacist leader to support improved prescribing.
 - 4. Act:** Take at least one prescribing improvement action, such as requiring reassessment within 48 hours, to check drug choice, dose, and duration.

- 5. Track:** Monitor prescribing and antibiotic resistance patterns.
 - 6. Report:** Regularly report to staff prescribing and resistance patterns, and steps to improve.
 - 7. Educate:** Offer education about antibiotic resistance and improving prescribing practices.
- ◇ Work with other health care facilities to prevent infections, transmission, and resistance.



Doctors and other hospital staff can

- ◇ Prescribe antibiotics correctly – get cultures, start the right drug promptly at the right dose for the right duration. Reassess the prescription within 48 hours based on tests and patient exam.
- ◇ Document the dose, duration and indication for every antibiotic prescription.
- ◇ Stay aware of antibiotic resistance patterns in your facility.
- ◇ Participate in and lead efforts within your hospital to improve prescribing practices.
- ◇ Follow hand hygiene and other infection control measures with every patient.



Hospital patients can

- ◇ Ask if tests will be done to make sure the right antibiotic is prescribed.
- ◇ Be sure everyone cleans their hands before touching you. If you have a catheter, ask each day if it is necessary.

For more information, please contact
Telephone: 1-800-CDC-INFO (232-4636)
TTY: 1-888-232-6348
Web: www.cdc.gov
Centers for Disease Control and Prevention
1600 Clifton Road NE, Atlanta, GA 30333
Publication date: 3/4/2014

www.cdc.gov/vitalsigns

www.cdc.gov/mmwr