



West Nile virus disease cases and presumptive viremic blood donors reported to ArboNET, United States, 2012

State	Neuroinvasive disease cases*	Non-neuroinvasive disease cases	Total cases†	Deaths	Presumptive viremic blood donors‡
Alabama	38	24	62	2	10
Arizona	87	46	133	7	27
Arkansas	44	20	64	7	4
California	297	182	479	20	63
Colorado	62	69	131	5	12
Connecticut	12	9	21	0	1
Delaware	2	7	9	1	0
District of Columbia	8	2	10	1	0
Florida	52	21	73	3	7
Georgia	46	53	99	6	20
Idaho	5	12	17	0	3
Illinois	187	103	290	12	31
Indiana	46	31	77	8	18
Iowa	11	20	31	0	6
Kansas	20	36	56	3	23
Kentucky	13	10	23	3	3
Louisiana	155	180	335	16	34
Maine	1	0	1	0	0
Maryland	25	22	47	6	13
Massachusetts	25	8	33	1	3
Michigan	141	61	202	17	38
Minnesota	34	36	70	1	35
Mississippi	103	144	247	5	21
Missouri	17	3	20	3	8
Montana	1	5	6	1	0
Nebraska	42	151	193	4	36
Nevada	5	4	9	1	1
New Hampshire	1	0	1	0	0
New Jersey	22	26	48	6	4
New Mexico	24	23	47	1	6
New York	61	46	107	9	15
North Carolina	7	0	7	2	2
North Dakota	39	50	89	1	16
Ohio	76	45	121	7	18
Oklahoma	103	88	191	17	38
Oregon	0	11	11	0	0
Pennsylvania	33	27	60	4	6
Puerto Rico	1	0	1	0	0
Rhode Island	2	2	4	0	0
South Carolina	20	9	29	4	12
South Dakota	62	141	203	3	42
Tennessee	19	14	33	1	1
Texas	844	1,024	1,868	89	108
Utah	3	2	5	1	0
Vermont	1	2	3	0	0
Virginia	20	10	30	4	3
Washington	4	0	4	0	0
West Virginia	5	5	10	0	1
Wisconsin	44	13	57	4	13
Wyoming	3	4	7	0	1
Totals	2,873	2,801	5,674	286	703

*Includes cases reported as meningitis, encephalitis, or acute flaccid paralysis.

†Includes confirmed and probable cases.

‡Presumptive viremic blood donors (PVDs) are people who had no symptoms at the time of donating blood through a blood collection agency, but whose blood tested positive when screened for the presence of West Nile virus. Some PVDs develop symptoms after donation; of the 703 PVDs, 98 (14%) developed clinical illness and are also included as disease cases.