
## FROM THE NATIONAL CENTER FOR HEALTH STATISTICS

## Age-Adjusted Rate\* for Suicide,† by Sex — National Vital Statistics System, United States, 1975–2015



<sup>\*</sup> Age adjusted rates are suicide deaths per 100,000 standard population.

There was an overall decline of 24% in the age-adjusted suicide rate from 1977 (13.7 per 100,000) to 2000 (10.4). The rate increased in most years from 2000 to 2015. The 2015 suicide rate (13.3) was 28% higher than in 2000. The rates for males and females followed the overall pattern; however, the rate for males was approximately 3–5 times higher than the rate for females throughout the study period.

Source: CDC. National Vital Statistics System. Mortality data. https://www.cdc.gov/nchs/nvss/deaths.htm.

Reported by: Sally C. Curtin, MA, sac2@cdc.gov, 301-458-4142; Holly Hedegaard, MD; Margaret Warner, PhD.

 $For more information on this topic, CDC \ recommends \ the following \ link: https://www.cdc.gov/violenceprevention/suicide/index.html.$ 

<sup>&</sup>lt;sup>†</sup> Suicides are identified using *International Classification of Diseases (ICD), 8th Revision* codes E950–E959 for 1975–1978; *ICD, 9th Revision* codes E950–E959 for 1979–1998; and *ICD, 10th Revision* codes U03, X60–X84, and Y87.0 for 1999–2015.